

RESEARCH

Open Access

Bounded solutions for the nonlinear wave equation

Tacksun Jung¹ and Q-Heung Choi^{2*}

*Correspondence:

qheung@inha.ac.kr

²Department of Mathematics
Education, Inha University, Incheon,
402-751, Korea

Full list of author information is
available at the end of the article

Abstract

We investigate the number of periodic weak solutions for the wave equation with nonlinearity decaying at the origin. We get a theorem which shows the existence of a bounded weak solution for this problem. We obtain this result by approaching the variational method and applying the critical point theory for the indefinite functional induced from the invariant subspaces and the invariant functional.

MSC: 35L05; 35L70

Keywords: wave equation; critical point theory; invariant function; invariant subspace; $(P.S.)_c$ condition; eigenvalue problem

1 Introduction and statement of the main result

Let $g(x, t, u)$ be a C^1 function from $[0, \pi] \times R \times R$ to R and T -periodic in t . In this paper we are concerned with the number of weak periodic solutions of the following wave equation with boundary and periodic conditions:

$$\begin{aligned}u_{tt} - u_{xx} &= g(x, t, u), \\u(0, t) &= u(\pi, t) = 0, \\u(x, t + T) &= u(x, t), \\u_t(x, 0) &= u_t(x, T) \quad \forall x \in [0, \pi],\end{aligned}\tag{1.1}$$

where T is a rational multiple of π . We assume that g satisfies the following conditions:

(g1) $g \in C^1([0, \pi] \times R \times R, R)$ is T -periodic in t ,

(g2) $g(x, t, 0) = 0$, $g(x, t, \xi) = o(|\xi|)$ uniformly with respect to $x \in [0, \pi]$ and $t \in R$,

(g3) there exists $C > 0$ such that $|g(x, t, \xi)| < C \forall x \in [0, \pi], t \in R, \xi \in R$.

Nonlinear problem of this type has been considered by many authors (cf. [1–5]).

The purpose of this paper is to show the existence of T -periodic weak solutions of problem (1.1).

Our main result is as follows.

Theorem 1.1 *Assume that g satisfies (g1)–(g3). Then problem (1.1) has at least one bounded solution.*

For the proof of our main result, we approach the variational method and apply the critical point theory induced from the invariant subspaces and the invariant functional.

The outline of the proof of Theorem 1.1 is as follows. In Section 2, we introduce two Banach spaces H and E of functions satisfying some symmetry properties, stable by A ($Au = u_{tt} - u_{xx}$), g such that the intersection of H with the kernel of A is reduced to 0. The search of a solution of problem (1.1) in the space H reduces the problem to a situation where A^{-1} is a compact operator. In Section 3, we introduce a functional I defined on E whose critical points and weak solutions of (1.1) possess one-to-one correspondence. We prove that $I \in C^1(E, R)$ and satisfies the Palais-Smale condition. By a critical point theorem for indefinite functionals (cf. [6]), we prove that there exists at least one solution of (1.1) which is bounded, and we prove Theorem 1.1.

2 Invariant Banach space

Let $\Omega = (0, \pi) \times (0, T)$; T is a rational multiple of π

$$T = \frac{2\pi b}{a},$$

where a and b are coprime integers. Let \mathcal{A} be the operator defined by

$$\begin{aligned} D(\mathcal{A}) = \{ & u \in C^2(\overline{\Omega}) \mid u(0, t) = u(\pi, t) = 0 \quad \forall t \in [0, T], \\ & u(x, 0) = u(x, T) \quad \forall x \in [0, \pi], \\ & u_t(x, 0) = u_t(x, T) \quad \forall x \in [0, \pi] \}, \\ \mathcal{A}u = & u_{tt} - u_{xx}. \end{aligned}$$

Let A be the adjoint of \mathcal{A} in $L^2(\Omega)$. We investigate the weak solutions of

$$Au = g(x, t, u).$$

We note that the eigenvalues of A are $\lambda_{jk} = j^2 - (\frac{2\pi k}{T})^2$, $j = 1, 2, \dots$ and $k = 0, 1, 2, \dots$, and the corresponding eigenfunctions are

$$\sin jx \sin \frac{2\pi kt}{T} \quad \text{and} \quad \sin jx \cos \frac{2\pi kt}{T}.$$

We also note that the set of functions $\sin jx \sin \frac{2\pi kt}{T}$, $\sin jx \cos \frac{2\pi kt}{T}$ is an orthogonal base for $L^2(\Omega)$. Let u be a function of $L^2(\Omega)$. Then there exists one and only one function of $L^2([0, \pi] \times R)$ which is T -periodic in t and equals u on Ω . We shall denote this function by u . Let us denote an element u , in $L^2(\Omega)$, by

$$u = \sum_{j>0, k} u_{jk} \sin jx \exp ik \frac{a}{b} t$$

with

$$u_{j,k} = \bar{u}_{j,-k}.$$

We assume that b is even and a is odd. Let H be the closed subspace of $L^2(\Omega)$ defined by

$$H = \left\{ u \in L^2(\Omega) \mid u(x, t) = -u\left(x, t + \frac{T}{2}\right) \text{ a.e. } x \in (0, \pi), t \in R \right\}.$$

Then H is invariant under shift: Let $h \in H$ and τ be a real number. If $v(x, t) = u(x, t + \tau)$, then $v \in H$. H is invariant under g : Let $u \in H$ such that $g(u) \in L^2(\Omega)$. Then $g(u) \in H$. Let $\tilde{u}(x, t) = u(x, t + \frac{T}{2})$. Then

$$u = \sum_{j>0,k} u_{j,k} (-1)^k \sin jx \exp ik \frac{a}{b} t.$$

Therefore

$$u \in H \iff u_{j,k} = 0 \text{ for any even } k. \tag{2.1}$$

Let A_1 be a linear operator of H defined by

$$D(A_1) = D(A) \cap H,$$

$$A_1 u = Au.$$

Then A_1 is self-adjoint in H and $H \cap N(A) = \{0\}$, where $N(A)$ is the kernel of A . In fact, let $u \in H \cap N(A)$. Then

$$u = \sum u_{j,k} \sin jx \exp ik \frac{a}{b} t,$$

$$j^2 - \frac{k^2 a^2}{b^2} \neq 0 \implies u_{j,k} = 0.$$

Let j and k be such that

$$j^2 - \frac{k^2 a^2}{b^2} = 0.$$

Since b is even and a is odd, k is even. Using (2.1), we have $u_{j,k} = 0$, and therefore $H \cap N(A) = \{0\}$. The eigenvalues of A_1 are $j^2 - (\frac{2\pi k}{T})^2$, where j is odd and k is even and $H \cap N(A) = \{0\}$. Given $u \in H$, we write

$$u = \sum_{j>0} u_{j,k} \sin jx \exp i \frac{2\pi kt}{T}, \quad j \text{ is odd, } k \text{ is even}$$

with

$$u_{j,k} = \bar{v}_{j,-k}.$$

Let

$$E = \left\{ u \in H \mid \sum_{j,k} \left| j^2 - \frac{a^2 k^2}{b^2} \right| \cdot |u_{j,k}|^2 < +\infty \right\},$$

$$(u, v) = \sum_{j,k} \left| j^2 - \frac{a^2 k^2}{b^2} \right| u_{j,k} \cdot \bar{v}_{j,k} \quad \text{for } u, v \in E,$$

where (\cdot, \cdot) is a scalar product on E . With this scalar product, E is a Hilbert space with a norm

$$\|u\| = (u, u)^{\frac{1}{2}}, \quad u \in E.$$

Let

$$\|u\|_r = \left(\int_{\Omega} |u|^r \right)^{\frac{1}{r}}, \quad r \geq 1.$$

By the classical theorem of Riesz (cf. [7, p.525]), we have

$$\|u\|_r \leq \left(\frac{\pi T'}{2} \right)^{\frac{1}{r}} \left(\sum_{j,k} |u_{j,k}|^{r'} \right)^{\frac{1}{r'}}, \quad r \geq 2, \frac{1}{r} + \frac{1}{r'} = 1.$$

Since for every $\epsilon > 0$

$$\sum_{j \text{ odd}, k \text{ even}} \frac{1}{|j^2 - \frac{a^2 k^2}{b^2}|^{1+\epsilon}} < \infty,$$

it follows that for every $r \in [2, +\infty)$, there is $c_r \in \mathbb{R}$ such that

$$\|u\|_r \leq c_r \|u\|. \tag{2.2}$$

Let

$$E_+ = \left\{ u \mid u \in E, u_{j,k} = 0 \text{ if } j^2 - \frac{a^2 k^2}{b^2} < 0 \right\},$$

$$E_- = \left\{ u \mid u \in E, u_{j,k} = 0 \text{ if } j^2 - \frac{a^2 k^2}{b^2} > 0 \right\}.$$

Then $E = E_+ \oplus E_-$. Let P_+ be the orthogonal projection from E onto E_+ and P_- be the orthogonal projection from E onto E_- . We can write $u = P_+ u + P_- u$ for $u \in E$.

3 Critical point theorem and the proof of Theorem 1.1

Now, we are going to seek a function u in E such that

$$A_1 u = g(x, t, u). \tag{3.1}$$

We consider the associated functional of (3.1),

$$I(u) = \frac{1}{2} \int_{\Omega} [-|u_t|^2 + |u_x|^2] dx dt - \int_{\Omega} G(x, t, u) dx dt$$

$$= \frac{1}{2} (\|P_+ u\|^2 - \|P_- u\|^2) - \int_{\Omega} G(x, t, u) dx dt, \tag{3.2}$$

where

$$G(x, t, u) = \int_0^u g(x, t, s) ds.$$

By (g1), I is well defined.

We recall the critical point theorem for the indefinite functional (cf. [6]).

Let

$$B_r = \{u \in H \mid \|u\| \leq r\}, \quad \partial B_r = \{u \in H \mid \|u\| = r\}.$$

Theorem 3.1 (Critical point theorem for the indefinite functional) *Let E be a real Hilbert space with $E = E_1 \oplus E_2$ and $E_2 = E_1^\perp$. Suppose that $I \in C^1(E, R)$ satisfies (PS), and that*

(I1) $I(u) = \frac{1}{2}(Lu, u) + bu$, where $Lu = L_1P_1u + L_2P_2u$ and $L_i : E_i \rightarrow E_i$ is bounded and self-adjoint, $i = 1, 2$,

(I2) b' is compact, and

(I3) there exist a subspace $\tilde{E} \subset E$ and sets $S \subset E$, $Q \subset \tilde{E}$ and constants $\alpha > \omega$ such that

(i) $S \subset E_1$ and $I|_S \geq \alpha$,

(ii) Q is bounded and $I|_{\partial Q} \leq \omega$,

(iii) S and ∂Q link.

Then I possesses a critical value $c \geq \alpha$.

The eigenvalues of A_1 are $\lambda_{jk} = j^2 - (\frac{2\pi k}{T})^2$, where j is odd and k is even, $j = 1, 2, \dots$ and $k = 0, 1, 2, \dots$. Thus A_1^{-1} is a compact operator.

It is convenient for the following to rearrange the eigenvalues λ_{jk} , where j is odd and k is even, by increasing magnitude: from now on we denote by $(\eta_i^-)_{i \geq 1} = \lambda_{jk} < 0$ the sequence of negative eigenvalues, by $(\eta_i^+)_{i \geq 1} = \lambda_{jk} > 0$ the sequence of positive ones, so that

$$\dots \leq \eta_i^- \leq \dots \leq \eta_2^- \leq \eta_1^- < \eta_1^+ \leq \eta_2^+ \leq \dots \leq \eta_i^+ \leq \dots \tag{3.3}$$

We note that each eigenvalue has a finite multiplicity and that $\eta_i^- \rightarrow -\infty$, $\eta_i^+ \rightarrow +\infty$ as $i \rightarrow \infty$.

We shall show that the functional I satisfies the geometrical assumptions of the critical point theorem for the indefinite functional.

By the following lemma, $I(u) \in C^1(E, R)$ and the solutions of (3.1) coincide with the nonzero critical points of $I(u)$.

Lemma 3.1 *Assume that g satisfies (g1)-(g3). Then $I(u)$ is continuous and Fréchet differentiable in E with the Fréchet derivative*

$$\begin{aligned} I'(u)h &= \int_{\Omega} [-u_t \cdot h_t + u_x \cdot h_x - g(x, t, u)h] \, dx \, dt \\ &= (P_+u, P_+h) - (P_-u, P_-h) - \int_{\Omega} g(x, t, u)h \, dx \, dt \end{aligned}$$

for all $h \in E$. Moreover, if we set

$$F(u) = \int_{\Omega} G(x, t, u) \, dx \, dt,$$

then $F'(u)$ is continuous with respect to weak convergence, $F'(u)$ is compact, and

$$F'(u)h = \int_{\Omega} g(x, t, u)h \, dx \, dt \quad \text{for all } h \in E.$$

This implies that $I \in C^1(E, R)$ and $F(u)$ is weakly continuous.

For the proof of Lemma 3.1, refer to [6].

Lemma 3.2 *Assume that g satisfies (g1)-(g3). The problem*

$$\begin{aligned}
 &u_{tt} - u_{xx} \quad \text{in } E, \\
 &u(0, t) = u(\pi, t) = 0, \\
 &u(x, t + T) = u(x, t), \\
 &u_t(x, 0) = u_t(x, T) \quad \forall x \in [0, \pi]
 \end{aligned} \tag{3.4}$$

has only a trivial solution.

Proof Let $Au = u_{tt} - u_{xx}$. Since $H \cap N(A) = \{0\}$, $E \cap N(A) = \{0\}$. Thus (3.4) has only a trivial solution. \square

Lemma 3.3 *Assume that g satisfies (g1)-(g3). Then $I(u)$ satisfies the Palais-Smale condition: If for a sequence (u_m) , $I(u_m)$ is bounded from above and $I'(u_m) \rightarrow 0$ as $m \rightarrow \infty$, then (u_m) has a convergent subsequence.*

Proof Let (u_m) be a sequence with $I(u_m) \leq M$ and $I'(u_m) \rightarrow 0$ as $m \rightarrow \infty$. It suffices to show that (u_m) is bounded. By contradiction, we suppose that $\|u_m\| \rightarrow \infty$ as $m \rightarrow \infty$. Let $w_m = \frac{u_m}{\|u_m\|}$. Then $\|w_m\| = 1$ and w_m converges weakly to an element, say w . Then we have

$$\frac{M}{\|u_m\|^2} \leq \frac{I(u_m)}{\|u_m\|^2} = \frac{1}{2} \int_{\Omega} [-|(w_m)_t|^2 + |(w_m)_{xx}|^2] - \int_{\Omega} \frac{G(x, t, u_m)}{\|u_m\|^2}. \tag{3.5}$$

Since $G(x, t, u)$ is bounded, letting $m \rightarrow \infty$ in (3.5), we have

$$0 \leq \lim P_+ w_m - \lim P_- w_m = \int_{\Omega} [-|w_t|^2 + |w_{xx}|^2]. \tag{3.6}$$

On the other hand, we have

$$0 \leftarrow \frac{I'(u_m)}{\|u_m\|^2} = \int_{\Omega} [-|(w_m)_t|^2 + |(w_m)_{xx}|^2] - \int_{\Omega} \frac{g(x, t, u_m)}{\|u_m\|^2}. \tag{3.7}$$

Letting $m \rightarrow \infty$ in (3.7), we have

$$0 = \lim_{m \rightarrow \infty} \|P_+ w_m\|^2 - \lim_{m \rightarrow \infty} \|P_- w_m\|^2 = \int_{\Omega} [-|w_t|^2 + |w_{xx}|^2] = \|P_+ w\|^2 - \|P_- w\|^2. \tag{3.8}$$

By (3.6) and (3.8), $\lim_{m \rightarrow \infty} \|P_+ w_m\|^2 = \|P_+ w\|^2$ and $\lim_{m \rightarrow \infty} \|P_- w_m\|^2 = \|P_- w\|^2$. Thus $\lim_{m \rightarrow \infty} \|w_m\| = \|w\|$, so w_m converges strongly to w and by (3.8), w is a weak solution of the problem

$$w_{tt} - w_{xx} = 0 \quad \text{in } E.$$

By Lemma 3.2, $w = 0$, which is absurd for the fact that $\|w\| = 1$. Thus (u_m) is bounded. \square

Let

$$Q = (\bar{B}_R \cap E_-) \oplus \{re | e \in \partial B_1 \cap E_+, 0 < r < R\}.$$

Lemma 3.4 *Assume that g satisfies the conditions (g1)-(g3). There exist a subspace $\tilde{E} \subset E$ and sets $\partial B_\rho \subset E$, $Q \subset \tilde{E}$ and constants $\rho > 0$ and $\alpha > 0$ such that*

- (i) $\partial B_\rho \subset E_+$ and $I|_{\partial B_\rho} \geq \alpha$,
- (ii) there are $e \in \partial B_1 \cap E_+$ and $R > \rho$ such that if $Q = (\bar{B}_R \cap E_-) \oplus \{re | 0 < r < R\}$, then $I|_{\partial Q} \leq 0$,
- (iii) there exists $u_0 \in E \setminus Q$ such that $\|u_0\| > R$ and $I(u_0) \leq 0$,
- (iv) ∂B_ρ and ∂Q link.

Proof (i) Let $u \in E_+$. Since $G(x, t, u)$ is bounded, there exists a constant $C > 0$ such that

$$\begin{aligned} I(u) &= \frac{1}{2} \|P_+ u\|^2 - \frac{1}{2} \|P_- u\|^2 - \int_{\Omega} G(x, t, u) \\ &\geq \frac{1}{2} \|P_+ u\|^2 - C \end{aligned}$$

for $C > 0$. Then there exist a constant $\rho > 0$ and a constant $\alpha > 0$ such that if $u \in \partial B_\rho \cap E_+$, then $I(u) \geq \alpha$.

(ii) Let us choose an element $e \in B_1 \cap E_+$. Let $u \in (\bar{B}_r \cap E_-) \oplus \{re | 0 < r\}$. Then $u = v + w$, $v \in B_r \cap E_-$, $w = re$. We note that

$$\text{if } v \in B_r \cap E_-, \text{ then } \int_{\Omega} [-|v_t|^2 + |v_x|^2] dx dt \leq \eta_1^- \|v\|_{L^2(\Omega)}^2 < 0.$$

Thus we have

$$\begin{aligned} I(u) &= \frac{1}{2} r^2 - \frac{1}{2} \|v\|^2 - \int_{\Omega} G(x, t, v + re) \\ &\leq \frac{1}{2} r^2 + \frac{1}{2} \eta_1^- \|v\|_{L^2(\Omega)}^2 + C_1 \end{aligned}$$

for $C_1 > 0$. Then there exists $R > \rho > 0$ such that if $u \in Q = (\bar{B}_R \cap E_-) \oplus \{re | 0 < r < R\}$, then $I(u)|_{\partial Q} \leq 0$.

(iii) Let us choose $u_0 \in E \setminus Q$. By (ii), $I(u_0) \leq 0$.

(iv) S and Q link at the set $\{e\}$. □

Proof of Theorem 1.1 By Lemma 3.1, $I(u)$ is continuous and Fréchet differentiable in E . By Lemma 3.3, the functional I satisfies the (PS) condition. We note that $I(0) = 0$. By Lemma 3.4, there are constants $\rho > 0$, $\alpha > 0$ and a bounded neighborhood B_ρ of 0 such that $I|_{\partial B_\rho \cap E_+} \geq \alpha$, and there are $e \in \partial B_1 \cap E_+$ and $R > \rho$ such that if $Q = (\bar{B}_R \cap E_-) \oplus \{re | 0 < r < R\}$. Let us set

$$\Gamma = \{\gamma \in C(\bar{Q}, H) | \gamma = \text{id on } \partial Q\}.$$

By Theorem 3.1, I possesses a critical value $c \geq \alpha$. Moreover, c can be characterized as

$$c = \inf_{\gamma \in \Gamma} \max_{u \in Q} I(\gamma(u)).$$

Thus we prove that I has at least one nontrivial critical point, we denote by \tilde{u} a critical point of I such that $I(\tilde{u}) = c$. We claim that c is bounded. In fact, we have

$$c \leq \max_{0 \leq \tau \leq 1} I(\tau u_0),$$

and by (g3),

$$\begin{aligned} I(\tau u_0) &= \tau^2 \left(\frac{1}{2} \|P_+ u_0\|^2 - \frac{1}{2} \|P_- u_0\|^2 \right) - \int_{\Omega} G(x, t, \tau u_0) \, dx \, dt \\ &\leq \tau^2 \|u_0\|^2 - \int_{\Omega} G(x, t, \tau u_0) \, dx \, dt \\ &\leq C_1 \tau^2 \|u_0\|^2 + C_1 \end{aligned}$$

for some constant $C_1 > 0$. Since $0 \leq \tau \leq 1$, c is bounded:

$$c < \tilde{C}. \tag{3.9}$$

We claim that \tilde{u} is bounded. In fact, by contradiction, $\tilde{u}_{tt} + \tilde{u}_{xx} = g(x, t, \tilde{u})$ and, for any $K > 0$, $\max_{\Omega} |\tilde{u}(x, t)| > K$ imply that

$$c = I(\tilde{u}) = \frac{1}{2} (\|P_+ \tilde{u}\|^2 - \|P_- \tilde{u}\|^2) - \int_{\Omega} G(x, t, \tilde{u}) \, dx \, dt$$

is not bounded, which is absurd to the fact that $c = I(\tilde{u})$ is bounded. Thus \tilde{u} is bounded, so (1.1) has at least one bounded weak solution, and hence we prove Theorem 1.1. \square

Competing interests

The authors did not provide this information.

Authors' contributions

The authors did not provide this information.

Author details

¹Department of Mathematics, Kunsan National University, Kunsan, 573-701, Korea. ²Department of Mathematics Education, Inha University, Incheon, 402-751, Korea.

Acknowledgements

This work (Choi) was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (KRF-2013010343).

Received: 27 June 2013 Accepted: 25 October 2013 Published: 25 Nov 2013

References

1. Amann, H, Zehnder, E: Multiple periodic solutions for a class of nonlinear-autonomous wave equations. *Houst. J. Math.* **7**, 147-174 (1981)
2. Brezis, H, Coron, JM: Periodic solutions of nonlinear wave equations and Hamiltonian systems. *Am. J. Math.* **103**, 559-570 (1981)
3. Brezis, H, Nirenberg, L: Forced vibrations for a nonlinear wave equation. *Commun. Pure Appl. Math.* **31**, 1-30 (1978)
4. Choi, QH, Jung, T: An application of a variational reduction method to a nonlinear wave equation. *J. Differ. Equ.* **7**, 390-410 (1995)
5. Rabinowitz, PH: Free vibrations for a semilinear wave equation. *Commun. Pure Appl. Math.* **31**, 31-68 (1968)
6. Benci, V, Rabinowitz, PH: Critical point theorems for indefinite functionals. *Invent. Math.* **52**, 241-273 (1979)
7. Fadell, ER, Rabinowitz, PH: Generalized cohomological index theories for the group actions with an application to bifurcation questions for Hamiltonian systems. *Invent. Math.* **45**, 139-174 (1978)

10.1186/1687-2770-2013-257

Cite this article as: Jung and Choi: Bounded solutions for the nonlinear wave equation. *Boundary Value Problems* 2013, **2013**:257