# RESEARCH

# **Open Access**



# The existence of solutions for *p*-Laplacian boundary value problems at resonance on the half-line

Weihua Jiang<sup>\*</sup>, Yi Zhang and Jiqing Qiu

\*Correspondence: weihuajiang@hebust.edu.cn College of Sciences, Hebei University of Science and Technology, Shijiazhuang, Hebei 050018, P.R. China

### Abstract

By using the extension of the continuation theorem of Ge and Ren and constructing suitable Banach spaces and operators, we investigate the existence of solutions for a *p*-Laplacian boundary value problem with integral boundary condition at resonance on the half-line.

**MSC:** 34B40

**Keywords:** continuation theorem; integral boundary condition; resonance; *p*-Laplacian; boundary value problem

## **1** Introduction

A boundary value problem is said to be at resonance one if the corresponding homogeneous boundary value problem has non-trivial solutions. Mawhin's continuation theorem [1] is an effective tool to investigate the boundary value problems at resonance with linear or semilinear differential operators (see [2–7] and the references cited therein). Boundary value problems with *p*-Laplacian have been widely studied owing to their importance in theory and application of mathematics and physics (see [8–13]). But the *p*-Laplacian boundary value problems at resonance cannot be solved by Mawhin's continuation theorem. In order to solve these problems, Ge and Ren extended Mawhin's continuation theorem and used it to study boundary value problems with *p*-Laplacian [11]. In their new theorem, two projectors (Definition 2.2) *P* and *Q* must be constructed. But it is difficult to give the projector *Q* in many boundary value problems with *p*-Laplacian. In [14], the author extended the theorem in [11] and studied the problem

$$\begin{cases} (\varphi_p(u''))'(t) = f(t, u(t), u'(t), u''(t)), & t \in [0, 1], \\ u''(0) = 0, & u'(0) = \int_0^1 g(t)u'(t) dt, & u'(1) = \int_0^1 h(t)u'(t) dt \end{cases}$$

in finite interval, where Q is not a projector but satisfies suitable conditions,  $\varphi_p(s) = |s|^{p-2}s$ , p > 1,  $\int_0^1 g(t) dt = 1$ ,  $\int_0^1 h(t) dt = 1$ .

Boundary value problems on the half-line arise in various applications such as in the study of the unsteady flow of gas through a semi-infinite porous medium, in analyzing the heat transfer in radial flow between circular disks, in the study of plasma physics, in an



© 2015 Jiang et al. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

analysis of the mass transfer on a rotating disk in a non-Newtonian fluid, *etc.* [15]. In [16], using the continuation theorem of Ge and Ren [11], the author investigated the existence of solutions for the problem

$$\begin{cases} (\varphi_p(u'))' + f(t, u, u') = 0, & 0 < t < +\infty, \\ u(0) = 0, & \varphi_p(u'(+\infty)) = \sum_{i=1}^n \alpha_i \varphi_p(u'(\xi_i)) \end{cases}$$

on the half-line, where *Q* is a projector,  $\alpha_i > 0$ , i = 1, 2, ..., n,  $\sum_{i=1}^{n} \alpha_i = 1$ . In this paper we study the boundary value problem

In this paper, we study the boundary value problem

$$\begin{cases} (\varphi_p(u'))'(t) = \psi(t)f(t, u(t), u'(t)), & t \in [0, +\infty), \\ u'(+\infty) = 0, & u(0) = \int_0^{+\infty} h(t)u(t) dt \end{cases}$$
(1.1)

in infinite interval, where *Q* is not a projector,  $\varphi_p(s) = |s|^{p-2}s$ , p > 1. To the best of our knowledge, this is the first paper to study the boundary value problems at resonance on the half-line where the operator *Q* is not a projector.

In this paper, we will always suppose that the following conditions hold.

- (H<sub>1</sub>)  $\int_0^{+\infty} h(t) dt = 1$ ,  $th(t) \in L^1[0, +\infty)$ ,  $\psi(t) \in L^1[0, +\infty) \cap C[0, +\infty)$ ,  $h(t) \ge 0$ ,  $\psi(t) > 0$ ,  $t \in [0, +\infty)$ .
- (H<sub>2</sub>) f(t, u, v) is continuous in  $[0, \infty) \times \mathbb{R}^2$ . For any r > 0, there exists a constant  $M_r > 0$ such that if  $\frac{|u|}{1+t} \le r$ ,  $|v| \le r$ ,  $t \in [0, +\infty)$  then  $|f(t, u, v)| \le M_r$ , and for any  $\varepsilon > 0$  there exists  $\delta > 0$  such that  $|f(t, u_1, v_1) - f(t, u_2, v_2)| < \varepsilon$  for  $t \in [0, +\infty)$ ,  $u_i, v_i \in \mathbb{R}$ , i = 1, 2, satisfying  $\frac{|u_1 - u_2|}{1+t} < \delta$ ,  $|v_1 - v_2| < \delta$  and  $\frac{|u_i|}{1+t} \le r$ ,  $|v_i| \le r$ .

The paper is organized as follows. The first section provides a short overview of the problem. Section 2 recalls some preliminary facts. Section 3 contains the main result of the paper.

#### 2 Preliminaries

**Definition 2.1** ([11]) Let *X* and *Z* be two Banach spaces with norms  $\|\cdot\|_X$ ,  $\|\cdot\|_Z$ , respectively. An operator  $M: X \cap \text{dom } M \to Z$  is said to be quasi-linear if

(i)  $\operatorname{Im} M := M(X \cap \operatorname{dom} M)$  is a closed subset of *Z*,

(ii) Ker  $M := \{x \in X \cap \text{dom } M : Mx = 0\}$  is linearly homeomorphic to  $\mathbb{R}^n$ ,  $n < \infty$ , where dom M denotes the domain of the operator M.

In this paper, an operator  $T: X \to Z$  is said to be bounded if  $T(V) \subset Z$  is bounded for any bounded subset  $V \subset X$ .

**Definition 2.2** *P* is a projector if  $P: Y \to Y$  is linear and  $P^2x = Px$ , where *Y* is a vector space.

Let  $X_1 = \text{Ker } M$ ,  $P : X \to X_1$  be a projector and  $X_2$  be the complement space of  $X_1$  in X with  $X = X_1 \oplus X_2$ . Let  $\Omega \subset X$  be an open and bounded set with the origin  $\theta \in \Omega$ .

**Definition 2.3** ([14]) Suppose that  $N_{\lambda} : \overline{\Omega} \to Z$ ,  $\lambda \in [0,1]$  is a continuous and bounded operator. Denote  $N_1$  by N. Let  $\Sigma_{\lambda} = \{x \in \overline{\Omega} : Mx = N_{\lambda}x\}$ .  $N_{\lambda}$  is said to be M-quasi-compact in  $\overline{\Omega}$  if there exists a vector subspace  $Z_1$  of Z satisfying dim  $Z_1$  = dim  $X_1$  and two operators Q and R such that the following conditions hold:

- (a) Ker Q = Im M,
- (b)  $QN_{\lambda}x = \theta, \lambda \in (0, 1) \Leftrightarrow QNx = \theta$ ,
- (c)  $R(\cdot, 0)$  is the zero operator and  $R(\cdot, \lambda)|_{\Sigma_{\lambda}} = (I P)|_{\Sigma_{\lambda}}$ ,
- (d)  $M[P+R(\cdot,\lambda)] = (I-Q)N_{\lambda}$ ,

where  $Q: Z \to Z_1$  is continuous, bounded with Q(I - Q) = 0,  $QZ = Z_1$  and  $R: \overline{\Omega} \times [0, 1] \to X_2$  is continuous and compact with  $Pu + R(u, \lambda) \in \text{dom } M$ ,  $u \in \overline{\Omega}$ ,  $\lambda \in [0, 1]$ .

**Theorem 2.1** ([14]) Let X and Z be two Banach spaces with the norms  $\|\cdot\|_X$ ,  $\|\cdot\|_Z$ , respectively, and  $\Omega \subset X$  be an open and bounded nonempty set. Suppose that

 $M: X \cap \operatorname{dom} M \to Z$ 

is a quasi-linear operator and that  $N_{\lambda} : \overline{\Omega} \to Z, \lambda \in [0,1]$  is *M*-quasi-compact. In addition, if the following conditions hold:

(C<sub>1</sub>)  $Mx \neq N_{\lambda}x, \forall x \in \partial \Omega \cap \text{dom} M, \lambda \in (0, 1),$ (C<sub>2</sub>)  $\text{deg}\{JQN, \Omega \cap \text{Ker} M, 0\} \neq 0,$ 

then the abstract equation Mx = Nx has at least one solution in dom  $M \cap \overline{\Omega}$ , where  $N = N_1$ ,  $J : \text{Im } Q \to \text{Ker } M$  is a homeomorphism with  $J(\theta) = \theta$ , deg is the Brouwer degree.

**Remark** In the proof of Theorem 2.1 in [14], the continuity of the operator *M* is not needed. And dom *M* may not be a linear space. But the operators *P*, *R* satisfy  $Pu + R(u, \lambda) \in \text{dom } M$ ,  $u \in \overline{\Omega}$ ,  $\lambda \in [0, 1]$ .

**Lemma 2.1** ([17]) Let  $\varphi_p : \mathbb{R} \to \mathbb{R}$  be a function given by the formula  $\varphi_p(s) = |s|^{p-2}s$ , where p > 1. Then, for any  $u, v \ge 0$ , we have

- (1)  $\varphi_p(u + v) \le \varphi_p(u) + \varphi_p(v), 1$
- (2)  $\varphi_p(u+v) \le 2^{p-2}(\varphi_p(u)+\varphi_p(v)), p \ge 2.$

#### 3 Main results

In the following, we will always assume that *q* satisfies 1/p + 1/q = 1.

Let  $X = \{u \in C^1[0, +\infty) : u'(+\infty) = 0, \sup_{t \in [0, +\infty)} \frac{|u(t)|}{1+t} < +\infty\}$  be endowed with the following norm  $||u||_X = \max\{||u'||_{\infty}, ||\frac{u(t)}{1+t}||_{\infty}\}, Y = \{y \in C[0, +\infty) : \sup_{t \in [0, +\infty)} |y(t)| < \infty\}$  be endowed with the following norm  $||y||_Y = \sup_{t \in [0, +\infty)} |y(t)| := ||y||_{\infty}$ . Take  $Z = \{\psi y : y \in Y\} \times \mathbb{R}$ , with norm  $||(\psi y, c)||_Z = \max\{||y||_{\infty}, |c|\}$ . We know that  $(X, ||\cdot||_X)$  and  $(Z, ||\cdot||_Z)$  are Banach spaces.

Define operator  $T: Y \to \mathbb{R}$  by Ty = c, for  $y \in Y$ , where *c* satisfies

$$\int_0^{+\infty} h(t) \int_0^t \varphi_q \left( \int_s^{+\infty} \psi(r) \left( y(r) - c \right) dr \right) ds \, dt = 0. \tag{3.1}$$

The following lemma shows that the operator T is well defined.

**Lemma 3.1** For  $y \in Y$ , there is only one constant  $c \in \mathbb{R}$  such that Ty = c with  $|c| \le ||y||_{\infty}$ ,  $T: Y \to \mathbb{R}$  is continuous and  $T(ky) = kT(y), k \in \mathbb{R}$ .

*Proof* For  $y \in Y$ , let

$$F(c) = \int_0^{+\infty} h(t) \int_0^t \varphi_q \left( \int_s^{+\infty} \psi(r) (y(r) - c) \, dr \right) \, ds \, dt.$$

Obviously, *F* is continuous and strictly decreasing in  $\mathbb{R}$ . If *y* is a constant, the results hold, clearly. Assume *y* is not a constant. Take  $a = \inf_{t \in [0,+\infty)} y(t)$ ,  $b = \sup_{t \in [0,+\infty)} y(t)$ . It is easy to see that F(a) > 0, F(b) < 0. So, there exists a unique constant  $c \in (a, b)$  such that F(c) = 0, *i.e.*, there is only one constant  $c \in \mathbb{R}$  such that Ty = c with  $|c| \le ||y||_{\infty}$ .

For  $y_1, y_2 \in Y$ , assume  $Ty_1 = a$ ,  $Ty_2 = b$ . By  $h(t) \ge 0$ ,  $\psi(t) > 0$ ,  $\int_0^{+\infty} h(t) dt = 1$  and  $\varphi_q$  being strictly increasing, we obtain that if  $b - a > \sup_{t \in [0, +\infty)} (y_2(t) - y_1(t))$ , then

$$0 = \int_{0}^{+\infty} h(t) \int_{0}^{t} \varphi_{q} \left( \int_{s}^{+\infty} \psi(r) (y_{2}(r) - b) dr \right) ds dt$$
  
=  $\int_{0}^{+\infty} h(t) \int_{0}^{t} \varphi_{q} \left( \int_{s}^{+\infty} \psi(r) [(y_{1}(r) - a) + (y_{2}(r) - y_{1}(r) - (b - a))] dr \right) ds dt$   
<  $\int_{0}^{+\infty} h(t) \int_{0}^{t} \varphi_{q} \left( \int_{s}^{+\infty} \psi(r) (y_{1}(r) - a) dr \right) ds dt = 0,$ 

a contradiction. On the other hand, if  $b - a < \inf_{t \in [0, +\infty)} (y_2(t) - y_1(t))$ , then

$$0 = \int_{0}^{+\infty} h(t) \int_{0}^{t} \varphi_{q} \left( \int_{s}^{+\infty} \psi(r) (y_{2}(r) - b) dr \right) ds dt$$
  
=  $\int_{0}^{+\infty} h(t) \int_{0}^{t} \varphi_{q} \left( \int_{s}^{+\infty} \psi(r) [(y_{1}(r) - a) + (y_{2}(r) - y_{1}(r) - (b - a))] dr \right) ds dt$   
>  $\int_{0}^{+\infty} h(t) \int_{0}^{t} \varphi_{q} \left( \int_{s}^{+\infty} \psi(r) (y_{1}(r) - a) dr \right) ds dt = 0,$ 

a contradiction, too. So, we have  $\inf_{t \in [0,+\infty)}(y_2(t) - y_1(t)) \le b - a \le \sup_{t \in [0,+\infty)}(y_2(t) - y_1(t))$ , *i.e.*,  $|b - a| \le ||y_2 - y_1||_{\infty}$ . So,  $T : Y \to \mathbb{R}$  is continuous. Obviously, T(ky) = kT(y),  $k \in \mathbb{R}$ . The proof is completed.

Define operators  $M: X \cap \operatorname{dom} M \to Z$ ,  $N_{\lambda}: X \to Z$  as follows:

$$Mu(t) = \left[ \left( \varphi_p(u') \right)'(t), T\left( \frac{(\varphi_p(u'))'(t)}{\psi(t)} \right) \right], \qquad N_{\lambda}u(t) = \left[ \lambda \psi(t) f\left( t, u(t), u'(t) \right), 0 \right],$$

where dom  $M = \{u \in X | \frac{(\varphi_p(u'))'}{\psi(t)} \in Y\}.$ 

**Definition 3.1** *u* is a solution of (1.1) if  $u \in \text{dom } M$  satisfies (1.1).

It is clear that  $u \in \text{dom } M$  is a solution of (1.1) if and only if it satisfies Mu = Nu, where  $N = N_1$ .

#### Lemma 3.2 M is a quasi-linear operator.

*Proof* It is easy to get that Ker  $M = \{c | c \in \mathbb{R}\} := X_1$ .

For  $u \in X \cap \text{dom } M$ , if  $Mu = (\psi y, c)$ , then *c* satisfies (3.1) with *y*. On the other hand, if  $y \in Y$ , Ty = c, take

$$u(t) = -\int_0^t \varphi_q\left(\int_s^{+\infty} \psi(r)y(r)\,dr\right)\,ds.$$

By a simple calculation, we get  $u \in X \cap \text{dom } M$  and  $Mu = (\psi y, c)$ . Thus

$$\operatorname{Im} M = \left\{ (\psi y, Ty) | y \in Y \right\} = \left\{ (\psi y, c) | y \in Y, c \text{ satisfying (3.1) with } y \right\}.$$

By the continuity of *T*, we get that  $\text{Im} M \subset Z$  is closed. So, *M* is quasi-linear. The proof is completed.

Take a projector  $P: X \to X_1$  and an operator  $Q: Z \to Z_1$  as follows:

$$(Pu)(t) = u(0), \qquad Q(\psi y, c) = (0, c - Ty),$$

where  $Z_1 = \{(0, c) | c \in \mathbb{R}\}$ . Obviously,  $QZ = Z_1$  and dim  $Z_1 = \dim X_1$ .

Define an operator *R* as

$$R(u,\lambda)(t) = -\int_0^t \varphi_q\left(\int_s^{+\infty} \lambda \psi(r) f(r,u(r),u'(r)) dr\right) ds, \quad (u,\lambda) \in X \times [0,1].$$

**Lemma 3.3** ([6])  $V \subset X$  is relatively compact if  $\{\frac{u(t)}{1+t}|u \in V\}$  and  $\{u'(t)|u \in V\}$  are both bounded, equicontinuous on any compact intervals of  $[0, +\infty)$  and equiconvergent at infinity.

**Lemma 3.4**  $R: \overline{\Omega} \times [0,1] \to X_2$  is continuous and compact,  $Pu + R(u, \lambda) \in \text{dom } M$ ,  $u \in \overline{\Omega}$ ,  $\lambda \in [0,1]$ , where  $X_2 = \{u \in X : u(0) = 0\}$ ,  $\Omega \subset X$  is an open bounded set.

Proof Firstly, we prove that  $R: \overline{\Omega} \times [0,1] \to X_2$  and  $Pu + R(u, \lambda) \in \text{dom } M, u \in \overline{\Omega}, \lambda \in [0,1]$ . Obviously,  $R(u, \lambda)(t) \in C^1[0, +\infty)$ ,  $R(u, \lambda)'(+\infty) = -\lim_{t \to +\infty} \varphi_q(\int_t^{+\infty} \lambda \psi(s)f(s, u(s), u'(s)) ds) = 0$ . By (H<sub>2</sub>), we get  $\frac{|R(u,\lambda)(t)|}{1+t} \le \varphi_q(||\psi||_1 M_{||u||_X}) < +\infty, u \in X$ . Therefore,  $R(u, \lambda) \in X$ . It is clear that  $R(u, \lambda)(0) = 0$ . Thus  $R(u, \lambda) \in X_2$ . Clearly,  $R(u, \lambda) + Pu \in X$ . It follows from  $(\varphi_p(R(u,\lambda)(t) + Pu(t))')' = \lambda \psi(t)f(t, u(t), u'(t))$  and (H<sub>2</sub>) that  $\frac{(\varphi_p(R(u,\lambda)(t) + Pu(t))')'}{\psi(t)} = \lambda f(t, u(t), u'(t)) \in Y$ . So,  $R(u, \lambda) + Pu \in \text{dom } M$ .

Secondly, we show that *R* is continuous.

Since  $\Omega$  is bounded, there exists a constant r > 0 such that  $||u||_X \leq r$ ,  $u \in \overline{\Omega}$ . By (H<sub>2</sub>), there exists a constant  $M_r > 0$  such that  $|f(t, u(t), u'(t))| \leq M_r$ ,  $u \in \overline{\Omega}$ ,  $t \in [0, +\infty)$ . So, we get

$$\left|\int_{t}^{+\infty}\psi(s)f(s,u(s),u'(s))\,ds\right|\leq \|\psi\|_{1}M_{r},\quad t\in[0,+\infty),u\in\overline{\Omega}.$$

By the uniform continuity of  $\varphi_q(x)$  in  $[-\|\psi\|_1 M_r, \max\{1, \|\psi\|_1 M_r\}]$ , we obtain that for any  $\varepsilon > 0$ , there exists a constant  $\delta_{\varepsilon} > 0$  such that

$$|\varphi_q(x_1) - \varphi_q(x_2)| < \varepsilon, \quad |x_1 - x_2| \le \delta_{\varepsilon}, x_1, x_2 \in [-\|\psi\|_1 M_r, \max\{1, \|\psi\|_1 M_r\}].$$

For  $\alpha = \frac{\delta_{\varepsilon}}{\|\psi\|_1}$ , by (H<sub>2</sub>), there exists a constant  $\delta_{\alpha} > 0$  such that if  $u, v \in \overline{\Omega}$ ,  $\|u - v\|_X < \delta_{\alpha}$ , then  $|f(t, u(t), u'(t)) - f(t, v(t), v'(t))| < \alpha$ ,  $t \in [0, \infty)$ . So, we have

$$\begin{aligned} \left| \int_{t}^{+\infty} \psi(s) f\left(s, u(s), u'(s)\right) ds - \int_{t}^{+\infty} \psi(s) f\left(s, v(s), v'(s)\right) ds \right| \\ \leq \int_{t}^{+\infty} \psi(s) \left| f\left(s, u(s), u'(s)\right) - f\left(s, v(s), v'(s)\right) \right| ds \leq \delta_{\varepsilon}, \quad \|u - v\|_{X} < \delta_{\alpha}. \end{aligned}$$

Take  $\delta = \min\{\delta_{\varepsilon}, \delta_{\alpha}\}$ . For  $u, v \in \overline{\Omega}, \lambda, \mu \in [0, 1]$ , if  $||u - v||_X < \delta, |\lambda - \mu| < \delta$ , then

$$\begin{aligned} \left| R(u,\lambda)'(t) - R(v,\mu)'(t) \right| \\ &= \left| \varphi_q \left( \int_t^{+\infty} \lambda \psi(s) f\left(s, u(s), u'(s)\right) ds \right) - \varphi_q \left( \int_t^{+\infty} \mu \psi(s) f\left(s, v(s), v'(s)\right) ds \right) \right| \\ &= \left| \varphi_q(\lambda) \varphi_q \left( \int_t^{+\infty} \psi(s) f\left(s, u(s), u'(s)\right) ds \right) - \varphi_q(\mu) \varphi_q \left( \int_t^{+\infty} \psi(s) f\left(s, v(s), v'(s)\right) ds \right) \right| \\ &\leq \left[ 1 + \varphi_q \left( \|\psi\|_1 M_r \right) \right] \varepsilon, \quad t \in [0, +\infty). \end{aligned}$$

This, together with

$$\begin{aligned} \frac{|R(u,\lambda)(t) - R(v,\mu)(t)|}{1+t} \\ &= \frac{|\int_0^t \varphi_q(\int_s^{+\infty} \lambda \psi(r)f(r,u(r),u'(r))\,dr)\,ds - \int_0^t \varphi_q(\int_s^{+\infty} \mu \psi(r)f(r,v(r),v'(r))\,dr)\,ds|}{1+t} \\ &\leq \frac{\int_0^t |\varphi_q(\int_s^{+\infty} \lambda \psi(r)f(r,u(r),u'(r))\,dr) - \varphi_q(\int_s^{+\infty} \mu \psi(r)f(r,v(r),v'(r))\,dr)|\,ds}{1+t} \\ &= \frac{\int_0^t |R(u,\lambda)'(s) - R(v,\mu)'(s)|\,ds}{1+t} \leq \left\|R(u,\lambda)' - R(v,\mu)'\right\|_{\infty}, \end{aligned}$$

means that  $R: \overline{\Omega} \times [0,1] \to X_2 \cap \operatorname{dom} M$  is continuous.

We will prove that  $R: \overline{\Omega} \times [0,1] \to X_2 \cap \operatorname{dom} M$  is compact.

It is easy to get that  $\{\frac{R(u,\lambda)(t)}{1+t} : u \in \overline{\Omega}, \lambda \in [0,1]\}$  and  $\{R(u,\lambda)'(t) : u \in \overline{\Omega}, \lambda \in [0,1]\}$  are bounded.

For any T > 0,  $t_1, t_2 \in [0, T]$ ,  $t_1 > t_2$ ,  $u \in \overline{\Omega}$ ,  $\lambda \in [0, 1]$ , we have

$$\begin{aligned} \left| \frac{R(u,\lambda)(t_1)}{1+t_1} - \frac{R(u,\lambda)(t_2)}{1+t_2} \right| \\ &= \left| \frac{1}{1+t_2} \int_0^{t_2} \varphi_q \left( \int_s^{+\infty} \lambda \psi(r) f(r,u(r),u'(r)) \, dr \right) ds \\ &- \frac{1}{1+t_1} \int_0^{t_1} \varphi_q \left( \int_s^{+\infty} \lambda \psi(r) f(r,u(r),u'(r)) \, dr \right) ds \right| \\ &\leq \left| \frac{1}{1+t_2} - \frac{1}{1+t_1} \right| \varphi_q \left( M_r \|\psi\|_1 \right) T + \varphi_q \left( M_r \|\psi\|_1 \right) |t_2 - t_1|. \end{aligned}$$

$$\begin{aligned} \left| R(u,\lambda)'(t_1) - R(u,\lambda)'(t_2) \right| \\ &= \left| \varphi_q \left( \int_{t_2}^{+\infty} \lambda \psi(r) f(r,u(r),u'(r)) \, dr \right) - \varphi_q \left( \int_{t_1}^{+\infty} \lambda \psi(r) f(r,u(r),u'(r)) \, dr \right) \right|. \end{aligned}$$

Take  $G(t) = \int_{t}^{+\infty} \lambda \psi(r) f(r, u(r), u'(r)) dr$ . We have

$$|G(t)| \le M_r \|\psi\|_1, \qquad |G(t_2) - G(t_1)| \le M_r \int_{t_2}^{t_1} \psi(r) dr.$$

It follows from the absolute continuity of integral and the uniform continuity of  $\varphi_q(t)$  in  $[-M_r \|\psi\|_1, M_r \|\psi\|_1]$  that  $\{R(u, \lambda)'(t), u \in \overline{\Omega}, \lambda \in [0, 1]\}$  is equicontinuous on [0, T].

For  $u \in \overline{\Omega}$ , since

$$\int_{s}^{+\infty} \lambda \psi(r) \left| f(r, u(r), u'(r)) \right| dr \leq M_{r} \int_{s}^{+\infty} \psi(r) dr$$

and

$$\lim_{s\to+\infty}\int_s^{+\infty}\psi(r)\,dr=0,$$

for any  $\varepsilon > 0$ , there exists a constant  $T_1 > 0$  such that

$$\varphi_q\left(\int_s^{+\infty}\lambda\psi(r)\big|f\big(r,u(r),u'(r)\big)\big|\right)dr < \frac{\varepsilon}{4}, \quad s > T_1, u \in \overline{\Omega}, \lambda \in [0,1].$$

Obviously, there exists a constant  $T > T_1$  such that, for any t > T,

$$\frac{1}{1+t}\varphi_q\big(M_r\|\psi\|_1\big)T_1<\frac{\varepsilon}{4}.$$

Thus, for any  $t_1, t_2 > T$ , we have

$$\begin{split} \frac{R(u,\lambda)(t_1)}{1+t_1} &- \frac{R(u,\lambda)(t_2)}{1+t_2} \bigg| \\ &= \left| \frac{1}{1+t_2} \int_0^{t_2} \varphi_q \left( \int_s^{+\infty} \lambda \psi(r) f(r,u(r),u'(r)) \, dr \right) ds \right| \\ &- \frac{1}{1+t_1} \int_0^{t_1} \varphi_q \left( \int_s^{+\infty} \lambda \psi(r) f(r,u(r),u'(r)) \, dr \right) ds \bigg| \\ &\leq \frac{1}{1+t_2} \int_0^{T_1} \varphi_q \left( \int_s^{+\infty} \lambda \psi(r) \left| f(r,u(r),u'(r)) \right| \, dr \right) ds \\ &+ \frac{1}{1+t_2} \int_{T_1}^{t_2} \varphi_q \left( \int_s^{+\infty} \lambda \psi(r) \left| f(r,u(r),u'(r)) \right| \, dr \right) ds \\ &+ \frac{1}{1+t_1} \int_0^{T_1} \varphi_q \left( \int_s^{+\infty} \lambda \psi(r) \left| f(r,u(r),u'(r)) \right| \, dr \right) ds \end{split}$$

$$+ \frac{1}{1+t_1} \int_{T_1}^{t_1} \varphi_q \left( \int_s^{+\infty} \lambda \psi(r) \left| f\left(r, u(r), u'(r)\right) \right| dr \right) ds$$
  
 
$$\leq \frac{1}{1+t_2} \varphi_q \left( M_r \|\psi\|_1 \right) T_1 + \frac{t_2 - T_1}{1+t_2} \frac{\varepsilon}{4} + \frac{1}{1+t_1} \varphi_q \left( M_r \|\psi\|_1 \right) T_1 + \frac{t_1 - T_1}{1+t_1} \frac{\varepsilon}{4} < \varepsilon$$

and

$$\begin{aligned} R(u,\lambda)'(t_1) - R(u,\lambda)'(t_2) \Big| \\ &= \left| \varphi_q \left( \int_{t_2}^{+\infty} \lambda \psi(r) f(r,u(r),u'(r)) \, dr \right) - \varphi_q \left( \int_{t_1}^{+\infty} \lambda \psi(r) f(r,u(r),u'(r)) \, dr \right) \right| \\ &\leq \varphi_q \left( \int_{t_2}^{+\infty} \lambda \psi(r) \left| f(r,u(r),u'(r)) \right| \, dr \right) + \varphi_q \left( \int_{t_1}^{+\infty} \lambda \psi(r) \left| f(r,u(r),u'(r)) \right| \, dr \right) \\ &< \varepsilon. \end{aligned}$$

By Lemma 3.3, we get that  $\{R(u, \lambda) | u \in \overline{\Omega}, \lambda \in [0, 1]\}$  is relatively compact. The proof is completed.

**Lemma 3.5** Assume that  $\Omega \subset X$  is an open bounded set. Then  $N_{\lambda}$  is M-quasi-compact in  $\overline{\Omega}$ .

*Proof* It is clear that Im *P* = Ker *M*, Ker *Q* = Im *M* and  $QN_{\lambda}x = \theta$ ,  $\lambda \in (0,1) \Leftrightarrow QNx = \theta$ , *i.e.*, Definition 2.3(a) and (b) are satisfied.

For  $u \in \Sigma_{\lambda} = \{\omega \in \overline{\Omega} \cap \operatorname{dom} M : M\omega = N_{\lambda}\omega\}$ , by (H<sub>1</sub>) and (H<sub>2</sub>), we get

$$R(u,\lambda) = -\int_0^t \varphi_q \left( \int_s^{+\infty} \lambda \psi(r) f(r,u(r),u'(r)) \, dr \right) ds$$
$$= -\int_0^t \varphi_q \left( \int_s^{+\infty} (\varphi_p(u'))'(r) \, dr \right) ds$$
$$= u(t) - u(0) = (I - P)u.$$

Clearly,  $R(\cdot, 0) = 0$ . Thus, Definition 2.3(c) is satisfied. For  $u \in \overline{\Omega}$ , we have

$$M[Pu + R(u,\lambda)](t) = [\lambda\psi(t)f(t,u(t),u'(t)), T(\lambda f(t,u(t),u'(t)))] = (I-Q)N_{\lambda}u(t).$$

So, Definition 2.3(d) is satisfied.

Considering (H<sub>2</sub>),

$$\left\|N_{\lambda}(u)-N_{\lambda}(v)\right\|_{Z} \leq \sup_{t\in[0,\infty)} \left|f\left(t,u(t),u'(t)\right)-f\left(t,v(t),v'(t)\right)\right|, \quad u,v\in\overline{\Omega},$$

and

$$\|N_{\lambda}(u)\|_{Z} \leq \sup_{t\in[0,\infty)} |f(t,u(t),u'(t))|, \quad u\in\overline{\Omega},$$

we can obtain that  $N_{\lambda}$  is continuous and bounded in  $\overline{\Omega}$ .

It follows from the continuity and boundedness of *T* that *Q* is continuous and bounded in *Z*. By a simple calculation, we can obtain that  $Q(I - Q)(\psi y, c) = (0, 0), (\psi y, c) \in Z$ .

These, together with Lemma 3.4, mean that  $N_{\lambda}$  is *M*-quasi-compact in  $\overline{\Omega}$ . The proof is completed.

**Theorem 3.1** *Suppose that* (H<sub>1</sub>), (H<sub>2</sub>) *and the following conditions hold:* 

(H<sub>3</sub>) There exist constants  $c_0 > 0$  and l > 0 such that

$$\int_0^{+\infty} h(t) \int_0^t \varphi_q\left(\int_s^{+\infty} \psi(r) f\left(r, u(r), u'(r)\right) dr\right) ds dt \neq 0, \quad |u(t)| > c_0, t \in [0, l], u \in X.$$

(H<sub>4</sub>) There exist nonnegative functions a(t), b(t), c(t) with  $(1 + t)^{p-1}a(t)\psi(t)$ ,  $b(t)\psi(t)$ ,  $c(t)\psi(t) \in L^1[0, +\infty)$  such that

$$|f(t,x,y)| \le a(t)|\varphi_p(x)| + b(t)|\varphi_p(y)| + c(t), \quad a.e. \ t \in [0, +\infty),$$

where  $\|(1+t)^{p-1}a(t)\psi(t)\|_1 l_0^{p-1} + \|b(t)\psi(t)\|_1 < 1$ , if  $1 ; <math>2^{p-2}\|(1+t)^{p-1}a(t) \times \psi(t)\|_1 l_0^{p-1} + \|b(t)\psi(t)\|_1 < 1$ , if  $p \ge 2$ , where  $l_0 = \max\{1, l\}$ .

- (H<sub>5</sub>) There exists a constant  $d_0 > 0$  such that if  $|d| > d_0$ , then one of the following inequalities holds:
  - (1) df(t,d,0) > 0, t ∈ [0,l);
     (2) df(t,d,0) < 0, t ∈ [0,l).</li>

    Then the boundary value problem (1.1) has at least one solution.

In order to prove Theorem 3.1, we show two lemmas.

**Lemma 3.6** Assume  $(H_1)$ - $(H_4)$  hold. Then the set

 $\Omega_1 = \left\{ u \in \operatorname{dom} M | Mu = N_\lambda u, \lambda \in (0, 1) \right\}$ 

is bounded in X.

*Proof* For  $u \in \Omega_1$ , we have  $QN_{\lambda}u = 0$ , *i.e.*, T(f(t, u(t), u'(t))) = 0. By (H<sub>3</sub>), there exists a constant  $t_0 \in [0, l]$  such that  $|u(t_0)| \le c_0$ . Since  $u(t) = u(t_0) + \int_{t_0}^t u'(s) ds$ , then

$$\frac{|u(t)|}{1+t} \leq \frac{c_0 + |t-t_0| \|u'\|_{\infty}}{1+t} \leq c_0 + \max\{1,l\} \|u'\|_{\infty} = c_0 + l_0 \|u'\|_{\infty}, \quad t \in [0,+\infty).$$

Thus

$$\left\|\frac{u}{1+t}\right\|_{\infty} \le c_0 + l_0 \left\|u'\right\|_{\infty}.$$
(3.2)

It follows from  $Mu = N_{\lambda}u$ , (H<sub>4</sub>) and (3.2) that

$$egin{aligned} &| arphi_pig(u'(t)ig) | \ &\leq \int_0^{+\infty} \psi(t) ig[ a(t) ig| arphi_pig(u(t)ig) ig| + b(t) ig| arphi_pig(u'(t)ig) ig| + c(t)ig] dt \end{aligned}$$

$$\leq \|\psi(t)a(t)(1+t)^{p-1}\|_{1}\varphi_{p}\left(\left\|\frac{u}{1+t}\right\|_{\infty}\right) + \|\psi b\|_{1}\varphi_{p}\left(\left\|u'\right\|_{\infty}\right) + \|\psi c\|_{1} \\ \leq \|\psi(t)a(t)(1+t)^{p-1}\|_{1}\varphi_{p}\left(c_{0}+l_{0}\left\|u'\right\|_{\infty}\right) + \|\psi b\|_{1}\varphi_{p}\left(\left\|u'\right\|_{\infty}\right) + \|\psi c\|_{1}.$$

Whenever 1 , by Lemma 2.1, we get

$$\left\| u' \right\|_{\infty} \leq \varphi_{q} \left( \frac{\|\psi c\|_{1} + \|\psi(t)a(t)(1+t)^{p-1}\|_{1}\varphi_{p}(c_{0})}{1 - \|\psi(t)a(t)(1+t)^{p-1}\|_{1}l_{0}^{p-1} - \|\psi b\|_{1}} \right).$$

Whenever p > 2, by Lemma 2.1, we get

$$\left\| u' \right\|_{\infty} \le \varphi_q \left( \frac{\|\psi c\|_1 + 2^{p-2} \|\psi(t)a(t)(1+t)^{p-1}\|_1 \varphi_p(c_0)}{1 - 2^{p-2} \|\psi(t)a(t)(1+t)^{p-1}\|_1 l_0^{p-1} - \|\psi b\|_1} \right)$$

These, together with (3.2), mean that  $\Omega_1$  is bounded in *X*.

Lemma 3.7 Assume (H<sub>1</sub>)-(H<sub>3</sub>) and (H<sub>5</sub>) hold. Then

$$\Omega_2 = \{ u \in \operatorname{Ker} M | QNu = 0 \}$$

*is bounded in X, where*  $N = N_1$ *.* 

*Proof* For  $u \in \Omega_2$ , we have u = a,  $a \in \mathbb{R}$  and Q(Nu) = 0, *i.e.*,

$$\int_0^{+\infty} h(t) \int_0^t \varphi_q \left( \int_s^{+\infty} \psi(r) f(r, a, 0) \, dr \right) ds \, dt = 0.$$

By (H<sub>5</sub>), we get that  $|a| \leq d_0$ . So,  $\Omega_2$  is bounded. The proof is completed.

*Proof of Theorem* 3.1 Let  $\Omega = \{u \in X | ||u|| < r\}$ , where  $r > d_0$  is large enough such that  $\Omega \supset \overline{\Omega}_1 \cup \overline{\Omega}_2$ .

By Lemmas 3.6 and 3.7, we know  $Mu \neq N_{\lambda}u$ ,  $u \in \text{dom } M \cap \partial \Omega$  and  $QNu \neq 0$ ,  $u \in \text{Ker } M \cap \partial \Omega$ .

Let  $H(u, \delta) = \rho \delta u + (1 - \delta)JQNu$ ,  $\delta \in [0, 1]$ ,  $u \in \text{Ker } M \cap \overline{\Omega}$ , where  $J : \text{Im } Q \to \text{Ker } M$  is a homeomorphism with J(0, a) = a,

$$\rho = \begin{cases} -1 & \text{if } (H_5)(1) \text{ holds,} \\ 1 & \text{if } (H_5)(2) \text{ holds.} \end{cases}$$

Define a function

$$\operatorname{sgn}(x) = \begin{cases} 1 & \text{if } x \ge 0, \\ -1 & \text{if } x < 0. \end{cases}$$

For  $u \in \operatorname{Ker} M \cap \partial \Omega$ , we have  $u = a \neq 0$ . Thus

$$H(u,\delta) = \rho \delta a + (1-\delta)(-T(f(t,a,0))).$$

If  $\delta = 1$ ,  $H(u, 1) = \rho a \neq 0$ . If  $\delta = 0$ , by  $QNu \neq 0$ , we get  $H(u, 0) = JQN(a) \neq 0$ . For  $0 < \delta < 1$ , we now prove that  $H(u, \delta) \neq 0$ . Otherwise, if  $H(u, \delta) = 0$ , then

$$T(f(t,a,0)) = \frac{\rho\delta}{1-\delta}a.$$
(3.3)

Since  $||u||_X = |a| = r > d_0$ , for  $t \in [0, l)$ , by (H<sub>5</sub>) and Lemma 3.1, we have

$$\operatorname{sgn}(af(t,a,0)) = \operatorname{sgn}(T(af(t,a,0))) = \operatorname{sgn}[a(T(f(t,a,0)))] = \operatorname{sgn}(\frac{\rho\delta}{1-\delta}a^2) = \operatorname{sgn}(\rho),$$

a contradiction with the definition of  $\rho$ . So,  $H(u, \delta) \neq 0$ ,  $u \in \text{Ker } M \cap \partial \Omega$ ,  $\delta \in [0, 1]$ .

By the homotopy of degree, we get that

$$deg(JQN, \Omega \cap \operatorname{Ker} M, 0)$$
  
=  $deg(H(\cdot, 0), \Omega \cap \operatorname{Ker} M, 0) = deg(H(\cdot, 1), \Omega \cap \operatorname{Ker} M, 0)$   
=  $deg(\rho I, \Omega \cap \operatorname{Ker} M, 0) = deg(\pm I, \Omega \cap \operatorname{Ker} M, 0) = \pm 1 \neq 0.$ 

By Theorem 2.1, we can get that Mu = Nu has at least one solution in  $\overline{\Omega}$ . The proof is completed.

#### 4 Example

7

Example 4.1 Let us consider the following boundary value problem at resonance:

$$\begin{cases} (\varphi_p(u'))'(t) = \psi(t)f(t, u(t), u'(t)), & t \in [0, +\infty), \\ u'(+\infty) = 0, & u(0) = \int_0^{+\infty} h(t)u(t) dt, \end{cases}$$
(4.1)

where  $p = \frac{4}{3}$ ,  $h(t) = e^{-t}$ ,

$$f(t, u, v) = \begin{cases} 0, & t > l, \\ (l-t)(u^{\frac{1}{3}} + \sin^{\frac{1}{3}}v), & t \le l, u, v \in \mathbb{R}, \end{cases}$$

 $\psi(t) = \frac{1}{3}(1+t)^{-\frac{4}{3}}(1+l_0)^{-\frac{4}{3}}e^{-t}$ ,  $l_0 = \max\{1, l\}$ . Take a(t) = b(t) = (l-t), c(t) = 0,  $d_0 = c_0 = 27$ . By a simple calculation, we can obtain  $||(1+t)^{p-1}a(t)\psi(t)||_1 l_0^{p-1} + ||b(t)\psi(t)||_1 < 1$ . It is easy to get that conditions (H<sub>1</sub>)-(H<sub>5</sub>) are satisfied. It follows from Theorem 3.1 that problem (4.1) has at least one solution.

#### **Competing interests**

The authors declare that there is no conflict of interests regarding the publication of this article.

#### Authors' contributions

The authors declare that the study was realized in collaboration with the same responsibility. All authors read and approved the final manuscript.

#### Acknowledgements

The author is grateful to anonymous referees for their constructive comments and suggestions which led to improvement of the original manuscript. This work is supported by the Natural Science Foundation of China (11171088) and the Natural Science Foundation of Hebei Province (A2013208108).

Received: 18 March 2015 Accepted: 14 September 2015 Published online: 05 October 2015

#### References

- 1. Gaines, R, Mawhin, J: Coincidence Degree and Nonlinear Differential Equation. Springer, Berlin (1977)
- Feng, W, Webb, JRL: Solvability of *m*-point boundary value problems with nonlinear growth. J. Math. Anal. Appl. 212, 467-480 (1997)
- 3. Feng, W, Webb, JRL: Solvability of three-point boundary value problems at resonance. Nonlinear Anal. **30**, 3227-3238 (1997)
- Du, Z, Lin, X, Ge, W: Some higher-order multi-point boundary value problem at resonance. J. Comput. Appl. Math. 177, 55-65 (2005)
- 5. Zhang, X, Feng, M, Ge, W: Existence result of second-order differential equations with integral boundary conditions at resonance. J. Math. Anal. Appl. 353, 311-319 (2009)
- Kosmatov, N: Multi-point boundary value problems on an unbounded domain at resonance. Nonlinear Anal. 68, 2158-2171 (2008)
- 7. Liu, Y, Ge, W: Solvability of nonlocal boundary value problems for ordinary differential equations of higher order. Nonlinear Anal. **57**, 435-458 (2004)
- 8. Yang, C, Yan, J: Positive solutions for third-order Sturm-Liouville boundary value problems with *p*-Laplacian. Comput. Math. Appl. **59**, 2059-2066 (2010)
- 9. Anderson, D, Avery, R, Henderson, J: Existence of solutions for a one-dimensional *p*-Laplacian on time scales. J. Differ. Equ. Appl. **10**, 889-896 (2004)
- Goodrich, C: The existence of a positive solution to a second-order delta-nabla p-Laplacian BVP on a time scale. Appl. Math. Lett. 25, 157-162 (2012)
- 11. Ge, W, Ren, J: An extension of Mawhin's continuation theorem and its application to boundary value problems with a *p*-Laplacian. Nonlinear Anal. TMA **58**, 477-488 (2004)
- Graef, JR, Kong, L: First-order singular boundary value problems with *p*-Laplacian on time scales. J. Differ. Equ. Appl. 17, 831-839 (2011)
- Goodrich, C: Existence of a positive solution to a first-order *p*-Laplacian BVP on a time scale. Nonlinear Anal. 74, 1926-1936 (2011)
- Jiang, W: Solvability of fractional differential equations with *p*-Laplacian at resonance. Appl. Math. Comput. 260, 48-56 (2015)
- Agarwal, RP, O'Regan, D: Infinite Interval Problems for Differential, Difference and Integral Equations. Kluwer Academic, Netherlands (2001)
- Jiang, W: Solvability for *p*-Laplacian boundary value problem at resonance on the half-line. Bound. Value Probl. 2013, 207 (2013). doi:10.1186/1687-2770-2013-207
- 17. Kuang, J: Applied Inequalities. Shandong Science and Technology Press, Jinan (2004)

# Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- ► Convenient online submission
- ► Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- ► High visibility within the field
- ► Retaining the copyright to your article

Submit your next manuscript at > springeropen.com