

Research Article

Multiple Positive Solutions for Singular Quasilinear Multipoint BVPs with the First-Order Derivative

Weihua Jiang,^{1,2} Bin Wang,³ and Yanping Guo¹

¹ College of Sciences, Hebei University of Science and Technology, Shijiazhuang, 050018 Hebei, China

² College of Mathematics and Science of Information, Hebei Normal University, Shijiazhuang, 050016 Hebei, China

³ Department of Basic Courses, Hebei Professional and Technological College of Chemical and Pharmaceutical Engineering, Shijiazhuang, 050031 Hebei, China

Correspondence should be addressed to Weihua Jiang, weihuajiang@hebust.edu.cn

Received 28 November 2007; Accepted 1 April 2008

Recommended by Wenming Zou

The existence of at least three positive solutions for differential equation $(\phi_p(u'(t)))' + g(t)f(t, u(t), u'(t)) = 0$, under one of the following boundary conditions: $u(0) = \sum_{i=1}^{m-2} a_i u(\xi_i)$, $\varphi_p(u'(1)) = \sum_{i=1}^{m-2} b_i \varphi_p(u'(\xi_i))$ or $\varphi_p(u'(0)) = \sum_{i=1}^{m-2} a_i \varphi_p(u'(\xi_i))$, $u(1) = \sum_{i=1}^{m-2} b_i u(\xi_i)$ is obtained by using the H. Amann fixed point theorem, where $\varphi_p(s) = |s|^{p-2}s$, $p > 1$, $0 < \xi_1 < \xi_2 < \dots < \xi_{m-2} < 1$, $a_i > 0$, $b_i > 0$, $0 < \sum_{i=1}^{m-2} a_i < 1$, $0 < \sum_{i=1}^{m-2} b_i < 1$. The interesting thing is that $g(t)$ may be singular at any point of $[0,1]$ and f may be noncontinuous.

Copyright © 2008 Weihua Jiang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In this paper, of multiple positive solutions for differential equation

$$(\varphi_p(u'(t)))' + g(t)f(t, u(t), u'(t)) = 0, \quad \text{a. e. } t \in (0, 1), \quad (1.1)$$

subject to boundary conditions:

$$u(0) = \sum_{i=1}^{m-2} a_i u(\xi_i), \quad \varphi_p(u'(1)) = \sum_{i=1}^{m-2} b_i \varphi_p(u'(\xi_i)), \quad (1.2)$$

$$\varphi_p(u'(0)) = \sum_{i=1}^{m-2} a_i \varphi_p(u'(\xi_i)), \quad u(1) = \sum_{i=1}^{m-2} b_i u(\xi_i), \quad (1.3)$$

respectively, where $\varphi_p(s) = |s|^{p-2}s$, $p > 1$, $0 < \xi_1 < \xi_2 < \dots < \xi_{m-2} < 1$, $a_i > 0$, $b_i > 0$, $0 < \sum_{i=1}^{m-2} a_i < 1$, $0 < \sum_{i=1}^{m-2} b_i < 1$, $g(t)$ may be singular at any point of $[0,1]$.

The multipoint boundary value problems for ordinary differential equations arise in a variety of different areas of applied mathematics and physics. The study of the multipoint boundary value problems for linear second-order ordinary differential equations was initiated by Il'in and Moiseev [1, 2]. Since then, nonlinear second-order multipoint boundary value problems have been studied by several authors. We refer the reader to [3–9] and references cited therein. Recently, in [10], Liang and Zhang studied the existence of positive solutions for differential equation

$$(\varphi(u'))' + a(t)f(u(t)) = 0, \quad 0 < t < 1, \quad (1.4)$$

under the boundary conditions (1.2) by using the fixed point index theory. Wang and Hou [11] investigated the multiplicity of solutions for the differential equation

$$(\varphi_p(u'(t)))' + f(t, u(t)) = 0, \quad t \in (0, 1), \quad (1.5)$$

under the boundary conditions (1.3) by utilizing the fixed point theorem for operators on a cone. Guo et al. [12] proved the existence of at least three positive solutions for differential equation

$$(\varphi_p(u'(t)))' + a(t)f(t, u(t)) = 0, \quad t \in (0, 1), \quad (1.6)$$

subject to (1.2) and (1.3), respectively, by using the five-functional-fixed-point theorem, where $a(t)$ and $f(t, u)$ are continuous.

All of the above work was done under the assumption that f is allowed to depend just on u , while the first-order derivative $u'(t)$ is not involved explicitly in the nonlinear term f .

In [13, 14], Wang and Ge and Sun et al. proved the existence of multiple positive solutions for (1.1) subject to boundary conditions:

$$\begin{aligned} u(0) &= \sum_{i=1}^{n-2} \alpha_i u(\xi_i), & u'(1) &= \sum_{i=1}^{n-2} \beta_i u'(\xi_i), \\ u'(0) &= \sum_{i=1}^n \alpha_i u'(\xi_i), & u(1) &= \sum_{i=1}^n \beta_i u(\xi_i), \end{aligned} \quad (1.7)$$

respectively, where g and f are continuous.

However, in the existing literature, few people considered the case where the nonlinear term is not only involved in the first-order derivative but also noncontinuous. Our paper will fill this gap in the literature. The purpose of this paper is to improve and generalize the results in the above mentioned references. We will prove that the problem (1.1), (1.2) and the problem (1.1), (1.3) have at least three positive solutions by using the H. Amann fixed point theorem, where $g(t)$ may be singular at any point of $[0,1]$ and $f(t, u, v)$ may be noncontinuous.

In this paper, we always suppose the following conditions are satisfied:

$$(H_1) \quad 0 = \xi_0 < \xi_1 < \xi_2 < \dots < \xi_{m-2} < \xi_{m-1} = 1, \quad a_i > 0, \quad b_i > 0, \quad 0 < \sum_{i=1}^{m-2} a_i < 1, \quad 0 < \sum_{i=1}^{m-2} b_i < 1;$$

$$(H_2) \quad g(t) \in L^1[0, 1], \quad g(t) > 0, \quad \text{a.e. } t \in [0, 1];$$

(H₃) $f(t, u, v) : [0, 1] \times R \times R \rightarrow [0, \infty)$ is bounded in a bounded subset of $[0, 1] \times R \times R$, $f(t, u(t), v(t))$ is measurable for $u(t), v(t) \in C[0, 1]$, and $f(t, \cdot, \cdot)$ is continuous for a.e. $t \in [0, 1]$.

Sometimes, we will make use of the following conditions.

(H₄) $f(t, \cdot, v)$ and $f(t, u, \cdot)$ are strictly increasing in $[0, \infty)$.

(H₅) $f(t, \cdot, v)$ is strictly increasing in $[0, \infty)$ and $f(t, u, \cdot)$ is strictly decreasing in $(-\infty, 0]$.

2. Preliminaries

Definition 2.1. Let X be a real Banach space and $P \subset X$ be a cone. For $u, v \in X$, we denote

$$\begin{aligned} u \preceq v &\iff u - v \in P, \\ u < v &\iff u - v \in P, \quad \text{but } u \neq v, \\ u \ll v &\iff u - v \in \text{int } P. \end{aligned} \tag{2.1}$$

If P has nonempty interior, then it is called a solid cone. If every ordered interval is bounded, then P is called a normal cone.

Definition 2.2. An operator $T : P \rightarrow X$ is called order preserving if $u \preceq v \Rightarrow Tu \preceq Tv$, strictly order preserving if $u < v \Rightarrow Tu < Tv$, and strongly order preserving if $u \ll v \Rightarrow Tu \ll Tv$.

Lemma 2.3 (see [15, 16]). *Let P be a normal solid cone of a real Banach space X and suppose there exist $y_1, z_1, y_2, z_2 \in X$ such that $y_1 < z_1 < y_2 < z_2$. In addition, suppose that $T : [y_1, z_2] \rightarrow X$ is a completely continuous and strongly order preserving operator such that*

$$y_1 \preceq Ty_1, \quad Tz_1 < z_1, \quad y_2 < Ty_2, \quad Tz_2 \preceq z_2. \tag{2.2}$$

Then T has at least three fixed points x_1, x_2 and x_3 satisfying

$$y_1 \preceq x_1 \ll z_1, \quad y_2 \ll x_2 \preceq z_2, \quad y_2 \not\preceq x_3 \not\preceq z_1. \tag{2.3}$$

3. The positive solutions for the problem (1.1), (1.2)

Let $X = C^1[0, 1]$ with norm $\|u\| = \max\{\max_{t \in [0, 1]} |u(t)|, \max_{t \in [0, 1]} |u'(t)|\}$. Define $P \subset X$ by

$$P = \{u \in X \mid u(t) \geq 0, u'(t) \geq 0, t \in [0, 1]\}. \tag{3.1}$$

Obviously, X is a Banach space and P is a normal solid cone of X .

We can easily get the following lemmas.

Lemma 3.1. *The boundary value problem (1.1), (1.2) has a solution $u(t)$ if and only if $u(t)$ satisfies the equation*

$$u(t) = \int_0^t \varphi_q \left(\int_s^1 g(\tau) f(\tau, u(\tau), u'(\tau)) d\tau + Bf \right) ds + Af, \tag{3.2}$$

where φ_q is the inverse function of φ_p , that is, $\varphi_q = \varphi_p^{-1}$, $1/p + 1/q = 1$, and

$$\begin{aligned} Bu &= \frac{1}{1 - \sum_{i=1}^{m-2} b_i} \sum_{i=1}^{m-2} b_i \int_{\xi_i}^1 g(\tau) u(\tau) d\tau, \quad u \in L^\infty[0, 1], \\ Au &= \frac{1}{1 - \sum_{i=1}^{m-2} a_i} \sum_{i=1}^{m-2} a_i \int_0^{\xi_i} \varphi_q \left(\int_s^1 g(\tau) u(\tau) d\tau + Bu \right) ds, \quad u \in L^\infty[0, 1]. \end{aligned} \quad (3.3)$$

Define an operator T in P by

$$Tu(t) = \int_0^t \varphi_q \left(\int_s^1 g(\tau) f(\tau, u(\tau), u'(\tau)) d\tau + Bf \right) ds + Af. \quad (3.4)$$

Evidently, $u(t)$ is a fixed point of T if and only if it is a solution of the problem (1.1), (1.2).

Lemma 3.2. Suppose (H_1) – (H_4) hold, $u(t) \in P$, then $Tu(t) \geq 0$, $(Tu)'(t) \geq 0$, $t \in [0, 1]$.

For convenience, we denote

$$Lu(t) = \int_0^t \varphi_q \left(\int_s^1 g(\tau) u(\tau) d\tau + Bu \right) ds + Au, \quad u \in L^\infty[0, 1]. \quad (3.5)$$

Obviously, we have $Tu(t) = Lf(t, u(t), u'(t))$.

Lemma 3.3. Suppose (H_2) holds. If $u_1(t), u_2(t) \in L^\infty[0, 1]$ and $u_1(t) \leq (<) u_2(t)$, a.e. $t \in [0, 1]$, then $L_1 u_1(t) \leq (<) L_1 u_2(t)$, $t \in [0, 1]$.

Assume (H_1) – (H_4) hold. By Lemmas 3.2 and 3.3, the absolute continuity of integral, Ascoli-Arzelà theorem and Lebesgue-Dominated-Convergence-theorem, we obtain that $T : P \rightarrow P$ is completely continuous and T is strongly order preserving.

Theorem 3.4. Suppose (H_1) – (H_4) hold. In addition, suppose there exist constants $0 < a < c < b < d$ and $u_0(t) \in P \setminus \{0\}$ satisfying $cLu_0(1) < bAu_0$ such that

- (A₁) $f(t, aAu_0, 0) \geq \varphi_p(a)u_0(t)$, a.e. $t \in [0, 1]$;
- (A₂) $f(t, cLu_0(1), c(Lu_0)'(0)) < \varphi_p(c)u_0(t)$, a.e. $t \in [0, 1]$;
- (A₃) $f(t, bAu_0, 0) > \varphi_p(b)u_0(t)$, a.e. $t \in [0, 1]$;
- (A₄) $f(t, dLu_0(1), d(Lu_0)'(0)) \leq \varphi_p(d)u_0(t)$, a.e. $t \in [0, 1]$.

Then the problem (1.1), (1.2) has at least three positive solutions $u_1(t)$, $u_2(t)$, and $u_3(t)$ satisfying

$$\begin{aligned} aLu_0(t) \preceq u_1(t) \ll cLu_0(t), \quad bLu_0(t) \ll u_2(t) \preceq dLu_0(t), \\ bLu_0(t) \not\preceq u_3(t) \not\preceq cLu_0(t). \end{aligned} \quad (3.6)$$

Proof. Let $y_1(t) = aLu_0(t)$, $z_1(t) = cLu_0(t)$, $y_2(t) = bLu_0(t)$, $z_2(t) = dLu_0(t)$. Obviously, we have

$$y_1(t) < z_1(t) < y_2(t) < z_2(t). \quad (3.7)$$

Firstly, we will show $y_1(t) \preceq Ty_1(t)$.

By $y_1(t) \geq aAu_0$, $y_1'(t) \geq 0$, (A_1) and Lemma 3.3, we have

$$\begin{aligned} Ty_1(t) &= Lf(t, y_1(t), y_1'(t)) \geq Lf(t, aAu_0, 0) \geq aLu_0(t) = y_1(t), \\ (Ty_1)'(t) &= \varphi_q \left(\int_t^1 g(\tau) f(\tau, y_1(\tau), y_1'(\tau)) d\tau + Bf \right) \geq a \left[\varphi_q \int_t^1 g(\tau) u_0(\tau) d\tau + Bu_0 \right] = y_1'(t). \end{aligned} \quad (3.8)$$

So, we get $y_1 \preceq Ty_1$.

Similarly, by (A_3) and Lemma 3.3, we get $y_2 < Ty_2$.

Next, we prove $Tz_1 < z_1$.

From $z_1(t) \leq cLu_0(1)$, $z_1'(t) \leq c(Lu_0)'(0)$, (A_2) , and Lemma 3.3, we have

$$\begin{aligned} Tz_1(t) &= Lf(t, z_1(t), z_1'(t)) \leq Lf(t, cLu_0(1), c(Lu_0)'(0)) < cLu_0(t) = z_1(t), \\ (Tz_1)'(t) &= \varphi_q \left(\int_t^1 g(\tau) f(\tau, z_1(\tau), z_1'(\tau)) d\tau + Bf \right) \leq c \left[\varphi_q \int_t^1 g(\tau) u_0(\tau) d\tau + Bu_0 \right] = z_1'(t). \end{aligned} \quad (3.9)$$

So, we get $Tz_1 < z_1$.

Similarly, by (A_4) and Lemma 3.3, we get $Tz_2 \preceq z_2$.

By Lemma 2.3, we get that the operator T has at least three fixed points $u_1(t)$, $u_2(t)$, and $u_3(t)$ satisfying

$$\begin{aligned} aLu_0(t) \preceq u_1(t) \ll cLu_0(t), \quad bLu_0(t) \ll u_2(t) \preceq dLu_0(t), \\ bLu_0(t) \not\preceq u_3(t) \not\preceq cLu_0(t). \end{aligned} \quad (3.10)$$

The proof is completed. □

4. The positive solutions for the problem (1.1), (1.3)

Let X be the same as the one in Section 3. Define $P_1 \subset X$ by

$$P_1 = \{u \in X \mid u(t) \geq 0, u'(t) \leq 0, t \in [0, 1]\}. \quad (4.1)$$

Evidently, P_1 is a normal solid cone of X .

We can easily get the following lemmas.

Lemma 4.1. *The boundary value problem (1.1), (1.3) has a solution $u(t)$ if and only if $u(t)$ satisfies the equation*

$$u(t) = - \int_0^t \varphi_q \left(\int_0^s g(\tau) f(\tau, u(\tau), u'(\tau)) d\tau + \tilde{B}f \right) ds + \tilde{A}f, \quad (4.2)$$

where φ_q is the same as the one in Lemma 3.1, and

$$\begin{aligned}\tilde{B}u &= \frac{1}{1 - \sum_{i=1}^{m-2} a_i} \sum_{i=1}^{m-2} a_i \int_0^{\xi_i} g(\tau)u(\tau)d\tau, \quad u \in L^\infty[0, 1], \\ \tilde{A}u &= \frac{1}{1 - \sum_{i=1}^{m-2} b_i} \left[\int_0^1 \varphi_q \left(\int_0^s g(\tau)u(\tau)d\tau + \tilde{B}u \right) ds \right. \\ &\quad \left. - \sum_{i=1}^{m-2} b_i \int_0^{\xi_i} \varphi_q \left(\int_0^s g(\tau)u(\tau)d\tau + \tilde{B}u \right) ds \right], \quad u \in L^\infty[0, 1].\end{aligned}\tag{4.3}$$

Define an operator T_1 in P_1 by

$$T_1u(t) = - \int_0^t \varphi_q \left(\int_0^s g(\tau)f(\tau, u(\tau), u'(\tau))d\tau + \tilde{B}f \right) ds + \tilde{A}f.\tag{4.4}$$

Obviously, $u(t) \in P_1$ is a fixed point of the operator T_1 if and only if it is a positive solution of the problem (1.1), (1.3).

Lemma 4.2. *Suppose (H_1) – (H_3) , (H_5) hold, and $u(t) \in P_1$, then $T_1u(t) \geq 0$, $(T_1u)'(t) \leq 0$, $t \in [0, 1]$.*

For convenience, we denote

$$L_1u(t) = - \int_0^t \varphi_q \left(\int_0^s g(\tau)u(\tau)d\tau + \tilde{B}u \right) ds + \tilde{A}u.\tag{4.5}$$

Clearly, $T_1u(t) = L_1f(t, u(t), u'(t))$.

Lemma 4.3. *Suppose (H_2) holds. If $u_1(t), u_2(t) \in L^\infty[0, 1]$, and $u_1(t) \leq (<)u_2(t)$, a.e. $t \in [0, 1]$, then $L_1u_1(t) \leq (<)L_1u_2(t)$, $t \in [0, 1]$.*

Assume (H_1) – (H_3) and (H_5) hold, by Lemmas 4.2 and 4.3, the absolute continuity of integral, Ascoli-Arzela theorem, and Lebesgue-Dominated-Convergence-theorem, we obtain that $T_1 : P_1 \rightarrow P_1$ is completely continuous, and T_1 is strongly order preserving.

Theorem 4.4. *Suppose (H_1) – (H_3) and (H_5) hold. In addition, suppose there exist constants $0 < \tilde{a} < \tilde{c} < \tilde{b} < \tilde{d}$ and function $u_0(t) \in P_1 \setminus \{0\}$ satisfying $\tilde{c}\tilde{A}u_0 < \tilde{b}L_1u_0(1)$ such that*

- (C₁) $f(t, \tilde{a}L_1u_0(1), 0) \geq \varphi_p(\tilde{a})u_0(t)$, a.e. $t \in [0, 1]$;
- (C₂) $f(t, \tilde{c}\tilde{A}u_0, \tilde{c}(L_1u_0)'(1)) < \varphi_p(\tilde{c})u_0(t)$, a.e. $t \in [0, 1]$;
- (C₃) $f(t, \tilde{b}L_1u_0(1), 0) > \varphi_p(\tilde{b})u_0(t)$, a.e. $t \in [0, 1]$;
- (C₄) $f(t, \tilde{d}\tilde{A}u_0, \tilde{d}(L_1u_0)'(1)) \leq \varphi_p(\tilde{d})u_0(t)$, a.e. $t \in [0, 1]$.

Then the problem (1.1), (1.3) has at least three positive solutions $v_1(t), v_2(t)$, and $v_3(t)$ satisfying

$$\begin{aligned}\tilde{a}L_1u_0(t) \preceq v_1(t) \ll \tilde{c}L_1u_0(t), \quad \tilde{b}L_1u_0(t) \ll v_2(t) \preceq \tilde{d}L_1u_0(t), \\ \tilde{b}L_1u_0(t) \not\preceq v_3(t) \not\preceq \tilde{c}L_1u_0(t).\end{aligned}\tag{4.6}$$

Proof. Let $y_1(t) = \tilde{a}L_1u_0(t)$, $z_1(t) = \tilde{c}L_1u_0(t)$, $y_2(t) = \tilde{b}L_1u_0(t)$, $z_2(t) = \tilde{d}L_1u_0(t)$. Obviously, we have

$$y_1(t) < z_1(t) < y_2(t) < z_2(t). \quad (4.7)$$

Firstly, we will show $y_1(t) \preceq T_1y_1(t)$.

By $y_1(t) \geq \tilde{a}L_1u_0(1)$, $y_1'(t) \leq 0$, (C₁), and Lemma 4.3, we have

$$\begin{aligned} T_1y_1(t) &= L_1f(t, y_1(t), y_1'(t)) \geq L_1f(t, \tilde{a}L_1u_0(1), 0) \geq \tilde{a}L_1u_0(t) = y_1(t), \\ (T_1y_1)'(t) &= -\varphi_q \left(\int_0^t g(\tau) f(\tau, y_1(\tau), y_1'(\tau)) d\tau + \tilde{B}f \right) \\ &\leq -\tilde{a}\varphi_q \left[\int_0^t g(\tau) u_0(\tau) d\tau + \tilde{B}u_0 \right] = y_1'(t). \end{aligned} \quad (4.8)$$

So, we get $y_1 \preceq T_1y_1$.

Similarly, by (C₃) and Lemma 4.3, we get $y_2 < T_1y_2$.

Next, we will prove $T_1z_1 < z_1$.

From $z_1(t) \leq \tilde{c}\tilde{A}u_0$, $z_1'(t) \geq \tilde{c}(L_1u_0)'(1)$, (C₂), and Lemma 4.3, we have

$$\begin{aligned} T_1z_1(t) &= L_1f(t, z_1(t), z_1'(t)) \leq L_1f(t, \tilde{c}\tilde{A}u_0, \tilde{c}(L_1u_0)'(1)) < \tilde{c}L_1u_0(t) = z_1(t), \\ (T_1z_1)'(t) &= -\varphi_q \left(\int_0^t g(\tau) f(\tau, z_1(\tau), z_1'(\tau)) d\tau + \tilde{B}f \right) \\ &\geq -\tilde{c}\varphi_q \left[\int_0^t g(\tau) u_0(\tau) d\tau + \tilde{B}u_0 \right] = z_1'(t). \end{aligned} \quad (4.9)$$

So, we get $T_1z_1 < z_1$.

Similarly, by (C₄) and Lemma 4.3, we get $T_1z_2 \preceq z_2$.

By Lemma 2.3, we get that the operator T_1 has at least three fixed points $v_1(t)$, $v_2(t)$, and $v_3(t)$ satisfying

$$\begin{aligned} \tilde{a}L_1u_0(t) \preceq v_1(t) << \tilde{c}L_1u_0(t), \quad \tilde{b}L_1u_0(t) << v_2(t) \preceq \tilde{d}L_1u_0(t), \\ \tilde{b}L_1u_0(t) \not\preceq v_3(t) \not\preceq \tilde{c}L_1u_0(t). \end{aligned} \quad (4.10)$$

The proof is completed. □

Acknowledgments

This work is supported by the Natural Science Foundation of China (10701032), Science and Technology Key Project of Hebei Province (07217169), and the Foundation of Hebei University of Science and Technology (XI200747).

References

- [1] V. A. Il'in and E. I. Moiseev, "Nonlocal boundary-value problem of the second kind for a Sturm-Liouville operator," *Differential Equations*, vol. 23, no. 8, pp. 979–987, 1987.
- [2] V. A. Il'in and E. I. Moiseev, "Nonlocal boundary value problem of the first kind for a Sturm-Liouville operator in its differential and finite-difference aspects," *Differential Equations*, vol. 23, no. 7, pp. 803–810, 1987.
- [3] H. Lü, D. O'Regan, and R. P. Agarwal, "Existence theorems for the one-dimensional singular p -Laplacian equation with a nonlinear boundary condition," *Journal of Computational and Applied Mathematics*, vol. 182, no. 1, pp. 188–210, 2005.
- [4] R. Ma, "Positive solutions of a nonlinear three-point boundary-value problem," *Electronic Journal of Differential Equations*, vol. 1999, no. 34, pp. 1–8, 1999.
- [5] W. Jiang and F. Li, "Several existence theorems of monotone positive solutions for third-order multipoint boundary value problems," *Boundary Value Problems*, vol. 2007, Article ID 17951, 9 pages, 2007.
- [6] Y. Guo and W. Ge, "Positive solutions for three-point boundary value problems with dependence on the first order derivative," *Journal of Mathematical Analysis and Applications*, vol. 290, no. 1, pp. 291–301, 2004.
- [7] W. Jiang and Y. Guo, "Multiple positive solutions for second-order m -point boundary value problems," *Journal of Mathematical Analysis and Applications*, vol. 327, no. 1, pp. 415–424, 2007.
- [8] W. Jiang, "Multiple positive solutions for n th-order m -point boundary value problems with all derivatives," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 68, no. 5, pp. 1064–1072, 2008.
- [9] W. Jiang, "Eigenvalue criteria for existence of multiple positive solutions of high-order nonlinear BVPs," *Nonlinear Analysis: Theory, Methods & Applications*. In press.
- [10] S. Liang and J. Zhang, "The existence of countably many positive solutions for nonlinear singular m -point boundary value problems," *Journal of Computational and Applied Mathematics*, vol. 214, no. 1, pp. 78–89, 2008.
- [11] Y. Wang and C. Hou, "Existence of multiple positive solutions for one-dimensional p -Laplacian," *Journal of Mathematical Analysis and Applications*, vol. 315, no. 1, pp. 144–153, 2006.
- [12] Y. Guo, Y. Ji, and X. Liu, "Multiple positive solutions for multi-point boundary value problems with p -Laplacian," *Journal of Computational and Applied Mathematics*, vol. 216, no. 1, pp. 144–156, 2008.
- [13] Y. Wang and W. Ge, "Multiple positive solutions for multipoint boundary value problems with one-dimensional p -Laplacian," *Journal of Mathematical Analysis and Applications*, vol. 327, no. 2, pp. 1381–1395, 2007.
- [14] B. Sun, W. Ge, and D. Zhao, "Three positive solutions for multipoint one-dimensional p -Laplacian boundary value problems with dependence on the first order derivative," *Mathematical and Computer Modelling*, vol. 45, no. 9-10, pp. 1170–1178, 2007.
- [15] H. Amann, "Fixed point equations and nonlinear eigenvalue problems in ordered Banach spaces," *SIAM Review*, vol. 18, no. 4, pp. 620–709, 1976.
- [16] D. Guo, *Nonlinear Functional Analysis*, Shandong Science and Technology Press, Jinan, China, 2004.