

Research Article

Positive Solutions of a Nonlinear Three-Point Integral Boundary Value Problem

Jessada Tariboon^{1,2} and Thanin Sitthiwirattham¹

¹ Department of Mathematics, Faculty of Applied Science, King Mongkut's University of Technology North Bangkok, Bangkok 10800, Thailand

² Centre of Excellence in Mathematics, CHE, Sri Ayutthaya Road, Bangkok 10400, Thailand

Correspondence should be addressed to Jessada Tariboon, jsdtrb@hotmail.com

Received 4 August 2010; Accepted 18 September 2010

Academic Editor: Raul F. Manasevich

Copyright © 2010 J. Tariboon and T. Sitthiwirattham. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study the existence of positive solutions to the three-point integral boundary value problem $u'' + a(t)f(u) = 0$, $t \in (0, 1)$, $u(0) = 0$, $\alpha \int_0^\eta u(s) ds = u(1)$, where $0 < \eta < 1$ and $0 < \alpha < 2/\eta^2$. We show the existence of at least one positive solution if f is either superlinear or sublinear by applying the fixed point theorem in cones.

1. Introduction

The study of the existence of solutions of multipoint boundary value problems for linear second-order ordinary differential equations was initiated by Il'in and Moiseev [1]. Then Gupta [2] studied three-point boundary value problems for nonlinear second-order ordinary differential equations. Since then, nonlinear second-order three-point boundary value problems have also been studied by several authors. We refer the reader to [3–19] and the references therein. However, all these papers are concerned with problems with three-point boundary condition restrictions on the slope of the solutions and the solutions themselves, for example,

$$\begin{aligned} u(0) &= 0, & \alpha u(\eta) &= u(1), \\ u(0) &= \beta u(\eta), & \alpha u(\eta) &= u(1), \\ u'(0) &= 0, & \alpha u(\eta) &= u(1), \\ u(0) - \beta u'(0) &= 0, & \alpha u(\eta) &= u(1), \\ \alpha u(0) - \beta u'(0) &= 0, & u'(\eta) + u'(1) &= 0, \end{aligned} \tag{1.1}$$

and so forth.

In this paper, we consider the existence of positive solutions to the equation

$$u'' + a(t)f(u) = 0, \quad t \in (0, 1), \quad (1.2)$$

with the three-point integral boundary condition

$$u(0) = 0, \quad \alpha \int_0^\eta u(s) ds = u(1), \quad (1.3)$$

where $0 < \eta < 1$. We note that the new three-point boundary conditions are related to the area under the curve of solutions $u(t)$ from $t = 0$ to $t = \eta$.

The aim of this paper is to give some results for existence of positive solutions to (1.2)-(1.3), assuming that $0 < \alpha < 2/\eta^2$ and f is either superlinear or sublinear. Set

$$f_0 = \lim_{u \rightarrow 0^+} \frac{f(u)}{u}, \quad f_\infty = \lim_{u \rightarrow \infty} \frac{f(u)}{u}. \quad (1.4)$$

Then $f_0 = 0$ and $f_\infty = \infty$ correspond to the superlinear case, and $f_0 = \infty$ and $f_\infty = 0$ correspond to the sublinear case. By the positive solution of (1.2)-(1.3) we mean that a function $u(t)$ is positive on $0 < t < 1$ and satisfies the problem (1.2)-(1.3).

Throughout this paper, we suppose the following conditions hold:

$$(H1) \quad f \in C([0, \infty), [0, \infty));$$

$$(H2) \quad a \in C([0, 1], [0, \infty)) \text{ and there exists } t_0 \in [\eta, 1] \text{ such that } a(t_0) > 0.$$

The proof of the main theorem is based upon an application of the following Krasnoselskii's fixed point theorem in a cone.

Theorem 1.1 (see [20]). *Let E be a Banach space, and let $K \subset E$ be a cone. Assume Ω_1, Ω_2 are open subsets of E with $0 \in \Omega_1$, $\overline{\Omega_1} \subset \Omega_2$, and let*

$$A : K \cap (\overline{\Omega_1} \setminus \Omega_2) \longrightarrow K \quad (1.5)$$

be a completely continuous operator such that

$$(i) \quad \|Au\| \leq \|u\|, \quad u \in K \cap \partial\Omega_1, \text{ and } \|Au\| \geq \|u\|, \quad u \in K \cap \partial\Omega_2; \text{ or}$$

$$(ii) \quad \|Au\| \geq \|u\|, \quad u \in K \cap \partial\Omega_1, \text{ and } \|Au\| \leq \|u\|, \quad u \in K \cap \partial\Omega_2.$$

Then A has a fixed point in $K \cap (\overline{\Omega_2} \setminus \Omega_1)$.

2. Preliminaries

We now state and prove several lemmas before stating our main results.

Lemma 2.1. *Let $\alpha\eta^2 \neq 2$. Then for $y \in C[0,1]$, the problem*

$$u'' + y(t) = 0, \quad t \in (0,1), \quad (2.1)$$

$$u(0) = 0, \quad \alpha \int_0^\eta u(s) ds = u(1), \quad (2.2)$$

has a unique solution

$$u(t) = \frac{2t}{2 - \alpha\eta^2} \int_0^1 (1-s)y(s) ds - \frac{\alpha t}{2 - \alpha\eta^2} \int_0^\eta (\eta - s)^2 y(s) ds - \int_0^t (t-s)y(s) ds. \quad (2.3)$$

Proof. From (2.1), we have

$$u''(t) = -y(t). \quad (2.4)$$

For $t \in [0, 1]$, integration from 0 to t , gives

$$u'(t) = u'(0) - \int_0^t y(s) ds. \quad (2.5)$$

For $t \in [0, 1]$, integration from 0 to t yields that

$$u(t) = u'(0)t - \int_0^t \left(\int_0^x y(s) ds \right) dx, \quad (2.6)$$

that is,

$$u(t) = u'(0)t - \int_0^t (t-s)y(s) ds. \quad (2.7)$$

So,

$$u(1) = u'(0) - \int_0^1 (1-s)y(s) ds. \quad (2.8)$$

Integrating (2.7) from 0 to η , where $\eta \in (0, 1)$, we have

$$\begin{aligned} \int_0^\eta u(s) ds &= u'(0) \frac{\eta^2}{2} - \int_0^\eta \left(\int_0^x (x-s)y(s) ds \right) dx \\ &= u'(0) \frac{\eta^2}{2} - \frac{1}{2} \int_0^\eta (\eta-s)^2 y(s) ds. \end{aligned} \quad (2.9)$$

From (2.2), we obtain that

$$u'(0) - \int_0^1 (1-s)y(s) ds = u'(0) \frac{\alpha\eta^2}{2} - \frac{\alpha}{2} \int_0^\eta (\eta-s)^2 y(s) ds. \quad (2.10)$$

Thus,

$$u'(0) = \frac{2}{2-\alpha\eta^2} \int_0^1 (1-s)y(s) ds - \frac{\alpha}{2-\alpha\eta^2} \int_0^\eta (\eta-s)^2 y(s) ds. \quad (2.11)$$

Therefore, (2.1)-(2.2) has a unique solution

$$u(t) = \frac{2t}{2-\alpha\eta^2} \int_0^1 (1-s)y(s) ds - \frac{\alpha t}{2-\alpha\eta^2} \int_0^\eta (\eta-s)^2 y(s) ds - \int_0^t (t-s)y(s) ds. \quad (2.12)$$

□

Lemma 2.2. *Let $0 < \alpha < 2/\eta^2$. If $y \in C(0, 1)$ and $y(t) \geq 0$ on $(0, 1)$, then the unique solution u of (2.1)-(2.2) satisfies $u \geq 0$ for $t \in [0, 1]$.*

Proof. If $u(1) \geq 0$, then, by the concavity of u and the fact that $u(0) = 0$, we have $u(t) \geq 0$ for $t \in [0, 1]$.

Moreover, we know that the graph of $u(t)$ is concave down on $(0, 1)$, we get

$$\int_0^\eta u(s) ds \geq \frac{1}{2} \eta u(\eta), \quad (2.13)$$

where $(1/2)\eta u(\eta)$ is the area of triangle under the curve $u(t)$ from $t = 0$ to $t = \eta$ for $\eta \in (0, 1)$.

Assume that $u(1) < 0$. From (2.2), we have

$$\int_0^\eta u(s) ds < 0. \quad (2.14)$$

By concavity of u and $\int_0^\eta u(s) ds < 0$, it implies that $u(\eta) < 0$.

Hence,

$$u(1) = \alpha \int_0^\eta u(s) ds \geq \frac{\alpha\eta}{2} u(\eta) > \frac{u(\eta)}{\eta}, \quad (2.15)$$

which contradicts the concavity of u . \square

Lemma 2.3. *Let $\alpha\eta^2 > 2$. If $y \in C(0, 1)$ and $y(t) \geq 0$ for $t \in (0, 1)$, then (2.1)-(2.2) has no positive solution.*

Proof. Assume (2.1)-(2.2) has a positive solution u .

If $u(1) > 0$, then $\int_0^\eta u(s) ds > 0$, it implies that $u(\eta) > 0$ and

$$\frac{u(1)}{1} = \alpha \int_0^\eta u(s) ds \geq \frac{\alpha\eta}{2} u(\eta) = \frac{\alpha\eta^2}{2} \frac{u(\eta)}{\eta} > \frac{u(\eta)}{\eta}, \quad (2.16)$$

which contradicts the concavity of u .

If $u(1) = 0$, then $\int_0^\eta u(s) ds = 0$, this is $u(t) \equiv 0$ for all $t \in [0, \eta]$. If there exists $\tau \in (\eta, 1)$ such that $u(\tau) > 0$, then $u(0) = u(\eta) < u(\tau)$, which contradicts the concavity of u . Therefore, no positive solutions exist. \square

In the rest of the paper, we assume that $0 < \alpha\eta^2 < 2$. Moreover, we will work in the Banach space $C[0, 1]$, and only the sup norm is used.

Lemma 2.4. *Let $0 < \alpha < 2/\eta^2$. If $y \in C[0, 1]$ and $y \geq 0$, then the unique solution u of the problem (2.1)-(2.2) satisfies*

$$\inf_{t \in [\eta, 1]} u(t) \geq \gamma \|u\|, \quad (2.17)$$

where

$$\gamma := \min \left\{ \eta, \frac{\alpha\eta^2}{2}, \frac{\alpha\eta(1-\eta)}{2-\alpha\eta^2} \right\}. \quad (2.18)$$

Proof. Set $u(\tau) = \|u\|$. We divide the proof into three cases.

Case 1. If $\eta \leq \tau \leq 1$ and $\inf_{t \in [\eta, 1]} u(t) = u(\eta)$, then the concavity of u implies that

$$\frac{u(\eta)}{\eta} \geq \frac{u(\tau)}{\tau} \geq u(\tau). \quad (2.19)$$

Thus,

$$\inf_{t \in [\eta, 1]} u(t) \geq \eta \|u\|. \quad (2.20)$$

Case 2. If $\eta \leq \tau \leq 1$ and $\inf_{t \in [\eta, 1]} u(t) = u(1)$, then (2.2), (2.13), and the concavity of u implies

$$u(1) = \alpha \int_0^\eta u(s) ds \geq \frac{\alpha \eta^2}{2} \left[\frac{u(\eta)}{\eta} \right] \geq \frac{\alpha \eta^2}{2} \frac{u(\tau)}{\tau} \geq \frac{\alpha \eta^2}{2} u(\tau). \quad (2.21)$$

Therefore,

$$\inf_{t \in [\eta, 1]} u(t) \geq \frac{\alpha \eta^2}{2} \|u\|. \quad (2.22)$$

Case 3. If $\tau \leq \eta < 1$, then $\inf_{t \in [\eta, 1]} u(t) = u(1)$. Using the concavity of u and (2.2), (2.13), we have

$$\begin{aligned} u(\sigma) &\leq u(1) + \frac{u(1) - u(\eta)}{1 - \eta} (0 - 1) \\ &\leq u(1) \left[1 - \frac{1 - 2/\alpha \eta}{1 - \eta} \right] \\ &= u(1) \frac{2 - \alpha \eta^2}{\alpha \eta (1 - \eta)}. \end{aligned} \quad (2.23)$$

This implies that

$$\inf_{t \in [\eta, 1]} u(t) \geq \frac{\alpha \eta (1 - \eta)}{2 - \alpha \eta^2} \|u\|. \quad (2.24)$$

This completes the proof. \square

3. Main Results

Now we are in the position to establish the main result.

Theorem 3.1. *Assume (H1) and (H2) hold. Then the problem (1.2)-(1.3) has at least one positive solution in the case*

- (i) $f_0 = 0$ and $f_\infty = \infty$ (superlinear), or
- (ii) $f_0 = \infty$ and $f_\infty = 0$ (sublinear).

Proof. It is known that $0 < \alpha < 2/\eta^2$. From Lemma 2.1, u is a solution to the boundary value problem (1.2)-(1.3) if and only if u is a fixed point of operator A , where A is defined by

$$\begin{aligned} Au(t) = & \frac{2t}{2-\alpha\eta^2} \int_0^1 (1-s)a(s)f(u(s))ds - \frac{\alpha t}{2-\alpha\eta^2} \int_0^\eta (\eta-s)^2 a(s)f(u(s))ds \\ & - \int_0^t (t-s)a(s)f(u(s))ds. \end{aligned} \quad (3.1)$$

Denote that

$$K = \left\{ u \mid u \in C[0,1], u \geq 0, \inf_{\eta \leq t \leq 1} u(t) \geq \gamma \|u\| \right\}, \quad (3.2)$$

where γ is defined in (2.18).

It is obvious that K is a cone in $C[0,1]$. Moreover, by Lemmas 2.2 and 2.4, $AK \subset K$. It is also easy to check that $A : K \rightarrow K$ is completely continuous.

Superlinear Case ($f_0 = 0$ and $f_\infty = \infty$).

Since $f_0 = 0$, we may choose $H_1 > 0$ so that $f(u) \leq \epsilon u$, for $0 < u \leq H_1$, where $\epsilon > 0$ satisfies

$$\frac{2\epsilon}{2-\alpha\eta^2} \int_0^1 (1-s)a(s)ds \leq 1. \quad (3.3)$$

Thus, if we let

$$\Omega_1 = \{u \in C[0,1] \mid \|u\| < H_1\}, \quad (3.4)$$

then, for $u \in K \cap \partial\Omega_1$, we get

$$\begin{aligned} Au(t) & \leq \frac{2t}{2-\alpha\eta^2} \int_0^1 (1-s)a(s)f(u(s))ds \\ & \leq \frac{2t\epsilon}{2-\alpha\eta^2} \int_0^1 (1-s)a(s)u(s)ds \\ & \leq \frac{2\epsilon}{2-\alpha\eta^2} \int_0^1 (1-s)a(s)ds \|u\| \\ & \leq \|u\|. \end{aligned} \quad (3.5)$$

Thus $\|Au\| \leq \|u\|$, $u \in K \cap \partial\Omega_1$.

Further, since $f_\infty = \infty$, there exists $\widehat{H}_2 > 0$ such that $f(u) \geq \rho u$, for $u \geq \widehat{H}_2$, where $\rho > 0$ is chosen so that

$$\rho\gamma \frac{2\eta}{2-\alpha\eta^2} \int_\eta^1 (1-s)a(s)ds \geq 1. \quad (3.6)$$

Let $H_2 = \max\{2H_1, \widehat{H}_2/\gamma\}$ and $\Omega_2 = \{u \in C[0,1] \mid \|u\| < H_2\}$. Then $u \in K \cap \partial\Omega_2$ implies that

$$\inf_{\eta \leq t \leq 1} u(t) \geq \gamma \|u\| = \gamma H_2 \geq \widehat{H}_2, \quad (3.7)$$

and so

$$\begin{aligned} Au(\eta) &= \frac{2\eta}{2-\alpha\eta^2} \int_0^1 (1-s)a(s)f(u(s))ds - \frac{\alpha\eta}{2-\alpha\eta^2} \int_0^\eta (\eta-s)^2 a(s)f(u(s))ds \\ &\quad - \int_0^\eta (\eta-s)a(s)f(u(s))ds \\ &= \frac{2\eta}{2-\alpha\eta^2} \int_0^1 (1-s)a(s)f(u(s))ds - \frac{\alpha\eta}{2-\alpha\eta^2} \int_0^\eta (\eta^2 - 2\eta s + s^2)a(s)f(u(s))ds \\ &\quad - \frac{1}{2-\alpha\eta^2} \int_0^\eta (2-\alpha\eta^2)(\eta-s)a(s)f(u(s))ds \\ &= \frac{2\eta}{2-\alpha\eta^2} \int_0^1 (1-s)a(s)f(u(s))ds + \frac{\alpha\eta^2}{2-\alpha\eta^2} \int_0^\eta sa(s)f(u(s))ds \\ &\quad - \frac{\alpha\eta}{2-\alpha\eta^2} \int_0^\eta s^2 a(s)f(u(s))ds - \frac{2\eta}{2-\alpha\eta^2} \int_0^\eta a(s)f(u(s))ds \\ &\quad + \frac{2}{2-\alpha\eta^2} \int_0^\eta sa(s)f(u(s))ds \\ &= \frac{2\eta}{2-\alpha\eta^2} \int_\eta^1 (1-s)a(s)f(u(s))ds + \frac{2(1-\eta)}{2-\alpha\eta^2} \int_0^\eta sa(s)f(u(s))ds \\ &\quad + \frac{\alpha\eta}{2-\alpha\eta^2} \int_0^\eta s(\eta-s)a(s)f(u(s))ds \\ &\geq \frac{2\eta}{2-\alpha\eta^2} \int_\eta^1 (1-s)a(s)f(u(s))ds \\ &\geq \frac{2\eta\rho}{2-\alpha\eta^2} \int_\eta^1 (1-s)a(s)u(s)ds \geq \frac{2\eta\rho\gamma}{2-\alpha\eta^2} \int_\eta^1 (1-s)a(s)ds \|u\| \geq \|u\|. \end{aligned} \quad (3.8)$$

Hence, $\|Au\| \geq \|u\|$, $u \in K \cap \partial\Omega_2$. By the first part of Theorem 1.1, A has a fixed point in $K \cap (\overline{\Omega}_2 \setminus \Omega_1)$ such that $H_1 \leq \|u\| \leq H_2$.

Sublinear Case ($f_0 = \infty$ and $f_\infty = 0$).

Since $f_0 = \infty$, choose $H_3 > 0$ such that $f(u) \geq Mu$ for $0 < u \leq H_3$, where $M > 0$ satisfies

$$\frac{2\eta\gamma M}{2 - \alpha\eta^2} \int_\eta^1 (1-s)a(s)ds \geq 1. \quad (3.9)$$

Let

$$\Omega_3 = \{u \in C[0,1] \mid \|u\| < H_3\}, \quad (3.10)$$

then for $u \in K \cap \partial\Omega_3$, we get

$$\begin{aligned} Au(\eta) &= \frac{2\eta}{2 - \alpha\eta^2} \int_0^1 (1-s)a(s)f(u(s))ds - \frac{\alpha\eta}{2 - \alpha\eta^2} \int_0^\eta (\eta-s)^2 a(s)f(u(s))ds \\ &\quad - \int_0^\eta (\eta-s)a(s)f(u(s))ds \\ &\geq \frac{2\eta}{2 - \alpha\eta^2} \int_\eta^1 (1-s)a(s)f(u(s))ds \\ &\geq \frac{2\eta\gamma M}{2 - \alpha\eta^2} \int_\eta^1 (1-s)a(s)ds \|u\| \geq \|u\|. \end{aligned} \quad (3.11)$$

Thus, $\|Au\| \geq \|u\|$, $u \in K \cap \partial\Omega_3$. Now, since $f_\infty = 0$, there exists $\widehat{H}_4 > 0$ so that $f(u) \leq \lambda u$ for $u \geq \widehat{H}_4$, where $\lambda > 0$ satisfies

$$\frac{2\lambda}{2 - \alpha\eta^2} \int_0^1 (1-s)a(s)ds \leq 1. \quad (3.12)$$

Choose $H_4 = \max\{2H_3, \widehat{H}_4/\gamma\}$. Let

$$\Omega_4 = \{u \in C[0,1] \mid \|u\| < H_4\}, \quad (3.13)$$

then $u \in K \cap \partial\Omega_4$ implies that

$$\inf_{\eta \leq t \leq 1} u(t) \geq \gamma \|u\| = \gamma H_4 \geq \widehat{H}_4. \quad (3.14)$$

Therefore,

$$\begin{aligned}
 Au(t) &= \frac{2t}{2-\alpha\eta^2} \int_0^1 (1-s)a(s)f(u(s))ds - \frac{\alpha t}{2-\alpha\eta^2} \int_0^\eta (\eta-s)^2 a(s)f(u(s))ds \\
 &\quad - \int_0^t (t-s)a(s)f(u(s))ds \\
 &\leq \frac{2t}{2-\alpha\eta^2} \int_0^1 (1-s)a(s)f(u(s))ds \\
 &\leq \frac{2\lambda\|u\|}{2-\alpha\eta^2} \int_0^1 (1-s)a(s)ds \leq \|u\|.
 \end{aligned} \tag{3.15}$$

Thus $\|Au\| \leq \|u\|$, $u \in K \cap \partial\Omega_4$. By the second part of Theorem 1.1, A has a fixed point u in $K \cap (\overline{\Omega_4} \setminus \Omega_3)$, such that $H_3 \leq \|u\| \leq H_4$. This completes the sublinear part of the theorem. Therefore, the problem (1.2)-(1.3) has at least one positive solution. \square

Acknowledgments

The authors would like to thank the referee for their comments and suggestions on the paper. Especially, the authors would like to thank Dr. Elvin James Moore for valuable advice. This research is supported by the Centre of Excellence in Mathematics, Thailand.

References

- [1] V. A. Il'in and E. I. Moiseev, "Nonlocal boundary-value problem of the first kind for a Sturm-Liouville operator in its differential and finite difference aspects," *Differential Equations*, vol. 23, pp. 803–810, 1987.
- [2] C. P. Gupta, "Solvability of a three-point nonlinear boundary value problem for a second order ordinary differential equations," *Journal of Mathematical Analysis and Applications*, vol. 168, no. 2, pp. 540–551, 1992.
- [3] Z. Chengbo, "Positive solutions for semi-positone three-point boundary value problems," *Journal of Computational and Applied Mathematics*, vol. 228, no. 1, pp. 279–286, 2009.
- [4] Y. Guo and W. Ge, "Positive solutions for three-point boundary value problems with dependence on the first order derivative," *Journal of Mathematical Analysis and Applications*, vol. 290, no. 1, pp. 291–301, 2004.
- [5] X. Han, "Positive solutions for a three-point boundary value problem," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 66, no. 3, pp. 679–688, 2007.
- [6] J. Li and J. Shen, "Multiple positive solutions for a second-order three-point boundary value problem," *Applied Mathematics and Computation*, vol. 182, no. 1, pp. 258–268, 2006.
- [7] S. Liang and L. Mu, "Multiplicity of positive solutions for singular three-point boundary value problems at resonance," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 71, no. 7-8, pp. 2497–2505, 2009.
- [8] R. Liang, J. Peng, and J. Shen, "Positive solutions to a generalized second order three-point boundary value problem," *Applied Mathematics and Computation*, vol. 196, no. 2, pp. 931–940, 2008.
- [9] B. Liu, "Positive solutions of a nonlinear three-point boundary value problem," *Applied Mathematics and Computation*, vol. 132, no. 1, pp. 11–28, 2002.
- [10] B. Liu, "Positive solutions of a nonlinear three-point boundary value problem," *Computers & Mathematics with Applications. An International Journal*, vol. 44, no. 1-2, pp. 201–211, 2002.
- [11] B. Liu, L. Liu, and Y. Wu, "Positive solutions for singular second order three-point boundary value problems," *Nonlinear Analysis. Theory, Methods & Applications.*, vol. 66, no. 12, pp. 2756–2766, 2007.

- [12] H. Luo and Q. Ma, "Positive solutions to a generalized second-order three-point boundary-value problem on time scales," *Electronic Journal of Differential Equations*, vol. 17, pp. 1–14, 2005.
- [13] R. Ma, "Multiplicity of positive solutions for second-order three-point boundary value problems," *Computers & Mathematics with Applications*, vol. 40, no. 2-3, pp. 193–204, 2000.
- [14] R. Ma, "Positive solutions for second-order three-point boundary value problems," *Applied Mathematics Letters*, vol. 14, no. 1, pp. 1–5, 2001.
- [15] R. Ma, "Positive solutions of a nonlinear three-point boundary-value problem," *Electronic Journal of Differential Equations*, vol. 34, pp. 1–8, 1999.
- [16] H. Pang, M. Feng, and W. Ge, "Existence and monotone iteration of positive solutions for a three-point boundary value problem," *Applied Mathematics Letters*, vol. 21, no. 7, pp. 656–661, 2008.
- [17] H.-R. Sun and W.-T. Li, "Positive solutions for nonlinear three-point boundary value problems on time scales," *Journal of Mathematical Analysis and Applications*, vol. 299, no. 2, pp. 508–524, 2004.
- [18] Y. Sun, L. Liu, J. Zhang, and R. P. Agarwal, "Positive solutions of singular three-point boundary value problems for second-order differential equations," *Journal of Computational and Applied Mathematics*, vol. 230, no. 2, pp. 738–750, 2009.
- [19] X. Xu, "Multiplicity results for positive solutions of some semi-positone three-point boundary value problems," *Journal of Mathematical Analysis and Applications*, vol. 291, no. 2, pp. 673–689, 2004.
- [20] M. A. Krasnoselskii, *Positive Solutions of Operator Equations*, Noordhoff, Groningen, The Netherlands, 1964.