RESEARCH

Open Access

Existence of solutions for integral boundary value problems of second-order ordinary differential equations

Hongyu Li^{1*} and Fei Sun²

*Correspondence:

sdlhy1978@163.com

¹Department of Applied Mathematics, Shandong University of Science and Technology, Qingdao, Shandong 266590, P.R. China

Full list of author information is available at the end of the article

Abstract

In this paper, we investigate the existence of solutions for some second-order integral boundary value problems, by applying new fixed point theorems in Banach spaces with the lattice structure derived by Sun and Liu.

MSC: 34B15; 34B18; 47H11

Keywords: lattice; fixed point; integral boundary value problem; sign-changing solution

1 Introduction

In this paper, we consider the following second-order integral boundary value problem:

$$\begin{cases} -x''(t) = f(t, x(t)), & 0 \leq t \leq 1, \\ x(0) = 0, & x(1) = \int_0^1 a(s)x(s)ds, \end{cases} \quad (1.1)$$

where $f \in C([0, 1] \times \mathbb{R}, \mathbb{R})$, $a \in L[0, 1]$ is nonnegative with $\int_0^1 a^2(s)ds < 1$.

The multi-point boundary value problems for ordinary differential equations have been well studied, especially on a compact interval. For example, the study of three-point boundary value problems for nonlinear second-order ordinary differential equations was initiated by Gupta (see [1]). Since then, the existence of solutions for nonlinear multi-point boundary value problems has received much attention from some authors; see [2–6] for reference.

The integral boundary value problems of ordinary differential equations arise in different areas of applied mathematics and physics such as heat conduction, underground water flow, thermo-elasticity and plasma physics (see [7, 8] and the references therein). Moreover, boundary value problems with Riemann-Stieltjes integral conditions constitute a very interesting and important class of problems. They include two, three, multi-point and integral boundary value problems as special cases (see [9, 10]). For boundary value problems with other integral boundary conditions, we refer the reader to the papers [11–21] and the references therein.

In [15], Zhang and Sun studied the following differential equation:

$$\begin{cases} -x''(t) = f(x(t)), & 0 \leq t \leq 1, \\ x(0) = 0, & x(1) = \int_0^1 a(s)x(s)ds, \end{cases}$$

where $f \in C([0, 1] \times \mathbb{R}, \mathbb{R})$, $a \in L[0, 1]$ is nonnegative with $\int_0^1 a^2(s) < 1$. By fixed-point index theory, the existence and multiplicity of sign-changing solutions was discussed.

As we know, nearly all the methods computing the topological degree depend on cone mappings. Recently, Sun and Liu introduced some new computation of topological degree when the concerned operators are not cone mappings in ordered Banach spaces with the lattice structure (for details, see [22–25]). To the best of our knowledge, there is only one paper to use this new computation of topological degree to study an integral boundary value problem with the asymptotically nonlinear term (see [16]).

Motivated by [15, 16, 22–25], this paper is concerned with the boundary value problem (1.1) under sublinear conditions. The method we use is based on some recent fixed point theorems derived by Sun and Liu [22, 23], which are different from [16] and the results we obtain are different from [11–21].

This paper is organized as follows. In Section 2, we recall some properties of the lattice, new fixed point theorems and some lemmas that will be used to prove the main results. In Section 3, we prove the main results and, finally, we give concrete examples to illustrate the applicability of our theory.

2 Preliminaries

We first give some properties of the lattice and give new fixed point theorems with the lattice structure (see [22–25]).

Let E be a Banach space with a cone P . Then E becomes an ordered Banach space under the partial ordering $\leq$ which is induced by P . P is said to be normal if there exists a positive constant N such that $\theta \leq x \leq y$ implies $\|x\| \leq N\|y\|$. P is called solid if it contains interior points, i.e., $\text{int } P \neq \emptyset$.

Definition 2.1 [22–25] We call E a lattice under the partial ordering $\leq$ if $\sup\{x, y\}$ and $\inf\{x, y\}$ exist for arbitrary $x, y \in E$.

For $x \in E$, let

$$x^+ = \sup\{x, \theta\}, \quad x^- = \inf\{x, \theta\},$$

x^+ and x^- are called the positive part and the negative part of x , respectively, and obviously $x = x^+ - x^-$. Take $|x| = x^+ + x^-$, then $|x| \in P$. For convenience, we use the notations $x_+ = x^+$, $x_- = x^-$.

Let $B : E \rightarrow E$ be a bounded linear operator. B is said to be positive if $B(P) \subset P$.

Definition 2.2 [22–25] Let $D \subset E$ and $A : D \rightarrow E$ be a nonlinear operator. A is said to be quasi-additive on lattice if there exists $v^* \in E$ such that

$$Ax = Ax_+ + Ax_- + v^*, \quad \forall x \in D.$$

Let P be a cone of a Banach space E . x is said to be a positive fixed point of A if $x \in (P \setminus \{\theta\})$ is a fixed point of A ; x is said to be a negative fixed point of A if $x \in ((-P) \setminus \{\theta\})$ is a fixed point of A ; x is said to be a sign-changing fixed point of A if $x \notin (P \cup (-P))$ is a fixed point of A .

Lemma 2.1 [22, 23] *Let P be a normal cone of E , and $A : E \rightarrow E$ be completely continuous and quasi-additive on lattice. Suppose that*

- (i) *there exist a positive bounded linear operator B_1 , $u^* \in P$ and $u_1 \in P$ such that*

$$-u^* \leq Ax \leq B_1x + u_1, \quad \forall x \in P;$$

- (ii) *there exist a positive bounded linear operator B_2 and $u_2 \in P$ such that*

$$Ax \geq B_2x - u_2, \quad \forall x \in (-P);$$

- (iii) $r(B_1) < 1$, $r(B_2) < 1$, where $r(B_i)$ is the spectral radius of B_i ($i = 1, 2$);

- (iv) $A\theta = \theta$, the Fréchet derivative A'_θ of A at θ exists, and 1 is not an eigenvalue of the linear operator A'_θ , the sum μ of the algebraic multiplicities for all eigenvalues of A'_θ lying in $(1, +\infty)$ is an odd number.

Then the operator A has at least one nonzero fixed point.

Lemma 2.2 [22] *Let the conditions (i), (ii) and (iii) of Lemma 2.1 be satisfied. Suppose, in addition, that $A\theta = \theta$; the Fréchet derivative A'_θ of A at θ exists; 1 is not an eigenvalue of the linear operator A'_θ ; the sum μ of the algebraic multiplicities for all eigenvalues of A'_θ lying in $(1, +\infty)$ is an even number and*

$$A(P \setminus \{\theta\}) \subset \text{int } P, \quad A((-P) \setminus \{\theta\}) \subset \text{int } (-P).$$

Then the operator A has at least three fixed points: one positive fixed point, one negative fixed point and one sign-changing fixed point.

Let $E = C[0, 1]$ with the normal $\|x(t)\| = \max_{t \in [0, 1]} |x(t)|$, then E is a Banach space. Let $P = \{x \in E : x(t) \geq 0, t \in [0, 1]\}$, then P is a cone of E . It is easy to see that E is a lattice under the partial ordering $\leq$ that is induced by P .

For convenience, list the following condition.

(H₀)

$$0 < \lambda_1 < \lambda_2 < \cdots < \lambda_n < \lambda_{n+1} < \cdots$$

is the sequence of positive solutions of the equation $\sin \sqrt{u} = \int_0^1 a(s) \sin(\sqrt{u}s) ds$.

Define the operators A , B and F :

$$(Bx)(t) = \int_0^1 G(t, s)x(s) ds, \tag{2.1}$$

$$(Fx)(t) = f(t, x(t)), \quad t \in [0, 1], \forall x \in E, \tag{2.2}$$

$$A = BF, \tag{2.3}$$

where

$$G(t, s) = k(t, s) + \frac{t}{1 - \sigma_1} \int_0^1 k(\tau, s) a(\tau) d\tau, \quad \sigma_1 = \int_0^1 sa(s) ds,$$

$$k(t, s) = \begin{cases} t(1 - s), & 0 \leq t \leq s \leq 1, \\ s(1 - t), & 0 \leq s \leq t \leq 1. \end{cases}$$

It is obvious that the fixed points of the operator A defined by (2.3) are the solutions of the boundary value problem (1.1) (see [15, 16]).

Lemma 2.3 [16]

- (i) $B : E \rightarrow E$ is a completely continuous linear operator;
- (ii) $A : E \rightarrow E$ is a completely continuous operator;
- (iii) $A = BF$ is quasi-additive on the lattice;
- (iv) the eigenvalues of the linear operator B are $\{\frac{1}{\lambda_n}, n = 1, 2, \dots\}$ and the algebraic multiplicity of $\frac{1}{\lambda_n}$ is equal to 1, where λ_n is defined by (H_0) ;
- (v) $r(B) = \frac{1}{\lambda_1}$, where $r(B)$ is the spectral radius of the operator B .

3 Main results

Let us list some conditions for convenience.

(H_1) There exists a constant $b > 0$ such that

$$f(t, x) \geq -b, \quad \forall t \in [0, 1], x \geq 0.$$

(H_2) There exists a constant $\epsilon > 0$ such that

$$\limsup_{|x| \rightarrow +\infty} \frac{f(t, x)}{x} \leq \lambda_1 - \epsilon \quad \text{uniformly on } t \in [0, 1],$$

where λ_1 is defined by (H_0) .

(H_3) $f(t, 0) = 0$ uniformly on $t \in [0, 1]$.

(H_4)

$$\lim_{x \rightarrow 0} \frac{f(t, x)}{x} = \lambda \quad \text{uniformly on } t \in [0, 1],$$

where $\lambda_n < \lambda < \lambda_{n+1}$, λ_n, λ_{n+1} is defined by (H_0) .

Theorem 3.1 Suppose that $(H_0), (H_1), (H_2), (H_3), (H_4)$ are satisfied and n is an odd number in (H_4) . Then the boundary value problem (1.1) has at least a nontrivial solution.

Proof Choose $0 < \delta < \epsilon$, then $h = \lambda_1 - \epsilon + \delta < \lambda_1$. By (H_2) , there exists a constant $M > 0$ such that

$$\frac{f(t, x)}{x} \leq h, \quad \forall t \in [0, 1], |x| \geq M. \quad (3.1)$$

So, by (3.1) and (H_1) , we know that

$$f(t, x) \geq hx - C, \quad \forall t \in [0, 1], x \leq 0, \quad (3.2)$$

$$-b \leq f(t, x) \leq hx + C, \quad \forall t \in [0, 1], x \geq 0, \quad (3.3)$$

where $C = \sup_{0 \leq t \leq 1, |x| < M} |f(t, x)|$.

By (3.2) and (3.3), we have

$$(Au)(t) \geq \bar{B}u(t) - C_1, \quad \forall u(t) \in (-P), t \in [0, 1], \quad (3.4)$$

$$-C^* \leq (Au)(t) \leq \bar{B}u(t) + C_1, \quad \forall u(t) \in P, t \in [0, 1], \quad (3.5)$$

where $C_1 = C \int_0^1 G(t, s) ds$, $C^* = b \int_0^1 G(t, s) ds$, $t \in [0, 1]$, $\bar{B} = hB$, B is defined by (2.1). Obviously, $C_1 \in P$, $C^* \in P$, $\bar{B} : P \rightarrow P$ is a positive completely continuous operator. By Lemma 2.3, we have $r(B) = \frac{1}{\lambda_1}$, so $r(\bar{B}) = hr(B) < \lambda_1 r(B) = 1$.

By (H_3) , we have $A\theta = \theta$, and

$$(A'_\theta u)(t) = \int_0^1 G(t, s) f'_x(s, 0) u(s) ds = \lambda \int_0^1 G(t, s) u(s) ds, \quad (3.6)$$

i.e., $A'_\theta = \lambda B$. By Lemma 2.3, 1 is not an eigenvalue of the linear operator A'_θ . Since $\lambda_n < \lambda < \lambda_{n+1}$, n is an odd number, the sum of the algebraic multiplicities for all eigenvalues of A'_θ lying in $(1, +\infty)$ is an odd number. By Lemma 2.1, the operator A has at least one nonzero fixed point. So, the boundary value problem (1.1) has at least one nontrivial solution. $\square$

Theorem 3.2 Suppose (H_0) , (H_2) , (H_3) , (H_4) are satisfied and n is an even number in (H_4) . In addition, assume that

$$f(t, x)x > 0, \quad \forall t \in [0, 1], x \in (-\infty, +\infty), x \neq 0. \quad (3.7)$$

Then the boundary value problem (1.1) has at least three nontrivial solutions: one positive solution, one negative solution and one sign-changing solution.

Proof By (3.7), we have

$$f(t, x) > 0, \quad \forall t \in [0, 1], \forall x > 0; \quad f(t, x) < 0, \quad \forall t \in [0, 1], \forall x < 0. \quad (3.8)$$

By (3.1) and (3.8), (3.4) and (3.5) hold. From (H_3) , (3.6) holds, and by Lemma 2.3, 1 is not an eigenvalue of the linear operator A'_θ . Since $\lambda_n < \lambda < \lambda_{n+1}$, n is an even number, the sum of the algebraic multiplicities for all eigenvalues of A'_θ lying in $(1, +\infty)$ is an even number.

Obviously, from (3.8) and (2.2), we easily get

$$F(P \setminus \{\theta\}) \subset P \setminus \{\theta\}, \quad F((-P) \setminus \{\theta\}) \subset (-P) \setminus \{\theta\}. \quad (3.9)$$

From (2.1), we easily know that $B(P \setminus \{\theta\}) \subset P \setminus \{\theta\}$, $B((-P) \setminus \{\theta\}) \subset (-P) \setminus \{\theta\}$.

So, by (3.9), we have

$$A(P \setminus \{\theta\}) \subset \text{int} P, \quad A((-P) \setminus \{\theta\}) \subset \text{int}(-P).$$

By Lemma 2.2, the boundary value problem (1.1) has at least three nontrivial solutions containing a positive solution, a negative solution and a sign-changing solution. $\square$

Remark By Theorem 3.1 and Theorem 3.2, we can see that the methods used in this paper are different from [11–21], and the results are different from [11–21].

Example 3.1 We consider the following integral boundary value problem:

$$\begin{cases} -x''(t) = f(t, x(t)), & 0 \leq t \leq 1, \\ x(0) = 0, & x(1) = \int_0^1 sx(s) ds, \end{cases} \quad (3.10)$$

where

$$f(t, x) = \begin{cases} 5x + t\sqrt{x} - 40, & t \in [0, 1], x \in (4, +\infty), \\ \frac{t-31}{3}(x-1) + 11 + t, & t \in [0, 1], x \in (1, 4], \\ 10x + (1+t)x^{\frac{5}{3}}, & t \in [0, 1], x \in [-1, 1], \\ \frac{t+29}{7}(x+1) - 11 - t, & t \in [0, 1], x \in (-8, -1], \\ 5x + t\sqrt[3]{x}, & t \in [0, 1], x \in (-\infty, -8]. \end{cases}$$

By simple calculations, we get that $\lambda_1 \approx 7.53$, $\lambda_2 \approx 37.41$, $\lambda_3 \approx 86.80$, $\lambda = 10$. So, $\lambda_1 < \lambda < \lambda_2$. It is easy to know that the nonlinear term f satisfies (H_1) , (H_2) , (H_3) , (H_4) . Thus, the boundary value problem (3.10) has at least a nontrivial solution by Theorem 3.1.

Example 3.2 We consider the following integral boundary value problem:

$$\begin{cases} -x''(t) = f(t, x(t)), & 0 \leq t \leq 1, \\ x(0) = 0, & x(1) = \int_0^1 sx(s) ds, \end{cases} \quad (3.11)$$

where

$$f(t, x) = \begin{cases} 6x + t\sqrt{x}, & t \in [0, 1], x \in (4, +\infty), \\ \frac{t-18}{3}(x-1) + 41 + t, & t \in [0, 1], x \in (1, 4], \\ 40x + (1+t)x^{\frac{5}{3}}, & t \in [0, 1], x \in [-1, 1], \\ \frac{t+7}{7}(x+1) - 41 - t, & t \in [0, 1], x \in (-8, -1], \\ 6x + t\sqrt[3]{x}, & t \in [0, 1], x \in (-\infty, -8]. \end{cases}$$

By simple calculations, we get that $\lambda_1 \approx 7.53$, $\lambda_2 \approx 37.41$, $\lambda_3 \approx 86.80$, $\lambda = 40$. So $\lambda_2 < \lambda < \lambda_3$. It is easy to know that the nonlinear term f satisfies (H_2) , (H_3) , (H_4) and $f(t, x)x > 0$, $\forall t \in [0, 1]$, $x \neq 0$. The boundary value problem (3.11) has at least three nontrivial solutions containing a positive solution, a negative solution and a sign-changing solution by Theorem 3.2.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors typed, read and approved the final manuscript.

Author details

¹Department of Applied Mathematics, Shandong University of Science and Technology, Qingdao, Shandong 266590, P.R. China. ²Jiamusi Hongqi Middle School, Jiamusi, Heilongjiang 154011, P.R. China.

Acknowledgements

The authors would like to thank the reviewers for carefully reading this article and making valuable comments and suggestions. The project is supported by the National Natural Science Foundation of P.R. China (10971179), Research Award Fund for Outstanding Young Scientists of Shandong Province (BS2012SF022, BS2010SF023), Natural Science Foundation of Shandong Province (ZR2010AM035) and SDUST CISE Research Fund.

Received: 24 June 2012 Accepted: 26 November 2012 Published: 17 December 2012

References

1. Gupta, CP: Solvability of a three-point nonlinear boundary value problem for a second order ordinary differential equations. *J. Math. Anal. Appl.* **168**, 540-551 (1992)
2. Zhang, GW, Sun, JX: Multiple positive solutions of singular second order three-point boundary value problems. *J. Math. Anal. Appl.* **317**, 442-447 (2006)
3. Zhang, GW, Sun, JX: Positive solutions of m -point boundary value problems. *J. Math. Anal. Appl.* **291**, 406-418 (2004)
4. Xu, X, Sun, JX: On sign-changing solution for some three-point boundary value problems. *Nonlinear Anal.* **59**, 491-505 (2004)
5. Ma, RY: Nodal solutions for a second-order m -point boundary value problem. *Czechoslov. Math. J.* **56**, 1243-1263 (2006)
6. Zhang, KM, Xie, XJ: Existence of sign-changing solutions for some asymptotically linear three-point boundary value problems. *Nonlinear Anal.* **70**, 2796-2805 (2009)
7. Gallardo, JM: Second order differential operators with integral boundary conditions and generation of semigroups. *Rocky Mt. J. Math.* **30**, 1265-1292 (2000)
8. Timoshenko, S: *Theory of Elastic Stability*. McGraw-Hill, New York (1961)
9. Webb, JRL, Infante, G: Positive solutions of nonlocal boundary value problems: a unified approach. *J. Lond. Math. Soc.* **74**, 673-693 (2006)
10. Karakostas, GL, Tsamatos, PC: Multiple positive solutions of some Fredholm integral equations arisen from nonlocal boundary-value problems. *Electron. J. Differ. Equ.* **30**, 1-17 (2002)
11. Yang, ZL: Existence of nontrivial solutions for a nonlinear Sturm-Liouville problem with integral boundary value conditions. *Nonlinear Anal.* **68**, 216-225 (2008)
12. Jankowski, T: Differential equations with integral boundary conditions. *J. Comput. Appl. Math.* **147**, 1-8 (2002)
13. Yang, ZL: Positive solutions of a second order integral boundary value problem. *J. Math. Anal. Appl.* **321**, 751-765 (2006)
14. Li, Y, Li, F: Sign-changing solutions for second-order integral boundary value problems. *Nonlinear Anal.* **69**, 1179-1187 (2008)
15. Zhang, XQ, Sun, JX: On multiple sign-changing solutions for some second-order integral boundary value problems. *Electron. J. Qual. Theory Differ. Equ.* **44**, 1-15 (2010)
16. Li, HT, Liu, YS: On sign-changing solutions for a second-order integral boundary value problem. *Comput. Math. Appl.* **62**, 651-656 (2011)
17. Zhang, XM, Feng, MQ, Ge, WG: Existence of solutions of boundary value problems with integral boundary conditions for second-order impulsive integro-differential equations in Banach spaces. *J. Comput. Appl. Math.* **233**, 1915-1926 (2010)
18. Feng, MQ, Zhang, XM, Ge, WG: New existence results for higher-order nonlinear fractional differential equation with integral boundary conditions. *Bound. Value Probl.* **2011**, Article ID 720702 (2011)
19. Feng, MQ, Ji, DH, Ge, WG: Positive solutions for a class of boundary value problem with integral boundary conditions in Banach spaces. *J. Comput. Appl. Math.* **222**, 351-363 (2008)
20. Feng, MQ: Existence of symmetric positive solutions for a boundary value problem with integral boundary conditions. *Appl. Math. Lett.* **24**, 1419-1427 (2011)
21. Feng, MQ, Pang, HH: A class of three-point boundary-value problems for second-order impulsive integro-differential equations in Banach spaces. *Nonlinear Anal.* **70**, 64-82 (2009)
22. Sun, JX: *Nonlinear Functional Analysis and Applications*. Science Press, Beijing (2008)
23. Sun, JX, Liu, XY: Computation of topological degree for nonlinear operators and applications. *Nonlinear Anal.* **69**, 4121-4130 (2008)
24. Sun, JX, Liu, XY: Computation of topological degree in ordered Banach spaces with lattice structure and its application to superlinear differential equations. *J. Math. Anal. Appl.* **348**, 927-937 (2008)
25. Liu, XY, Sun, JX: Computation of topological degree of unilaterally asymptotically linear operators and its applications. *Nonlinear Anal.* **71**, 96-106 (2009)

doi:10.1186/1687-2770-2012-147

Cite this article as: Li and Sun: Existence of solutions for integral boundary value problems of second-order ordinary differential equations. *Boundary Value Problems* 2012 **2012**:147.