

RESEARCH

Open Access

# Existence of solutions for nonlocal $p$ -Laplacian thermistor problems on time scales

Wenjing Song<sup>1</sup> and Wenjie Gao<sup>2\*</sup>

\*Correspondence: wjgao@jlu.edu.cn

<sup>2</sup>Institute of Mathematics, Jilin University, Changchun, 130012, P.R. China

Full list of author information is available at the end of the article

## Abstract

In this paper, a nonlocal initial value problem to a  $p$ -Laplacian equation on time scales is studied. The existence of solutions for such a problem is obtained by using the topological degree method.

**Keywords:** existence;  $p$ -Laplacian; time scales; topological degree

## 1 Introduction

In this paper, we are concerned with the existence of solutions of the following nonlocal  $p$ -Laplacian dynamic equation on a time scale  $\mathbb{T}$ :

$$-(\phi_p(u^\Delta(t)))^\nabla = \frac{\lambda a(t)f(u(t))}{(\int_0^T f(u(s))\nabla s)^k}, \quad \forall t \in (0, T)_{\mathbb{T}}, \quad (1.1)$$

with integral initial value

$$\begin{aligned} u(0) &= \int_0^T g(s)u(s)\nabla s, \\ u^\Delta(0) &= A, \end{aligned} \quad (1.2)$$

where  $\phi_p(\cdot)$  is the  $p$ -Laplace operator defined by  $\phi_p(s) = |s|^{p-2}s$ ,  $p > 1$ ,  $\phi_p^{-1} = \phi_q$  with  $q$  the Hölder conjugate of  $p$ , i.e.,  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $\lambda > 0$ ,  $k > 0$ ,  $f : [0, T]_{\mathbb{T}} \rightarrow \mathbb{R}^{+*}$  is continuous ( $\mathbb{R}^{+*}$  denotes positive real numbers),  $a : [0, T]_{\mathbb{T}} \rightarrow \mathbb{R}^+$  is left dense continuous,  $g(s) \in L^1([0, T]_{\mathbb{T}})$  and  $A$  is a real constant.

This model arises in ohmic heating phenomena, which occur in shear bands of metals which are deformed at high strain rates [1, 2], in the theory of gravitational equilibrium of polytropic stars [3], in the investigation of the fully turbulent behavior of real flows, using invariant measures for the Euler equation [4], in modeling aggregation of cells via interaction with a chemical substance (chemotaxis) [5]. For the one-dimensional case, problems with the nonlocal initial condition appear in the investigation of diffusion phenomena for a small amount of gas in a transparent tube [6, 7]; nonlocal initial value problems in higher dimension are important from the point of view of their practical applications to modeling and investigating of pollution processes in rivers and seas, which are caused by sew-age [8].

The study of dynamic equations on time scales has led to some important applications [9–11], and an amount of literature has been devoted to the study the existence of solutions of second-order nonlinear boundary value problems (e.g., see [12–18]).

Motivated by the above works, in this paper, we study the existence of solutions to Problem (1.1), (1.2). Compared with the works mentioned above, this article has the following new features: firstly, the main technique used in this paper is the topological degree method; secondly, Problem (1.1), (1.2) involves the integral initial condition.

The paper is organized as follows. We introduce some necessary definitions and lemmas in the rest of this section. In Section 2, we provide some necessary preliminaries, and in Section 3, the main results are stated and proved.

**Definition 1.1** For  $t < \sup \mathbb{T}$  and  $r > \inf \mathbb{T}$ , define the forward jump operator  $\sigma$  and the backward jump operator  $\rho$ , respectively,

$$\sigma(t) = \inf\{\tau \in \mathbb{T} \mid \tau > t\} \in \mathbb{T}, \quad \rho(r) = \sup\{\tau \in \mathbb{T} \mid \tau < r\} \in \mathbb{T}$$

for all  $t, r \in \mathbb{T}$ . If  $\sigma(t) > t$ ,  $t$  is said to be right scattered, and if  $\rho(r) < r$ ,  $r$  is said to be left scattered. If  $\sigma(t) = t$ ,  $t$  is said to be right dense, and if  $\rho(r) = r$ ,  $r$  is said to be left dense. If  $\mathbb{T}$  has a right scattered minimum  $m$ , define  $\mathbb{T}_k = \mathbb{T} - \{m\}$ ; otherwise, set  $\mathbb{T}_k = \mathbb{T}$ . If  $\mathbb{T}$  has a left scattered maximum  $M$ , define  $\mathbb{T}^k = \mathbb{T} - \{M\}$ ; otherwise, set  $\mathbb{T}^k = \mathbb{T}$ .

**Definition 1.2** For  $x: \mathbb{T} \rightarrow \mathbb{R}$  and  $t \in \mathbb{T}^k$ , we define the delta derivative of  $x(t)$ ,  $x^\Delta(t)$ , to be the number (when it exists) with the property that for any  $\varepsilon > 0$ , there is a neighborhood  $U$  of  $t$  such that

$$\left| [x(\sigma(t)) - x(s)] - x^\Delta(t)[\sigma(t) - s] \right| < \varepsilon |\sigma(t) - s|$$

for all  $s \in U$ . For  $x: \mathbb{T} \rightarrow \mathbb{R}$  and  $t \in \mathbb{T}_k$ , we define the nabla derivative of  $x(t)$ ,  $x^\nabla(t)$ , to be the number (when it exists) with the property that for any  $\varepsilon > 0$ , there is a neighborhood  $V$  of  $t$  such that

$$\left| [x(\rho(t)) - x(s)] - x^\nabla(t)[\rho(t) - s] \right| < \varepsilon |\rho(t) - s|$$

for all  $s \in V$ .

**Definition 1.3** If  $F^\Delta(t) = f(t)$ , then we define the delta integral by

$$\int_a^t f(s) \Delta s = F(t) - F(a).$$

If  $\Phi^\nabla(t) = f(t)$ , then we define the nabla integral by

$$\int_a^t f(s) \nabla s = \Phi(t) - \Phi(a).$$

Throughout this paper, we assume that  $\mathbb{T}$  is a nonempty closed subset of  $\mathbb{R}$  with  $0 \in \mathbb{T}_k$ ,  $T \in \mathbb{T}^k$ .

**Lemma 1.1** (Alternative theorem) *Suppose that  $X$  is a Banach space and  $A$  is a completely continuous operator from  $X$  to  $X$ . Then for any  $\lambda \neq 0$ , only one of the following statements holds:*

(i) *For any  $y \in X$ , there exists a unique  $x \in X$ , such that*

$$(A - \lambda I)x = y;$$

(ii) *There exists an  $x \in X$ ,  $x \neq 0$ , such that*

$$(A - \lambda I)x = 0.$$

## 2 Preliminaries

Let  $E = C_{ld}([0, T]_{\mathbb{T}}, R)$  be a Banach space equipped with the maximum norm  $\|u\| = \max \lim_{[0, T]_{\mathbb{T}}} |u(t)|$ .

Consider the following problem:

$$-(\phi_p(x^\Delta(t)))^\nabla = y(t), \quad \forall t \in (0, T)_{\mathbb{T}}, \tag{2.1}$$

$$x(0) = \int_0^T g(s)x(s)\nabla s, \tag{2.2}$$

$$x^\Delta(0) = A,$$

where  $y \in C([0, T]_{\mathbb{T}})$ ,  $\int_0^T g(s)\nabla s \neq 1$ .

Integrating Eq. (2.1) from 0 to  $t$ , one obtains

$$\phi_p(x^\Delta(t)) - \phi_p(x^\Delta(0)) = - \int_0^t y(s)\nabla s.$$

Using the initial condition (2.2), we have

$$x^\Delta(t) = \phi_p^{-1} \left( \phi_p(A) - \int_0^t y(s)\nabla s \right).$$

Integrating the above equality from 0 to  $t$  again, we obtain

$$x(t) - \int_0^T g(s)x(s)\nabla s = \int_0^t \phi_p^{-1} \left( \phi_p(A) - \int_0^\tau y(s)\nabla s \right) \Delta \tau. \tag{2.3}$$

Let  $F(t) := \int_0^t \phi_p^{-1}(\phi_p(A) - \int_0^\tau y(s)\nabla s) \Delta \tau$ .

Define an operator  $K : C_{ld}([0, T]_{\mathbb{T}}) \rightarrow C_{ld}([0, T]_{\mathbb{T}})$  by

$$(Kx) = \int_0^T g(s)x(s)\nabla s,$$

then (2.3) can be rewritten as

$$(I - K)x(t) = F(t). \tag{2.4}$$

Thus,  $x(t)$  is a solution to (2.1), (2.2) if and only if it is a solution to (2.4).

**Lemma 2.1**  *$I - K$  is a Fredholm operator.*

*Proof* To prove that  $I - K$  is a Fredholm operator, we need only to show that  $K$  is completely continuous.

It is easy to see from the definition of  $K$  that  $K$  is a bounded linear operator from  $C_{ld}([0, T]_{\mathbb{T}})$  to  $C_{ld}([0, T]_{\mathbb{T}})$ . Obviously,  $\dim R(K) = 1$ . So,  $K$  is a completely continuous operator. This completes the proof.  $\square$

**Lemma 2.2** *Problem (2.1), (2.2) admits a unique solution.*

*Proof* Since Problem (2.1), (2.2) is equivalent to Problem (2.4), we need only to show that Problem (2.4) has a unique solution.

Using Lemma 2.1 and the alternative theorem, it is sufficient to prove that

$$(I - K)x(t) = 0 \tag{2.5}$$

has a trivial solution  $x \equiv 0$  only.

On the contrary, suppose (2.5) has a nontrivial solution  $\mu$ , then  $\mu$  is a constant, and we have

$$I\mu = K\mu = \mu.$$

The definition of  $K$  and the above equality yield

$$\left[ 1 - \int_0^T g(s) \nabla s \right] \mu = 0,$$

which is a contradiction to the assumptions  $\int_0^T g(s) \nabla s \neq 1$  and  $\mu \neq 0$ .

Thus, we complete the proof.  $\square$

### 3 Main results

Throughout this section, we assume that the following conditions hold.

- (H1)  $\int_0^T |g(s)| \nabla s = M < 1$ ;
- (H2)  $f : [0, T]_{\mathbb{T}} \rightarrow \mathbb{R}^{**}$  is continuous;
- (H3)  $a : [0, T]_{\mathbb{T}} \rightarrow \mathbb{R}^+$  is left dense continuous and  $\max_{t \in [0, T]_{\mathbb{T}}} a(t) \leq M_1$ ;
- (H4)  $f(y) \leq [c_1 \phi_p(|y|) + c_2]^{\frac{1}{1-k}}$ ,  $c_1, c_2 > 0$  and  $c_1 < \frac{\phi_p(\frac{1-M}{2q-1T})}{\lambda M_1 T^{1-k}}$ , when  $k < 1$ ;
- (H5)  $f(y) \geq [c_3 \phi_p(|y|)]^{\frac{1}{1-k}}$ ,  $c_3 > 0$  and  $c_3 < \frac{\phi_p(\frac{1-M}{2q-1T})}{\lambda M_1 T^{1-k}}$ , when  $k > 1$ .

From Lemma 2.2 we know that  $u(t)$  is a solution to Problem (1.1), (1.2) if and only if it is a solution to the following integral equation:

$$(I - K)u(t) = \int_0^t \phi_p^{-1} \left( \phi_p(A) - \int_0^\tau \frac{\lambda a(s) f(u(s))}{(\int_0^T f(u(s)) \nabla s)^k} \nabla s \right) \Delta \tau. \tag{3.1}$$

Define an operator  $F : C_{ld}([0, T]_{\mathbb{T}}) \rightarrow C_{ld}([0, T]_{\mathbb{T}})$  by

$$(Fu)(t) = \int_0^t \phi_p^{-1} \left( \phi_p(A) - \int_0^\tau \frac{\lambda a(s) f(u(s))}{(\int_0^T f(u(s)) \nabla s)^k} \nabla s \right) \Delta \tau,$$

then (3.1) can be rewritten as

$$(I - K)u(t) = (Fu)(t).$$

In order to prove the existence of solutions to (3.1), we need the following lemmas.

**Lemma 3.1** *F is completely continuous.*

*Proof* Let  $R_1$  be an arbitrary positive real number and denote  $B_1 = \{u \in C_{ld}([0, T]_{\mathbb{T}}); \|u\| \leq R_1\}$ . Then we have for any  $u \in B_1$ ,

$$\begin{aligned} |(Fu)(t)| &\leq \int_0^t \left| \phi_p^{-1} \left( \phi_p(A) - \int_0^\tau \frac{\lambda a(s)f(u(s))}{(\int_0^T f(u(s))\nabla s)^k} \nabla s \right) \right| \Delta \tau \\ &\leq \int_0^T \phi_p^{-1} \left( |\phi_p(A)| + \left| \int_0^T \frac{\lambda a(s) \sup_{u \in B_1} f}{(T \inf_{u \in B_1} f)^k} \nabla s \right| \right) \Delta \tau \\ &\leq \phi_p^{-1} \left( |\phi_p(A)| + M_1 T \frac{\lambda \sup_{u \in B_1} f}{(T \inf_{u \in B_1} f)^k} \right) T. \end{aligned}$$

This shows that  $F(B_1)$  is uniformly bounded.

Moreover, for any  $t \in [0, T]_{\mathbb{T}}$ , we have

$$\begin{aligned} |(Fu)^\Delta(t)| &= \left| \phi_p^{-1} \left( \phi_p(A) - \int_0^t \frac{\lambda a(s)f(u(s))}{(\int_0^T f(u(s))\nabla s)^k} \nabla s \right) \right| \\ &\leq \phi_p^{-1} \left( |\phi_p(A)| + M_1 T \frac{\lambda \sup_{u \in B_1} f}{(T \inf_{u \in B_1} f)^k} \right). \end{aligned}$$

Thus, it is easy to prove that  $F(B_1)$  is equicontinuous. This together with the Ascoli-Arzelà theorem guarantees that  $F(B_1)$  is relatively compact in  $C_{ld}([0, T]_{\mathbb{T}})$ .

Therefore,  $F$  is completely continuous. The proof of Lemma 3.1 is completed. □

**Theorem 3.1** *Assume that conditions (H1)-(H5) hold. Then Problem (1.1), (1.2) has at least one solution.*

*Proof* Lemma 2.1 and Lemma 3.1 imply that the operator  $K + F$  is completely continuous. It suffices for us to prove that the equation

$$(I - (K + F))u = 0 \tag{3.2}$$

has at least one solution.

Define  $H : [0, 1] \times C_{ld}([0, T]_{\mathbb{T}}) \rightarrow C_{ld}([0, T]_{\mathbb{T}})$  as

$$H(\sigma, u) = (K + \sigma F)u,$$

and it is clear that  $H$  is completely continuous.

Set  $h_\sigma(u) = u - H(\sigma, u)$ , then we have

$$h_0(u) = (I - K)u,$$

$$h_1(u) = [I - (K + F)]u.$$

To apply the Leray-Schauder degree to  $h_\sigma$ , we need only to show that there exists a ball  $B_R(\theta)$  in  $C_{id}([0, T]_{\mathbb{T}})$ , whose radius  $R$  will be fixed later, such that  $\theta \notin h_\sigma(\partial B_R(\theta))$ .

If  $k < 1$ , choosing  $R > \frac{2^{q-1}\phi_p^{-1}(|\phi_p(A)| + \lambda M_1 T^{1-k} c_2) T}{1 - M - 2^{q-1}\phi_p^{-1}(\lambda M_1 T^{1-k} c_1) T}$ , then for any fixed  $u \in \partial B_R(\theta)$ , there exists a  $t_0 \in [0, T]_{\mathbb{T}}$  such that  $|u(t_0)| = R$ . By direct calculation, we have

$$\begin{aligned} & |(h_\sigma u)(t_0)| \\ &= \left| u(t_0) - \left[ \int_0^T g(s)u(s)\nabla s + \sigma \int_0^{t_0} \phi_p^{-1} \left( \phi_p(A) - \int_0^\tau \frac{\lambda a(s)f(u(s))}{(\int_0^T f(u(s))\nabla s)^k} \nabla s \right) \Delta \tau \right] \right| \\ &\geq |u(t_0)| - \left| \int_0^T g(s)u(s)\nabla s \right| - \left| \int_0^{t_0} \phi_p^{-1} \left( \phi_p(A) - \int_0^\tau \frac{\lambda a(s)f(u(s))}{(\int_0^T f(u(s))\nabla s)^k} \nabla s \right) \Delta \tau \right| \\ &\geq (1 - M)R - \int_0^T \phi_p^{-1} \left( |\phi_p(A)| + \int_0^T \frac{\lambda a(s)f(u(s))}{(\int_0^T f(u(s))\nabla s)^k} \nabla s \right) \Delta \tau. \end{aligned} \tag{3.3}$$

From (H4), we have

$$\begin{aligned} |(h_\sigma u)(t_0)| &\geq (1 - M)R - \int_0^T \phi_p^{-1} \left( |\phi_p(A)| + \lambda M_1 \left( \int_0^T f(u(s))\nabla s \right)^{1-k} \right) \Delta \tau \\ &\geq (1 - M)R - \int_0^T \phi_p^{-1} [|\phi_p(A)| + \lambda M_1 T^{1-k} (c_1 \phi_p(\|u\|) + c_2)] \Delta \tau \\ &> 0. \end{aligned} \tag{3.4}$$

If  $k > 1$ , choosing  $R > \frac{2^{q-1}|A|T}{1 - M - 2^{q-1}\phi_p^{-1}(\lambda M_1 T^{1-k} c_3) T}$ , then for any fixed  $u \in \partial B_R(\theta)$ , there exists a  $t_0 \in [0, T]_{\mathbb{T}}$  such that  $|u(t_0)| = R$ . From (H5), we have

$$\begin{aligned} |(h_\sigma u)(t_0)| &\geq (1 - M)R - \int_0^T \phi_p^{-1} \left( |\phi_p(A)| + \frac{\lambda M_1}{(\int_0^T f(u(s))\nabla s)^{k-1}} \right) \Delta \tau \\ &\geq (1 - M)R - \int_0^T \phi_p^{-1} [|\phi_p(A)| + \lambda M_1 T^{1-k} c_3 \phi_p(\|u\|)] \Delta \tau \\ &> 0. \end{aligned} \tag{3.5}$$

If  $k = 1$ , choosing  $R > \frac{\phi_p^{-1}(|\phi_p(A)| + \lambda M_1) T}{1 - M}$ , then for any fixed  $u \in \partial B_R(\theta)$ , there exists a  $t_0 \in [0, T]_{\mathbb{T}}$  such that  $|u(t_0)| = R$ . By direct calculation, we have

$$\begin{aligned} |(h_\sigma u)(t_0)| &\geq (1 - M)R - \int_0^T \phi_p^{-1} (|\phi_p(A)| + \lambda M_1) \Delta \tau \\ &> 0. \end{aligned} \tag{3.6}$$

This implies  $h_\sigma u \neq \theta$  and hence we obtain  $\theta \notin h_\sigma(\partial B_R(\theta))$ .

Since  $\deg(h_1, B_R(\theta), \theta) = \deg(h_0, B_R(\theta), \theta) = \pm 1 \neq 0$ , we know that (3.2) admits a solution  $u \in B_R(\theta)$ , which implies that (1.1), (1.2) also admits a solution in  $B_R(\theta)$ .  $\square$

**Competing interests**

All authors declare that they have no competing interests.

#### Authors' contributions

WS drafted this paper and WG checked and corrected the manuscript.

#### Author details

<sup>1</sup>Institute of Applied Mathematics, Jilin University of Finance and Economics, Changchun, 130117, P.R. China. <sup>2</sup>Institute of Mathematics, Jilin University, Changchun, 130012, P.R. China.

#### Acknowledgements

This work was supported by NSFC (11271154) and by Key Lab of Symbolic Computation and Knowledge Engineering of Ministry of Education and by the 985 program of Jilin University, and the first author is also supported by the Youth Studies Program of Jilin University of Finance and Economics (XJ2012006).

Received: 1 September 2012 Accepted: 18 December 2012 Published: 4 January 2013

#### References

1. Bebernes, JW, Lacey, AA: Global existence and finite-time blow-up for a class of non-local parabolic problems. *Adv. Differ. Equ.* **2**, 927-953 (1997)
2. Bebernes, JW, Li, C, Talaga, P: Single-point blow-up for non-local parabolic problems. *Physica D* **134**, 48-60 (1999)
3. Krzywicki, A, Nadzieja, T: Some results concerning the Poisson-Boltzmann equation. *Zastos. Mat.* **21**(2), 265-272 (1991)
4. Caglioti, E, Lions, P-L, Marchioro, C, Pulvirenti, M: A special class of stationary flows for two-dimensional Euler equations: a statistical mechanics description. *Commun. Math. Phys.* **143**, 501-525 (1992)
5. Wolansky, G: A critical parabolic estimate and application to non-local equations arising in chemotaxis. *Appl. Anal.* **66**, 291-321 (1997)
6. Kerefov, AA: Nonlocal boundary value problems for parabolic equations. *Differ. Uravn.* **15**, 74-78 (1979)
7. Vabishchevich, PN: Nonlocal parabolic problems and the inverse heat-conduction problem. *Differ. Uravn.* **17**, 1193-1199 (1981)
8. Gordeziani, DG, Avalishvili, G: Investigation of the nonlocal initial boundary value problems for some hyperbolic equations. *Hiroshima Math. J.* **31**, 345-366 (2001)
9. Agarwal, R, Bohner, M, O'Regan, D, Peterson, A: Dynamic equations on time scales: a survey. *J. Comput. Appl. Math.* **141**, 1-26 (2002)
10. Atici, FM, Guseinov, GS: On Green's functions and positive solutions for boundary value problems on time scales. *J. Comput. Appl. Math.* **141**, 75-99 (2002)
11. Bohner, M, Peterson, A: *Advances in Dynamic Equations on Time Scales*. Birkhäuser, Boston (2003)
12. Hamal, NA, Yoruk, F: Positive solutions of nonlinear  $m$ -point boundary value problems on time scales. *J. Comput. Appl. Math.* **231**, 92-105 (2009)
13. Li, WT, Sun, HR: Positive solutions for second order  $m$ -point boundary value problems on time scales. *Acta Math. Sin. Engl. Ser.* **22**(6), 1797-1804 (2006)
14. Sidi Ammi, MR, Torres, DFM: Existence of positive solutions for non local  $p$ -Laplacian thermistor problems on time scales. *J. Inequal. Pure Appl. Math.* **8**(3), 69 (2007)
15. Sidi Ammi, MR, Torres, DFM: Existence of three positive solutions to some  $p$ -Laplacian boundary value problems. *Discrete Dyn. Nat. Soc.* **2012**, Art. ID 145050 (2012)
16. Sidi Ammi, MR, Torres, DFM: Existence and uniqueness of a positive solution to generalized nonlocal thermistor problems with fractional order derivatives. *Differ. Equ. Appl.* **4**, 267-276 (2012)
17. Wang, DB: Existence multiplicity and infinite solvability of positive solutions for  $p$ -Laplacian dynamic equations on time scales. *Electron. J. Differ. Equ.* **2006**, 96 (2006)
18. Yang, YT, Meng, FW: Positive solutions of the singular semipositone boundary value problem on time scales. *Math. Comput. Model.* **52**, 481-489 (2010)

doi:10.1186/1687-2770-2013-1

**Cite this article as:** Song and Gao: Existence of solutions for nonlocal  $p$ -Laplacian thermistor problems on time scales. *Boundary Value Problems* 2013 **2013**:1.

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)