

RESEARCH

Open Access

Study of solutions to an initial and boundary value problem for certain systems with variable exponents

Yunzhu Gao^{1*} and Wenjie Gao^{2*}

*Correspondence:
yzgao_2008@163.com;
wjgao@jlu.edu.cn

¹Department of Mathematics and
Statistics, Beihua University, Jilin
City, P.R. China

²Institute of Mathematics, Jilin
University, Changchun, 130012, P.R.
China

Abstract

In this paper, the existence and blow-up property of solutions to an initial and boundary value problem for a nonlinear parabolic system with variable exponents is studied. Meanwhile, the blow-up property of solutions for a nonlinear hyperbolic system is also obtained.

Keywords: existence; blow-up; parabolic system; hyperbolic system; variable exponent

1 Introduction

In this paper, we first consider the initial and boundary value problem to the following nonlinear parabolic system with variable exponents:

$$\begin{cases} u_t = \Delta u + f_1(u, v), & (x, t) \in Q_T, \\ v_t = \Delta v + f_2(u, v), & (x, t) \in Q_T, \\ u(x, t) = 0, \quad v(x, t) = 0, & (x, t) \in S_T, \\ u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), & x \in \Omega, \end{cases} \quad (1.1)$$

where $\Omega \subset \mathbb{R}^N$ is a bounded domain with smooth boundary $\partial\Omega$ and $0 < T < \infty$, $Q_T = \Omega \times [0, T)$, S_T denotes the lateral boundary of the cylinder Q_T , and the source terms f_1, f_2 are in the form

$$f_1(u, v) = a_1(x)v^{p_1(x)} \quad \text{and} \quad f_2(u, v) = a_2(x)u^{p_2(x)},$$

or

$$f_1(u, v) = a_1(x) \int_{\Omega} v^{p_1(y)}(y, t) dy \quad \text{and} \quad f_2(u, v) = a_2(x) \int_{\Omega} u^{p_2(y)}(y, t) dy,$$

respectively, where p_1, p_2, a_1, a_2 are functions satisfying conditions (2.1) below.

In the case when p_1, p_2 are constants, system (1.1) provides a simple example of a reaction-diffusion system. It can be used as a model to describe heat propagation in a two-component combustible mixture. There have been many results about the existence,

boundedness and blow-up properties of the solutions; we refer the readers to the bibliography given in [1–7].

The motivation of this work is due to [2], where the following system of equations is studied.

$$\begin{cases} u_t - \Delta u = v^p, \\ v_t - \Delta v = u^q, \end{cases} \quad (1.2)$$

where $x \in \mathbb{R}^N$ ($N \geq 1$), $t > 0$, and p, q are positive numbers. The authors investigated the boundedness and blow-up of solutions to problem (1.2). Furthermore, the authors also studied the uniqueness and global existence of solutions (see [3]).

Besides, this work is also motivated by [8] in which the following problem is considered:

$$\begin{cases} u_t = \Delta u + f(x, u), & (x, t) \in \Omega \times [0, T], \\ u(x, 0) = u_0(x), & x \in \Omega, \\ u(x, t) = 0, & (x, t) \in \partial\Omega \times [0, T], \end{cases} \quad (1.3)$$

where $\Omega \in \mathbb{R}^n$ is a bounded domain with smooth boundary $\partial\Omega$, and the source term is of the form $f(x, u) = a(x)u^{p(x)}$ or $f(x, u) = a(x) \int_{\Omega} u^{q(y)}(y, t) dy$. The author studied the blow-up property of solutions for parabolic and hyperbolic problems. Parabolic problems with sources like the ones in (1.3) appear in several branches of applied mathematics, which can be used to model chemical reactions, heat transfer or population dynamics *etc.* We also refer the interested reader to [9–23] and the references therein.

We also study the following nonlinear hyperbolic system of equations:

$$\begin{cases} u_{tt} = \Delta u + f_1(u, v), & (x, t) \in Q_T, \\ v_{tt} = \Delta v + f_2(u, v), & (x, t) \in Q_T, \\ u(x, t) = 0, \quad v(x, t) = 0, & (x, t) \in S_T, \\ u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), & x \in \Omega, \\ u_t(x, 0) = u_1(x), \quad v_t(x, 0) = v_1(x), & x \in \Omega. \end{cases} \quad (1.4)$$

The aim of this paper is to extend the results in [2, 8] to the case of parabolic system (1.1) and hyperbolic system (1.4). As far as we know, this seems to be the first paper, where the blow-up phenomenon is studied with variable exponents for the initial and boundary value problem to some parabolic and hyperbolic systems. The main method of the proof is similar to that in [3, 8].

We conclude this introduction by describing the outline of this paper. Some preliminary results, including existence of solutions to problem (1.1), are gathered in Section 2. The blow-up property of solutions are stated and proved in Section 3. Finally, in Section 4, we prove the blow-up property of solutions for hyperbolic problem (1.4).

2 Existence of solutions

In this section, we first state some assumptions and definitions needed in the proof of our main result and then prove the existence of solutions.

Throughout the paper, we assume that the exponents $p_1(x), p_2(x) : \Omega \rightarrow (1, +\infty)$ and the continuous functions $a_1(x), a_2(x) : \Omega \rightarrow \mathbb{R}$ satisfy the following conditions:

$$\begin{aligned}
 1 < p_1^- &= \inf_{x \in \Omega} p_1(x) \leq p_1(x) \leq p_1^+ = \sup_{x \in \Omega} p_1(x) < +\infty, \\
 1 < p_2^- &= \inf_{x \in \Omega} p_2(x) \leq p_2(x) \leq p_2^+ = \sup_{x \in \Omega} p_2(x) < +\infty, \\
 0 < c_1 &\leq a_1(x) \leq C_1 < +\infty, \quad 0 < c_2 \leq a_2(x) \leq C_2 < +\infty.
 \end{aligned} \tag{2.1}$$

Definition 2.1 We say that the solution $(u(x, t), v(x, t))$ for problem (1.1) blows up in finite time if there exists an instant $T^* < \infty$ such that

$$\| (u, v) \| \rightarrow \infty \quad \text{as } t \rightarrow T^*,$$

where

$$\| (u, v) \| = \sup_{t \in [0, T]} \{ \| u(\cdot, t) \|_\infty + \| v(\cdot, t) \|_\infty \}.$$

Our first result here is the following.

Theorem 2.1 Let $\Omega \subset \mathbb{R}^N$ be a bounded smooth domain, $p_1(x), p_2(x), a_1(x), a_2(x)$ satisfy the conditions in (2.1), and assume that $u_0(x)$ and $v_0(x)$ are nonnegative, continuous and bounded. Then there exists a $T^0, 0 < T^0 \leq \infty$, such that problem (1.1) has a nonnegative and bounded solution (u, v) in Q_{T^0} .

Proof We only prove the case when $f_1(u, v) = a_1(x)v^{p_1(x)}$ and $f_2(u, v) = a_2(x)u^{p_2(x)}$, and the proofs to the cases $f_1(u, v) = a_1(x) \int_\Omega v^{p_1(y)}(y, t) dy$ and $f_2(u, v) = a_2(x) \int_\Omega u^{p_2(y)}(y, t) dy$ are similar.

Let us consider the equivalent systems of (1.1)

$$\begin{cases}
 u(x, t) = \int_\Omega g(x, y, t) u_0(y) dy + \int_0^t \int_\Omega g(x, y, t-s) a_1(y) v^{p_1(y)} dy ds, \\
 v(x, t) = \int_\Omega g(x, y, t) v_0(y) dy + \int_0^t \int_\Omega g(x, y, t-s) a_2(y) u^{p_2(y)} dy ds,
 \end{cases}$$

where $g(x, y, t)$ is the corresponding Green function. Then the existence and uniqueness of solutions for a given $(u_0(x), v_0(x))$ could be obtained by a fixed point argument.

We introduce the following iteration scheme:

$$\begin{aligned}
 u_1(x, t) &= 0, \quad v_1(x, t) = 0, \\
 u_{n+1}(x, t) &= \int_\Omega g(x, y, t) u_0(y) dy + \int_0^t \int_\Omega g(x, y, t-s) a_1(y) v_n^{p_1(y)} dy ds, \\
 v_{n+1}(x, t) &= \int_\Omega g(x, y, t) v_0(y) dy + \int_0^t \int_\Omega g(x, y, t-s) a_2(y) u_n^{p_2(y)} dy ds,
 \end{aligned}$$

and the convergence of the sequence $\{(u_n, v_n)\}$ follows by showing that

$$\begin{cases}
 \Phi_1(v) = \int_0^t \int_\Omega g(x, y, t-s) a_1(y) v_n^{p_1(y)} dy ds, \\
 \Phi_2(u) = \int_0^t \int_\Omega g(x, y, t-s) a_2(y) u_n^{p_2(y)} dy ds
 \end{cases}$$

is a contraction in the set E_T to be defined below.

Now, we define

$$\Psi(u, v) = (\Phi_1(v), \Phi_2(u)),$$

where

$$\begin{aligned} \Phi_1(v) &= \int_0^t \int_{\Omega} g(x, y, t-s) v_n^{p_1(y)} dy ds, \\ \Phi_2(u) &= \int_0^t \int_{\Omega} g(x, y, t-s) u_n^{p_2(y)} dy ds. \end{aligned}$$

We denote

$$\Psi(u, v) - \Psi(w, z) = (\Phi_1(v) - \Phi_1(z), \Phi_2(u) - \Phi_2(w)),$$

and for arbitrary $T > 0$, define the set

$$E_T = \{C^{1,2}(\Omega_T) \cap C(\overline{\Omega_T}) \mid \|(u, v)\| \leq M\},$$

where $\Omega_T = \Omega \times [0, T]$, $M > \|(u_0(x), v_0(x))\|$ is a fixed positive constant.

We claim that Ψ is a contraction on E_T . In fact, for any $x \in \Omega$ fixed, we have

$$\begin{aligned} \xi_1^{p_1(x)} - \eta_1^{p_1(x)} &= p_1(x) w_1^{p_1(x)-1} (\xi_1 - \eta_1), \quad \text{where } w_1 = s_1(x)\xi_1 + (1-s_1(x))\eta_1, s_1(x) \in (0, 1), \\ \xi_2^{p_2(x)} - \eta_2^{p_2(x)} &= p_2(x) w_2^{p_2(x)-1} (\xi_2 - \eta_2), \\ &\text{where } w_2 = s_2(x)\xi_2 + (1-s_2(x))\eta_2, s_2(x) \in (0, 1), \end{aligned}$$

and we always have

$$\|p_i(x) w_i^{p_i(x)-1} (\xi_i - \eta_i)\| \leq p_i^+ (2M)^{p_i^+-1} \|\xi_i - \eta_i\|_{\infty}, \quad i = 1, 2. \tag{2.2}$$

Now, we define

$$\mu(t) = \sup_{x \in \overline{\Omega}, 0 \leq \tau < t} \int_0^{\tau} \int_{\Omega} g(x, y, t-s) dy ds.$$

It is obvious that $\mu(t) \rightarrow 0$ when $t \rightarrow 0^+$.

Then, by using inequality (2.2), we get

$$\begin{aligned} &\|\Phi_1(v) - \Phi_1(z)\|_{\infty} + \|\Phi_2(u) - \Phi_2(w)\|_{\infty} \\ &\leq \left\| \int_0^t \int_{\Omega} a_1(y) g(x, y, t-s) (v_n^{p_1(y)} - z_n^{p_1(y)}) dy ds \right\|_{\infty} \\ &\quad + \left\| \int_0^t \int_{\Omega} a_2(y) g(x, y, t-s) (u_n^{p_2(y)} - w_n^{p_2(y)}) dy ds \right\|_{\infty} \\ &\leq (2\mu(t))(2M)^{\max\{p_1^+, p_2^+\}-1} (C_1 + C_2) (\|p_1\|_{\infty} + \|p_2\|_{\infty}) (\|v - z\|_{\infty} + \|u - w\|_{\infty}) \\ &\leq (2\mu(t))(2M)^{\max\{p_1^+, p_2^+\}-1} (C_1 + C_2) (\|p_1\|_{\infty} + \|p_2\|_{\infty}) \sup_{t \in [0, T]} (\|v - z\|_{\infty} + \|u - w\|_{\infty}) \\ &= (2\mu(t))(2M)^{\max\{p_1^+, p_2^+\}-1} (C_1 + C_2) (\|p_1\|_{\infty} + \|p_2\|_{\infty}) \|(v - z, u - w)\|. \end{aligned}$$

Hence, for sufficiently small t , we have

$$\begin{aligned} & \| \Psi(u, v) - \Psi(w, z) \| \\ &= \| (\Phi_1(v) - \Phi_1(z), \Phi_2(u) - \Phi_2(w)) \| \\ &\leq \sup_{t \in [0, T]} \{ \| \Phi_1(v) - \Phi_1(z) \|_\infty + \| \Phi_2(u) - \Phi_2(w) \|_\infty \} \\ &\leq (2\mu(t))(2M)^{\max\{p_1^+, p_2^+\}-1} (C_1 + C_2) (\|p_1\|_\infty + \|p_2\|_\infty) \| (v - z, u - w) \| \\ &< \theta \| (v - z, u - w) \|, \end{aligned}$$

where $\theta < 1$ is a constant. Then Ψ is a strict contraction. □

3 Blow-up of solutions

In this section, we study the blow-up property of the solutions to problem (1.1). We need the following lemma.

Lemma 3.1 *Let $y(t)$ be a solution of*

$$y'(t) \geq -\lambda y(t) + ay^r(t) - C,$$

where $\lambda > 0$, $a > 0$, $r > 1$ and $C > 0$ are given constants. Then, there exists a constant $K > 0$ such that if $y(0) \geq K$, then $y(t)$ cannot be globally defined; in fact,

$$y(t) \geq \left(y(0)^{1-r} - \frac{a(r-1)}{2} t \right)^{-1/(r-1)}. \tag{3.1}$$

Proof It is sufficient to take $K > 0$ such that

$$-\lambda\eta + a\eta^r - C \geq \frac{a}{2}\eta^r, \quad \forall \eta \geq K.$$

Hence, we have

$$y'(t) \geq \frac{a}{2}y(t)^r. \tag{3.2}$$

By a direct integration to (3.2), then we get immediately (3.1), which gives an upper bound for the blow-up time $t^* = \frac{2y^{1-r}(0)}{a(r-1)}$. □

The next theorem gives the main result of this section.

Theorem 3.1 *Let $\Omega \subset \mathbb{R}^N$ be a bounded smooth domain, and let (u, v) be a positive solution of problem (1.1), with $p_1(x), p_2(x), a_1(x), a_2(x)$ satisfying conditions in (2.1). Then any solutions of problem (1.1) will blow up at finite time T^* if the initial datum $(u_0(x), v_0(x))$ satisfies*

$$\int_{\Omega} (u_0(x) + v_0(x))\varphi > C,$$

where $\varphi > 0$ is the first eigenfunction of the homogeneous Dirichlet Laplacian on Ω and $C > 0$ is a constant depending only on the domain Ω and the bounds C_1, C_2 given in condition (2.1).

Proof Let λ_1 be the first eigenvalue of

$$-\Delta \varphi = \lambda \varphi, \quad x \in \Omega,$$

with the homogeneous Dirichlet boundary condition, and let φ be a positive function satisfying

$$\int_{\Omega} \varphi \, dx = 1.$$

We introduce the function $\eta(t) = \int_{\Omega} (u + v)\varphi$. First of all, we consider the case $f_1(u, v) = a_1(x)v^{p_1(x)}, f_2(u, v) = a_2(x)u^{p_2(x)}$. Then

$$\begin{aligned} \eta'(t) &= \int_{\Omega} (u_t + v_t)\varphi \\ &= \int_{\Omega} (\Delta u + \Delta v)\varphi + \int_{\Omega} (a_1(x)v^{p_1(x)} + a_2(x)u^{p_2(x)})\varphi \\ &= -\lambda_1 \eta + \int_{\Omega} (a_1(x)v^{p_1(x)} + a_2(x)u^{p_2(x)})\varphi. \end{aligned}$$

We now deal with the term $\int_{\Omega} (a_1(x)v^{p_1(x)} + a_2(x)u^{p_2(x)})\varphi$. For each $t > 0$, we divide Ω into the following four sets:

$$\begin{aligned} \Omega_{11} &= \{x \in \Omega : v(x, t) < 1, u(x, t) < 1\}, & \Omega_{12} &= \{x \in \Omega : v(x, t) < 1, u(x, t) \geq 1\}, \\ \Omega_{21} &= \{x \in \Omega : v(x, t) \geq 1, u(x, t) < 1\}, & \Omega_{22} &= \{x \in \Omega : v(x, t) \geq 1, u(x, t) \geq 1\}. \end{aligned}$$

Then we have

$$\begin{aligned} &\int_{\Omega} (a_1(x)v^{p_1(x)} + a_2(x)u^{p_2(x)})\varphi \\ &= \int_{\Omega_{11}} (a_1(x)v^{p_1(x)} + a_2(x)u^{p_2(x)})\varphi + \int_{\Omega_{12}} (a_1(x)v^{p_1(x)} + a_2(x)u^{p_2(x)})\varphi \\ &\quad + \int_{\Omega_{21}} a_1(x)(v^{p_1(x)} + a_2(x)u^{p_2(x)})\varphi + \int_{\Omega_{22}} (a_1(x)v^{p_1(x)} + a_2(x)u^{p_2(x)})\varphi \\ &\geq \int_{\Omega_{12}} a_2(x)u^p \varphi + \int_{\Omega_{21}} a_1(x)v^p \varphi + \int_{\Omega_{22}} (a_1(x)v^p + a_2(x)u^p)\varphi \\ &\geq \int_{\Omega_{12}} a_2(x)u^p \varphi + \int_{\Omega_{21}} a_1(x)v^p \varphi \\ &\quad + \left(\int_{\Omega} (a_1(x)v^p + a_2(x)u^p)\varphi - \int_{\Omega_{12}} (a_1(x)v^p + a_2(x)u^p)\varphi \right. \\ &\quad \left. - \int_{\Omega_{21}} (a_1(x)v^p + a_2(x)u^p)\varphi - \int_{\Omega_{11}} (a_1(x)v^p + a_2(x)u^p)\varphi \right) \end{aligned}$$

$$\begin{aligned} &\geq \gamma \int_{\Omega} (v^p + u^p)\varphi - \Gamma \int_{\Omega_{12}} v^p\varphi - \Gamma \int_{\Omega_{21}} u^p\varphi - \Gamma \int_{\Omega_{11}} (v^p + u^p)\varphi \\ &\geq \gamma \int_{\Omega} (v^p + u^p)\varphi - 4\Gamma \int_{\Omega} \varphi, \end{aligned}$$

where $p = \min\{p_1^-, p_2^-\}$ and $\gamma = \min\{c_1, c_2\}$, $\Gamma = \max\{C_1, C_2\}$.

From the convex property of the function $f(w) = w^r$, $r > 1$ and Jensen's inequality, we obtain

$$\begin{aligned} \int_{\Omega} (a_1(x)v^{p_1(x)} + a_2(x)u^{p_2(x)})\varphi &\geq \gamma \int_{\Omega} (v^p + u^p)\varphi - 4\Gamma \int_{\Omega} \varphi \\ &\geq \gamma \int_{\Omega} 2\left(\frac{v+u}{2}\right)^p \varphi - 4\Gamma \int_{\Omega} \varphi \\ &\geq \frac{\gamma}{2^{p-1}}\eta^p(t) - 4\Gamma. \end{aligned}$$

Then we get

$$\eta'(t) \geq -\lambda_1\eta(t) + \frac{\gamma}{2^{p-1}}\eta^p(t) - 4\Gamma.$$

Note that

$$\eta(t) = \int_{\Omega} (u + v)\varphi \leq (\|u(\cdot, t)\|_{L^\infty(\Omega)} + \|v(\cdot, t)\|_{L^\infty(\Omega)}) \int_{\Omega} \varphi \leq \|(u, v)\|.$$

Hence, for $\eta(0)$ big enough, the result follows from Lemma 3.1.

Next, we state briefly the proof to the theorem in the case $f_1(u, v) = a_1(x) \int_{\Omega} v^{p_1(y)}(y, t) dy$ and $f_2(u, v) = a_2(x) \int_{\Omega} u^{p_2(y)}(y, t) dy$. We repeat the previous argument under defining $\eta(t) = \int_{\Omega} (u + v)\varphi$, and we obtain in much the same way

$$\eta'(t) \geq -\lambda_1\eta(t) + \int_{\Omega} \left(a_1(x) \int_{\Omega} v^{p_1(y)}(y, t) dy + a_2(x) \int_{\Omega} u^{p_2(y)}(y, t) dy \right) \varphi(x) dx.$$

In view of the property of φ , we get

$$\begin{aligned} &\int_{\Omega} \left(a_1(x) \int_{\Omega} v^{p_1(y)}(y, t) dy + a_2(x) \int_{\Omega} u^{p_2(y)}(y, t) dy \right) \varphi(x) dx \\ &= \int_{\Omega} v^{p_1(y)}(y, t) dy \left(\int_{\Omega} a_1(x)\varphi(x) dx \right) + \int_{\Omega} u^{p_2(y)}(y, t) dy \left(\int_{\Omega} a_2(x)\varphi(x) dx \right) \\ &\geq c_1 \int_{\Omega} v^{p_1(y)}(y, t) \frac{\varphi(y)}{\|\varphi\|_\infty} dy + c_2 \int_{\Omega} u^{p_2(y)}(y, t) \frac{\varphi(y)}{\|\varphi\|_\infty} dy \\ &\geq \gamma \int_{\Omega} (v^{p_1(y)}(y, t) + u^{p_2(y)}(y, t)) \frac{\varphi(y)}{\|\varphi\|_\infty} dy. \end{aligned}$$

According to the convex property of the function $f(w) = w^r$, $r > 1$, and by using Jensen's inequality, by considering again Ω_{11} , Ω_{12} , Ω_{21} , Ω_{22} as before, we obtain

$$\gamma \int_{\Omega} (v^{p_1(y)}(y, t) + u^{p_2(y)}(y, t)) \frac{\varphi(y)}{\|\varphi\|_\infty} dy \geq \gamma_0\eta^p(t) - 4\Gamma|\Omega|,$$

where γ_0 depends only on γ, p and $\|\varphi\|_\infty, |\Omega|$ denotes the measure of Ω . Hence,

$$\eta'(t) \geq -\lambda_1 \eta(t) + \gamma_0 \eta^p(t) - 4\Gamma |\Omega|.$$

By Lemma 3.1, the proof is complete. □

4 Blow-up of solutions for a hyperbolic system

Lemma 4.1 [15] *Let $y(t) \in C^2$ satisfying*

$$y''(t) \geq h(y(t)),$$

$y(0) = a > 0, y'(0) = b > 0$, and $h(s) \geq 0$ for all $s \geq a$. Then $y'(t) > 0$ whenever y exists; and

$$t \leq \int_a^{y(t)} \left(b^2 + 2 \int_a^s h(x) dx \right)^{-1/2} ds. \tag{4.1}$$

Now, let us study the following problem:

$$\begin{cases} u_{tt} = \Delta u + f_1(u, v), & (x, t) \in Q_T, \\ v_{tt} = \Delta v + f_2(u, v), & (x, t) \in Q_T, \\ u(x, t) = 0, \quad v(x, t) = 0, & (x, t) \in S_T, \\ u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), & x \in \Omega, \\ u_t(x, 0) = u_1(x), \quad v_t(x, 0) = v_1(x), & x \in \Omega, \end{cases} \tag{4.2}$$

where $u_0(x), v_0(x), u_1(x), v_1(x) \geq 0$ and they are not identically zero, and $f_1(u, v), f_2(u, v)$ as above respectively.

Theorem 4.1 *Let $(u, v) \in C^2 \times C^2$ be a solution of problem (4.2), and let the conditions in (2.1) hold. Then there exist sufficiently large initial data u_0, v_0, u_1, v_1 such that any solutions of problem (4.1) blew up at finite time T^* .*

Proof Let (λ_1, φ) be the first eigenvalue and eigenfunction of Laplacian in Ω with homogeneous Dirichlet boundary conditions as before. We assume that $f_1(u, v) = a_1(x)v^{p_1(x)}$, $f_2(u, v) = a_2(x)u^{p_2(x)}$, the other is similar. We also define the function $\eta(t) = \int_\Omega (u + v)\varphi$, so we have

$$\begin{aligned} \eta''(t) &= \int_\Omega (u_{tt} + v_{tt})\varphi \\ &= \int_\Omega (\Delta u + \Delta v)\varphi + \int_\Omega (a_1(x)v^{p_1(x)} + a_2(x)u^{p_2(x)})\varphi \\ &= -\lambda_1 \eta + \int_\Omega (a_1(x)v^{p_1(x)} + a_2(x)u^{p_2(x)})\varphi. \end{aligned}$$

The term $\int_\Omega (a_1(x)v^{p_1(x)} + a_2(x)u^{p_2(x)})\varphi$ is dealt with as before, then we get

$$\int_\Omega (a_1(x)v^{p_1(x)} + a_2(x)u^{p_2(x)})\varphi \geq \gamma \int_\Omega (v^p + u^p)\varphi - 4\Gamma \int_\Omega \varphi.$$

By virtue of the convex property of the function $f(w) = w^r$, $r > 1$, and Jensen's inequality, we still obtain

$$\gamma \int_{\Omega} (v^p + u^p) \varphi - 4\Gamma \int_{\Omega} \varphi \geq \gamma \int_{\Omega} 2 \left(\frac{v+u}{2} \right)^p \varphi - 4\Gamma \int_{\Omega} \varphi \geq \frac{\gamma}{2^{p-1}} \eta^p(t) - 4\Gamma.$$

Then we have

$$\eta''(t) \geq -\lambda_1 \eta(t) + \frac{\gamma}{2^{p-1}} \eta^p(t) - 4\Gamma.$$

Now, we can apply Lemma (4.1) for $a = \eta(0)$, $b = \eta'(0)$ large enough such that $-\lambda_1 \eta(t) + \frac{\gamma}{2^{p-1}} \eta^p(t) - 4\Gamma > 0$, and note that

$$a = \eta(0) = \int_{\Omega} (u_0 + v_0) \varphi,$$

$$b = \eta'(0) = \int_{\Omega} (u_1 + v_1) \varphi.$$

Moreover,

$$\eta(t) = \int_{\Omega} (u + v) \varphi \leq (\|u(\cdot, t)\|_{L^\infty(\Omega)} + \|v(\cdot, t)\|_{L^\infty(\Omega)}) \int_{\Omega} \varphi \leq \|(u, v)\|.$$

Hence, (u, v) blows up before the maximal time of existence defined in inequality (4.1) is reached. □

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

YG performed the calculations and drafted the manuscript. WG supervised and participated in the design of the study and modified the draft versions. All authors read and approved the final manuscript.

Acknowledgements

Supported by NSFC (11271154) and by Key Lab of Symbolic Computation and Knowledge Engineering of Ministry of Education and by the 985 program of Jilin University.

Received: 15 February 2013 Accepted: 18 March 2013 Published: 5 April 2013

References

- Chen, Y, Levine, S, Rao, M: Variable exponent, linear growth functions in image restoration. *SIAM J. Appl. Math.* **66**, 1383-1406 (2006)
- Escobedo, M, Herrero, MA: Boundedness and blow up for a semilinear reaction-diffusion system. *J. Differ. Equ.* **89**, 176-202 (1991)
- Escobedo, M, Herrero, MA: A semilinear parabolic system in a bounded domain. *Ann. Mat. Pura Appl.* **CLXV**, 315-336 (1998)
- Friedman, A, Giga, Y: A single point blow up for solutions of nonlinear parabolic systems. *J. Fac. Sci. Univ. Tokyo Sect. I* **34**(1), 65-79 (1987)
- Galaktionov, VA, Kurdyumov, SP, Samarskii, AA: A parabolic system of quasilinear equations I. *Differ. Equ.* **19**(12), 2133-2143 (1983)
- Galaktionov, VA, Vázquez, JL: A Stability Technique for Evolution Partial Differential Equations. *Progress in Nonlinear Differential Equations and Their Applications*, vol. 56. Birkhäuser, Boston (2004)
- Kufner, A, Oldrich, J, Fucik, S: *Function Space*. Kluwer Academic, Dordrecht (1977)
- Pinasco, JP: Blow-up for parabolic and hyperbolic problems with variable exponents. *Nonlinear Anal.* **71**, 1094-1099 (2009)
- Andreu-Vailló, F, Caselles, V, Mazón, JM: *Parabolic Quasilinear Equations Minimizing Linear Growth Functions*. *Progress in Mathematics*, vol. 223. Birkhäuser, Basel (2004)
- Antontsev, SN, Shmarev, SI: Anisotropic parabolic equations with variable nonlinearity. *CMAF, University of Lisbon, Portugal* **013**, 1-34 (2007)

11. Antontsev, SN, Shmarev, SI: Blow-up of solutions to parabolic equations with nonstandard growth conditions. *CMAF, University of Lisbon, Portugal* **02**, 1-16 (2009)
12. Antontsev, SN, Shmarev, SI: Parabolic equations with anisotropic nonstandard growth conditions. *Int. Ser. Numer. Math.* **154**, 33-44 (2007)
13. Antontsev, SN, Shmarev, S: Blow-up of solutions to parabolic equations with nonstandard growth conditions. *J. Comput. Appl. Math.* **234**, 2633-2645 (2010)
14. Erdem, D: Blow-up of solutions to quasilinear parabolic equations. *Appl. Math. Lett.* **12**, 65-69 (1999)
15. Glassey, RT: Blow-up theorems for nonlinear wave equations. *Math. Z.* **132**, 183-203 (1973)
16. Kalashnikov, AS: Some problems of the qualitative theory of nonlinear degenerate second-order parabolic equations. *Russ. Math. Surv.* **42**(2), 169-222 (1987)
17. Levine, HA: Some nonexistence and instability theorems for solutions of formally parabolic equations of the form $Pu_t = -Au + F(u)$. *Arch. Ration. Mech. Anal.* **51**, 371-386 (1973)
18. Levine, HA, Payne, LE: Nonexistence of global weak solutions for classes of nonlinear wave and parabolic equations. *J. Math. Anal. Appl.* **55**, 329-334 (1976)
19. Lian, SZ, Gao, WJ, Cao, CL, Yuan, HJ: Study of the solutions to a model porous medium equation with variable exponents of nonlinearity. *J. Math. Anal. Appl.* **342**, 27-38 (2008)
20. Ruzicka, M: *Electrorheological Fluids: Modelling and Mathematical Theory*. Lecture Notes in Math., vol. 1748. Springer, Berlin (2000)
21. Simon, J: Compact sets in the space $L^p(0, T; B)$. *Ann. Mat. Pura Appl.* **4**(146), 65-96 (1987)
22. Tsutsumi, M: Existence and nonexistence of global solutions for nonlinear parabolic equations. *Publ. Res. Inst. Math. Sci.* **8**, 211-229 (1972)
23. Zhao, JN: Existence and nonexistence of solutions for $u_t = \operatorname{div}(|\nabla u|^{p-2} \nabla u) + f(\nabla u, u, x, t)$. *J. Math. Anal. Appl.* **172**, 130-146 (1993)

doi:10.1186/1687-2770-2013-76

Cite this article as: Gao and Gao: **Study of solutions to an initial and boundary value problem for certain systems with variable exponents.** *Boundary Value Problems* 2013 **2013**:76.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
