

RESEARCH

Open Access

Nonlocal boundary value hyperbolic problems involving integral conditions

Allaberen Ashyralyev^{1,2} and Necmettin Aggez^{1*}

*Correspondence:

naggez@fatih.edu.tr

¹ Department of Mathematics, Fatih University, Istanbul, 34500, Turkey
Full list of author information is available at the end of the article

Abstract

Stability estimates for the solution of the nonlocal boundary value problem with two integral conditions for hyperbolic equations in a Hilbert space H are established. In applications, stability estimates for the solution of the nonlocal boundary value problems for hyperbolic equations are obtained.

MSC: 35L10

Keywords: hyperbolic equation; stability; nonlocal boundary value problems

1 Introduction

It is well known that nonlocal boundary value problems with integral conditions are widely used for thermo-elasticity, chemical engineering, heat conduction, and plasma physics [1–4]. Some problems arising in dynamics of ground waters are defined as hyperbolic equations with nonlocal conditions [5] and [6]. The authors of [7] investigate nonclassical problems for multidimensional hyperbolic equation with integral boundary conditions and the uniqueness of classical solution. In [8] a linear second-order hyperbolic equation with forcing and integral constraints on the solution is converted to a nonlocal hyperbolic problem. Using the Riesz representation theorem and the Schauder fixed point theorem, existence and uniqueness of a generalized solution are proved. The solutions of hyperbolic equations with nonlocal integral conditions were investigated in [9–16]. The method of operators as a tool for investigation of the solution to hyperbolic equations in Hilbert and Banach spaces has been used extensively in [17–29].

In [30] the nonlocal boundary value problem

$$\begin{cases} \frac{d^2 u(t)}{dt^2} + Au(t) = f(t), & 0 \leq t \leq 1, \\ u(0) = \sum_{r=1}^n \alpha_r u(\lambda_r) + \varphi, & u_t(0) = \sum_{r=1}^n \beta_r u_t(\lambda_r) + \psi, \\ 0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n \leq 1 \end{cases}$$

was investigated. Stability estimates for the solution of the problem were established. First order of accuracy difference schemes for the approximate solutions of the problem were presented. Stability estimates for the solution of these difference schemes were established. Theoretical statements were supported by numerical examples.

In the present paper, we consider the nonlocal boundary value problem with integral conditions

$$\begin{cases} \frac{d^2 u(t)}{dt^2} + Au(t) = f(t), & 0 \leq t \leq 1, \\ u(0) = \int_0^1 \alpha(\rho)u(\rho) d\rho + \varphi, \\ u_t(0) = \int_0^1 \beta(\rho)u_t(\rho) d\rho + \psi \end{cases} \quad (1)$$

in a Hilbert space H with a self-adjoint positive definite operator A . We are interested in studying the stability of solutions of problem (1) under the assumption

$$\left| 1 + \int_0^1 \alpha(s)\beta(s) ds \right| > \int_0^1 (|\alpha(s)| + |\beta(s)|) ds. \quad (2)$$

As in [30], a function $u(t)$ is called a *solution* of problem (1) if the following conditions are satisfied:

- (i) $u(t)$ is twice continuously differentiable on the interval $(0, 1)$ and continuously differentiable on the segment $[0, 1]$.
- (ii) The element $u(t)$ belongs to $D(A)$ for all $t \in [0, 1]$, and the function $Au(t)$ is continuous on the segment $[0, 1]$.
- (iii) $u(t)$ satisfies the equation and nonlocal boundary conditions (1).

2 The main theorem

Let H be a Hilbert space, A be a positive definite self-adjoint operator with $A \geq \delta I$, where $\delta > \delta_0 > 0$. Throughout this paper, $\{c(t), t \geq 0\}$ is a strongly continuous cosine operator-function defined by

$$c(t) = \frac{e^{itA^{1/2}} + e^{-itA^{1/2}}}{2}.$$

Then, from the definition of sine operator-function $s(t)$,

$$s(t)u = \int_0^t c(\rho)u d\rho$$

it follows that

$$s(t) = A^{-1/2} \frac{e^{itA^{1/2}} - e^{-itA^{1/2}}}{2i}.$$

For the theory of cosine operator-function we refer to [23] and [24].

Lemma 2.1 *The following estimates hold:*

$$\|c(t)\|_{H \rightarrow H} \leq 1, \quad \|A^{1/2}s(t)\|_{H \rightarrow H} \leq 1. \quad (3)$$

Lemma 2.2 *Suppose that assumption (2) holds. Then the operator T ,*

$$T = \left(1 + \int_0^1 \int_0^1 \alpha(s)\beta(s) ds ds \right) I - \int_0^1 (\beta(s) + \alpha(s))c(s) ds,$$

has the inverse

$$T^{-1} = \left[\left(1 + \int_0^1 \int_0^1 \alpha(s)\beta(s) ds ds \right) I - \int_0^1 (\beta(s) + \alpha(s))c(s) ds \right]^{-1}$$

and the following estimate is satisfied:

$$\|T^{-1}\|_{H \rightarrow H} \leq \frac{1}{|1 + \int_0^1 \alpha(s)\beta(s) ds| - \int_0^1 (|\alpha(s)| + |\beta(s)|) ds}. \tag{4}$$

Proof Applying the triangle inequality and estimates (3), we obtain

$$\begin{aligned} & \left\| \left(1 + \int_0^1 \int_0^1 \alpha(s)\beta(s) ds ds \right) I - \int_0^1 (\beta(s) + \alpha(s))c(s) ds \right\|_{H \rightarrow H} \\ & \geq \left| 1 + \int_0^1 \alpha(s)\beta(s) ds \right| - \left\| \int_0^1 (\beta(s) + \alpha(s))c(s) ds \right\|_{H \rightarrow H} \\ & \geq \left| 1 + \int_0^1 \alpha(s)\beta(s) ds \right| - \int_0^1 (|\alpha(s)| + |\beta(s)|) \|c(s)\|_{H \rightarrow H} ds \\ & \geq \left| 1 + \int_0^1 \alpha(s)\beta(s) ds \right| - \int_0^1 (|\alpha(s)| + |\beta(s)|) ds > 0. \end{aligned}$$

Estimate (4) follows from this estimate. Lemma 2.2 is proved. □

Now, we will obtain the formula for the solution of problem (1) for $\varphi \in D(A)$ and $\psi \in D(A^{1/2})$. It is clear that [23] the initial value problem

$$\frac{d^2u}{dt^2} + Au(t) = f(t), \quad 0 < t < 1, \quad u(0) = u_0, \quad u'(0) = u'_0 \tag{5}$$

has a unique solution,

$$u(t) = c(t)u_0 + s(t)u'_0 + \int_0^t s(t-\lambda)f(\lambda) d\lambda, \tag{6}$$

where the function $f(t)$ is not only continuous but also continuously differentiable on $[0, 1]$, $u_0 \in D(A)$ and $u'_0 \in D(A^{1/2})$.

Using (6) and the nonlocal boundary condition

$$u_0 = \int_0^1 \alpha(y)u(y) dy + \varphi,$$

we get

$$u_0 = \int_0^1 \alpha(y) \left(c(y)u_0 + s(y)u'_0 + \int_0^y s(y-\lambda)f(\lambda) d\lambda \right) dy + \varphi.$$

Then

$$\left(I - \int_0^1 \alpha(y)c(y) dy \right) u_0 - \int_0^1 \alpha(y)s(y) dy u'_0 = \int_0^1 \alpha(y) \int_0^y s(y-\lambda)f(\lambda) d\lambda dy + \varphi. \tag{7}$$

Differentiating both sides of (6), we obtain

$$u'(t) = -As(t)u_0 + c(t)u'_0 + \int_0^t c(t-\lambda)f(\lambda) d\lambda.$$

Using this formula and the integral condition

$$u'_0 = \int_0^1 \beta(y)u'(y) dy + \psi,$$

we get

$$u'_0 = \int_0^1 \beta(y) \left(-As(y)u_0 + c(y)u'_0 + \int_0^y c(y-\lambda)f(\lambda) d\lambda \right) dy + \psi.$$

Thus,

$$\begin{aligned} & \int_0^1 \beta(y)As(y) dy u_0 + \left(I - \int_0^1 \beta(y)c(y) dy \right) u'_0 \\ & = \int_0^1 \beta(y) \int_0^y c(y-\lambda)f(\lambda) d\lambda dy + \psi. \end{aligned} \tag{8}$$

Now, we have a system of equations (7) and (8) for the solution of u_0 and u'_0 . Solving it, we get

$$\begin{aligned} u_0 = T^{-1} \left\{ \left[I - \int_0^1 \beta(y)c(y) dy \right] \left[\int_0^1 \alpha(y) \int_0^y s(y-\lambda)f(\lambda) d\lambda dy + \varphi \right] \right. \\ \left. + \int_0^1 \alpha(y)s(y) dy \left[\int_0^1 \beta(y) \int_0^y c(y-\lambda)f(\lambda) d\lambda dy + \psi \right] \right\} \end{aligned} \tag{9}$$

and

$$\begin{aligned} u'_0 = T^{-1} \left\{ \left[I - \int_0^1 \alpha(y)c(y) dy \right] \left[\int_0^1 \beta(y) \int_0^y c(y-\lambda)f(\lambda) d\lambda dy + \psi \right] \right. \\ \left. - \int_0^1 \beta(y)As(y) dy \left[\int_0^1 \alpha(y) \int_0^y s(y-\lambda)f(\lambda) d\lambda dy + \varphi \right] \right\}. \end{aligned} \tag{10}$$

Hence, for the solution of the nonlocal boundary value problem (1) we have (6), (9), and (10).

Theorem 2.1 *Suppose that $\varphi \in D(A)$, $\psi \in D(A^{1/2})$, and $f(t)$ is a continuously differentiable on $[0, 1]$; assumption (2) holds. Then there is a unique solution of problem (1) and the following stability inequalities:*

$$\max_{0 \leq t \leq 1} \|u(t)\|_H \leq M \left[\|\varphi\|_H + \|A^{-1/2}\psi\|_H + \int_0^1 \|A^{-1/2}f(\lambda)\|_H d\lambda \right], \tag{11}$$

$$\begin{aligned} & \max_{0 \leq t \leq 1} \left\| \frac{du(t)}{dt} \right\|_H + \max_{0 \leq t \leq 1} \|A^{1/2}u(t)\|_H \\ & \leq M \left[\|A^{1/2}\varphi\|_H + \|\psi\|_H + \int_0^1 \|f(\lambda)\|_H d\lambda \right], \end{aligned} \tag{12}$$

$$\begin{aligned} & \max_{0 \leq t \leq 1} \left\| \frac{d^2 u(t)}{dt^2} \right\|_H + \max_{0 \leq t \leq 1} \|Au(t)\|_H \\ & \leq M \left\{ \|A\varphi\|_H + \|A^{1/2}\psi\|_H + \|f(0)\|_H + \int_0^1 \|f'(\lambda)\|_H d\lambda \right\} \end{aligned} \tag{13}$$

are valid, where M does not depend on $f(t)$, $t \in [0, 1]$, φ , and ψ .

Proof We take the estimates

$$\max_{0 \leq t \leq 1} \|u(t)\|_H \leq \left[\|u(0)\|_H + \|A^{-1/2}u'(0)\|_H + \int_0^1 \|A^{-1/2}f(t)\|_H dt \right], \tag{14}$$

$$\begin{aligned} & \max_{0 \leq t \leq 1} \|u'(t)\|_H + \max_{0 \leq t \leq 1} \|A^{1/2}u(t)\|_H \\ & \leq M \left[\|A^{1/2}u(0)\|_H + \|u'(0)\|_H + \int_0^1 \|f(t)\|_H dt \right], \end{aligned} \tag{15}$$

$$\begin{aligned} & \max_{0 \leq t \leq 1} \left\| \frac{d^2 u(t)}{dt^2} \right\|_H + \max_{0 \leq t \leq 1} \|Au(t)\|_H + \max_{0 \leq t \leq 1} \|A^{1/2}u'(t)\|_H \\ & \leq M \left[\|Au(0)\|_H + \|A^{1/2}u'(0)\|_H + \|f(0)\|_H + \int_0^1 \|f'(t)\|_H dt \right] \end{aligned} \tag{16}$$

from [26] for the solution of problem (5). The proof of Theorem 2.1 is based on estimates (14), (15), (16), and the estimates for the norms of u_0 , $A^{-1/2}u'_0$, $A^{1/2}u_0$, u'_0 , Au_0 , $A^{1/2}u'_0$.

First of all, let us find an estimate for $\|u(0)\|_H$. By using (9) and estimates (3), (4), we obtain

$$\|u(0)\|_H \leq M_1 \left[\int_0^1 \|A^{-1/2}f(\lambda)\|_H d\lambda + \|\varphi\|_H + \|A^{-1/2}\psi\|_H \right]. \tag{17}$$

Applying $A^{-1/2}$ to (10), we get

$$\begin{aligned} A^{-1/2}u'_0 &= T^{-1} \left\{ \left[I - \int_0^1 \alpha(y)c(y) dy \right] \left[\int_0^1 \beta(y) \int_0^y c(y-\lambda)A^{-1/2}f(\lambda) d\lambda dy + A^{-1/2}\psi \right] \right. \\ & \quad \left. - \int_0^1 \beta(y)A^{1/2}s(y) dy \right. \\ & \quad \left. \times \left[\int_0^1 \alpha(y) \int_0^y A^{1/2}s(y-\lambda)A^{-1/2}f(\lambda) d\lambda dy + \varphi \right] \right\}. \end{aligned}$$

Using estimates (3), (4), we obtain

$$\|A^{-1/2}u'(0)\|_H \leq M_2 \left[\int_0^1 \|A^{-1/2}f(\lambda)\|_H d\lambda + \|A^{-1/2}\psi\|_H + \|\varphi\|_H \right]. \tag{18}$$

Thus, estimates (14), (17), and (18) yield estimate (11).

Second, applying operator $A^{1/2}$ to (9), we get

$$\begin{aligned} A^{1/2}u_0 &= T^{-1} \left\{ \left[I - \int_0^1 \beta(y)c(y) dy \right] \left[\int_0^1 \alpha(y) \int_0^y A^{1/2}s(y-\lambda)f(\lambda) d\lambda dy + A^{1/2}\varphi \right] \right. \\ & \quad \left. + \int_0^1 \alpha(y)A^{1/2}s(y) dy \left[\int_0^1 \beta(y) \int_0^y c(y-\lambda)f(\lambda) d\lambda dy + \psi \right] \right\}. \end{aligned}$$

Using estimates (3) and (4), we obtain

$$\|A^{1/2}u(0)\|_H \leq M_1 \left[\int_0^1 \|f(\lambda)\|_H d\lambda + \|A^{1/2}\varphi\|_H + \|\psi\|_H \right]. \tag{19}$$

Using (10), and estimates (3), (4), we get

$$\|u'(0)\|_H \leq M_2 \left[\int_0^1 \|f(\lambda)\|_H d\lambda + \|\psi\|_H + \|A^{1/2}\varphi\|_H \right]. \tag{20}$$

Then estimate (12) follows from estimates (15), (19), and (20).

Third, applying A to (9) and using Abel's formula, we have

$$\begin{aligned} Au_0 = T^{-1} & \left\{ \left[I - \int_0^1 \beta(y)c(y) dy \right] \right. \\ & \times \left[\int_0^1 \alpha(y) \left[f(y) - c(y)f(0) - \int_0^y c(y-\lambda)f'(\lambda) d\lambda \right] dy + A\varphi \right] \\ & + \int_0^1 \alpha(y)A^{1/2}s(y) dy \\ & \left. \times \left[\int_0^1 \beta(y) \left[A^{1/2}s(y)f(0) + \int_0^y A^{1/2}s(y-\lambda)f'(\lambda) d\lambda \right] dy + A^{1/2}\psi \right] \right\} \end{aligned}$$

and using estimates (3), (4), we get

$$\|Au(0)\|_H \leq M_3 \left[\|A\varphi\|_H + \|A^{1/2}\psi\|_H + \|f(0)\|_H + \int_0^1 \|f'(\lambda)\|_H d\lambda \right]. \tag{21}$$

In the same manner, applying $A^{1/2}$ to (10) and using Abel's formula, and estimates (3), (4), we obtain

$$\|A^{1/2}u'(0)\|_H \leq M_4 \left[\|A\varphi\|_H + \|A^{1/2}\psi\|_H + \|f(0)\|_H + \int_0^1 \|f'(\lambda)\|_H d\lambda \right]. \tag{22}$$

Thus, estimate (13) follows from estimates (16) and (21), and (22). □

3 Applications

Now, we consider the applications of Theorem 2.1. First, a nonlocal boundary value problem for a hyperbolic equation

$$\begin{cases} u_{tt} - (a(x)u_x)_x + \sigma u = f(t, x), & 0 < t < 1, 0 < x < 1, \\ u(0, x) = \int_0^1 \alpha(\rho)u(\rho, x) d\rho + \varphi(x), \\ u_t(0, x) = \int_0^1 \beta(\rho)u_t(\rho, x) d\rho + \psi(x), & 0 \leq x \leq 1, \\ u(t, 0) = u(t, 1), \quad u_x(t, 0) = u_x(t, 1), & 0 \leq t \leq 1 \end{cases} \tag{23}$$

under assumption (2) is considered. Problem (23) has a unique smooth solution $u(t, x)$ for (2), smooth functions $a(x) \geq a > 0$ ($x \in (0, 1)$), $a(0) = a(1)$, $\varphi(x)$, $\psi(x)$ ($x \in [0, 1]$) and $f(t, x)$ ($t, x \in [0, 1]$), σ a positive constant and under some conditions. This allows us to reduce problem (23) to nonlocal boundary value problem (1) in the Hilbert space $H = L_2[0, 1]$ with a self-adjoint positive definite operator A^x defined by (23).

Theorem 3.1 *For the solution of problem (23), we have the following stability inequalities:*

$$\max_{0 \leq t \leq 1} \|u_x(t, \cdot)\|_{L_2[0,1]} \leq M \left[\max_{0 \leq t \leq 1} \|f(t, \cdot)\|_{L_2[0,1]} + \|\varphi_x\|_{L_2[0,1]} + \|\psi\|_{L_2[0,1]} \right], \tag{24}$$

$$\begin{aligned} & \max_{0 \leq t \leq 1} \|u_{xx}(t, \cdot)\|_{L_2[0,1]} + \max_{0 \leq t \leq 1} \|u_{tt}(t, \cdot)\|_{L_2[0,1]} \\ & \leq M \left[\max_{0 \leq t \leq 1} \|f_t(t, \cdot)\|_{L_2[0,1]} + \|f(0, \cdot)\|_{L_2[0,1]} + \|\varphi_{xx}\|_{L_2[0,1]} + \|\psi_x\|_{L_2[0,1]} \right], \end{aligned} \tag{25}$$

where M does not depend on $\varphi(x)$, $\psi(x)$, and $f(t, x)$.

The proof of Theorem 3.1 is based on Theorem 2.1 and the symmetry properties of the space operator generated by problem (23).

Proof Problem (23) can be written in the abstract form

$$\begin{cases} \frac{d^2 u(t)}{dt^2} + Au(t) = f(t) & (0 \leq t \leq 1), \\ u(0) = \int_0^1 \alpha(\rho)u(\rho) d\rho + \varphi, \\ u_t(0) = \int_0^1 \beta(\rho)u_t(\rho) d\rho + \psi \end{cases} \tag{26}$$

in the Hilbert space $L_2[0, 1]$ of all square integrable functions defined on $[0, 1]$ with a self-adjoint positive definite operator $A = A^x$ defined by the formula

$$A^x u(x) = -(a(x)u_x)_x + \sigma u(x)$$

with the domain

$$D(A^x) = \left\{ \begin{array}{l} u(x) : u, u_x, (a(x)u_x)_x \in L_2[0, 1], \\ u(0) = u(1), u'(0) = u'(1) \end{array} \right\}.$$

Here, $f(t) = f(t, x)$ and $u(t) = u(t, x)$ are known and unknown abstract functions defined on $[0, 1]$ with the values in $H = L_2[0, 1]$. Therefore, estimates (24) and (25) follow from estimates (11), (12), and (13). Thus, Theorem 3.1 is proved. \square

Second, let Ω be the unit open cube in the m -dimensional Euclidean space $\mathbb{R}^m : \{x = (x_1, \dots, x_m) : 0 < x_j < 1, 1 \leq j \leq m\}$ with boundary $S, \overline{\Omega} = \Omega \cup S$. In $[0, 1] \times \Omega$, let us consider a boundary value problem for the multidimensional hyperbolic equation

$$\begin{cases} \frac{\partial^2 u(t,x)}{\partial t^2} - \sum_{r=1}^m (a_r(x)u_{x_r})_{x_r} + \sigma u(x) = f(t, x), \\ x = (x_1, \dots, x_m) \in \Omega, 0 < t < 1, \\ u(0, x) = \int_0^1 \alpha(\rho)u(\rho, x) d\rho + \varphi(x), \quad x \in \overline{\Omega}, \\ u_t(0, x) = \int_0^1 \beta(\rho)u_t(\rho, x) d\rho + \psi(x), \quad x \in \overline{\Omega}, \\ u(t, x) = 0, \quad x \in S, \end{cases} \tag{27}$$

under assumption (2). Here, $a_r(x)$ ($x \in \Omega$), $\varphi(x)$, $\psi(x)$ ($x \in \overline{\Omega}$) and $f(t, x)$, $t \in (0, 1)$, $x \in \Omega$ are given smooth functions and $a_r(x) \geq a > 0$.

Let us introduce the Hilbert space $L_2(\overline{\Omega})$ of all square integrable functions defined on $\overline{\Omega}$, equipped with the norm

$$\|f\|_{L_2(\overline{\Omega})} = \left\{ \int \cdots \int_{x \in \overline{\Omega}} |f(x)|^2 dx_1 \cdots dx_m \right\}^{\frac{1}{2}}.$$

Theorem 3.2 *For the solution of problem (27), the following stability inequalities hold:*

$$\begin{aligned} & \max_{0 \leq t \leq 1} \sum_{r=1}^m \|u_{x_r}(t, \cdot)\|_{L_2(\overline{\Omega})} \\ & \leq M \left[\max_{0 \leq t \leq 1} \|f(t, \cdot)\|_{L_2(\overline{\Omega})} + \sum_{r=1}^m \|\varphi_{x_r}\|_{L_2(\overline{\Omega})} + \|\psi\|_{L_2(\overline{\Omega})} \right], \end{aligned} \tag{28}$$

$$\begin{aligned} & \max_{0 \leq t \leq 1} \sum_{r=1}^m \|u_{x_r x_r}(t, \cdot)\|_{L_2(\overline{\Omega})} + \max_{0 \leq t \leq 1} \|u_{tt}(t, \cdot)\|_{L_2(\overline{\Omega})} \\ & \leq M \left[\max_{0 \leq t \leq 1} \|f_t(t, \cdot)\|_{L_2(\overline{\Omega})} + \|f(0, \cdot)\|_{L_2(\overline{\Omega})} \right. \\ & \quad \left. + \sum_{r=1}^m \|\varphi_{x_r x_r}\|_{L_2(\overline{\Omega})} + \sum_{r=1}^m \|\psi_{x_r}\|_{L_2(\overline{\Omega})} \right], \end{aligned} \tag{29}$$

where M does not depend on $\varphi(x)$, $\psi(x)$, and $f(t, x)$ ($t \in (0, 1)$, $x \in \Omega$).

Proof Problem (27) can be written in the abstract form (26) in the Hilbert space $L_2(\overline{\Omega})$ with a self-adjoint positive definite operator $A = A^x$ defined by the formula

$$A^x u(x) = - \sum_{r=1}^m (a_r(x) u_{x_r})_{x_r} + \sigma u(x)$$

with domain

$$D(A^x) = \{u(x) : u, u_{x_r}, (a_r(x) u_{x_r})_{x_r} \in L_2(\overline{\Omega}), 1 \leq r \leq m, u(x) = 0, x \in S\}.$$

Here, $f(t) = f(t, x)$ and $u(t) = u(t, x)$ are known and unknown abstract functions defined on $\overline{\Omega}$ with the values in $H = L_2(\overline{\Omega})$. So, estimates (28) and (29) follow from estimates (11), (12), (13), and the following theorem. \square

Theorem 3.3 [29] *For the solution of the elliptic differential problem*

$$A^x u(x) = \omega(x), \quad x \in \Omega,$$

$$u(x) = 0, \quad x \in S,$$

the following coercivity inequality holds:

$$\sum_{r=1}^m \|u_{x_r x_r}\|_{L_2(\overline{\Omega})} \leq M \|\omega\|_{L_2(\overline{\Omega})},$$

where M is independent of ω .

4 Conclusion

This work is devoted to the study of the stability of the nonlocal boundary value problem with integral conditions for hyperbolic equations. For the solution of nonlocal boundary problem (1) in a Hilbert space H with a self-adjoint positive definite operator A , Theorem 2.1 is established. Two applications of Theorem 2.1 are given. Of course, stable two-step difference schemes for approximate solution of problem (1) can be presented. The methods given above permit us to establish the stability of these difference schemes. Applying [31], we can give a numerical support of the theoretical results.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

Author details

¹Department of Mathematics, Fatih University, Istanbul, 34500, Turkey. ²Department of Mathematics, ITTU, Gerogly Street, Ashgabat, Turkmenistan.

Acknowledgements

This work is supported by the Scientific Research Fund of Fatih University (Project No: P50041203-B).

Received: 15 June 2014 Accepted: 19 August 2014 Published online: 25 September 2014

References

1. Shi, P: Weak solution to evolution problem with a nonlocal constraint. *SIAM J. Math. Anal.* **24**, 46-58 (1993)
2. Choi, YS, Chan, KY: A parabolic equation with nonlocal boundary conditions arising from electrochemistry. *Nonlinear Anal.* **18**, 317-331 (1992)
3. Cahlon, B, Kulkarni, DM, Shi, P: Stepwise stability for the heat equation with a nonlocal constraint. *SIAM J. Numer. Anal.* **32**, 571-593 (1995)
4. Samarski, AA: Some problems in the modern theory of differential equations. *Differ. Uravn.* **16**, 1221-1228 (1980)
5. Beilin, SA: Existence of solutions for one-dimensional wave equations with non-local conditions. *Electron. J. Differ. Equ.* **2001**, 76 (2001)
6. Dehghan, M: On the solution of an initial-boundary value problem that combines Neumann and integral condition for the wave equation. *Numer. Methods Partial Differ. Equ.* **21**, 24-40 (2005)
7. Avalishvili, G, Avalishvili, M, Gordeziani, D: On integral nonlocal boundary value problems for some partial differential equations. *Bull. Georgian Acad. Sci.* **5**(1), 31-37 (2011)
8. Pulkina, LS: A nonlocal problem with integral conditions for hyperbolic equations. *Electron. J. Differ. Equ.* **1999**, 45 (1999)
9. Ramezani, M, Dehghan, M, Razzaghi, M: Combined finite difference and spectral methods for the numerical solution of hyperbolic equation with an integral condition. *Numer. Methods Partial Differ. Equ.* **24**, 1-8 (2008)
10. Pulkina, LS: On solvability in L_2 of nonlocal problem with integral conditions for hyperbolic equations. *Differ. Uravn.* **36**(2), 279-280 (2000)
11. Mesloub, S, Bouziani, A: On a class of singular hyperbolic equation with a weighted integral condition. *Int. J. Math. Math. Sci.* **22**(3), 511-519 (1999)
12. Shakeri, F, Dehghan, M: The method of lines for solution of the one-dimensional wave equation subject to an integral conservation condition. *Comput. Math. Appl.* **56**, 2175-2188 (2008)
13. Dehghan, M, Saadatmandi, A: Variational iteration method for solving the wave equation subject to an integral conservation condition. *Chaos Solitons Fractals* **41**, 1448-1453 (2009)
14. Dehghan, M, Shokri, A: A meshless method for numerical solution of the one-dimensional wave equation with an integral condition using radial basis functions. *Numer. Algorithms* **52**, 461-477 (2009)
15. Dehghan, M, Saadatmandi, A: Numerical solution of the one-dimensional wave equation with an integral condition. *Numer. Methods Partial Differ. Equ.* **23**, 282-292 (2007)
16. Yousefi, SA, Barikbin, Z, Dehghan, M: Bernstein Ritz-Galerkin method for solving an initial-boundary value problem that combines Neumann and integral condition for the wave equation. *Numer. Methods Partial Differ. Equ.* **26**, 1236-1246 (2010)
17. Ashyralyev, A, Aggez, N: A note on the difference schemes of the nonlocal boundary value problems for hyperbolic equations. *Numer. Funct. Anal. Optim.* **25**(5-6), 1-24 (2004)
18. Sobolevskii, PE, Semenov, S: On some approach to investigation of singular hyperbolic equations. *Dokl. Akad. Nauk SSSR* **270**(1), 555-558 (1983)
19. Agarwal, RP, Bohner, M, Shakhmurov, VB: Maximal regular boundary value problems in Banach-valued weighted space. *Bound. Value Probl.* **1**, 9-42 (2004)
20. Ashyralyev, A, Sobolevskii, PE: *New Difference Schemes for Partial Differential Equations*. Birkhäuser, Basel (2004)
21. Ashyralyev, A, Sobolevskii, PE: *Well-Posedness of Parabolic Difference Equations*. Operator Theory Advances and Applications. Birkhäuser, Basel (1994)
22. Kostin, VA: On analytic semigroups and strongly continuous cosine functions. *Dokl. Akad. Nauk SSSR* **307**(4), 796-799 (1989)

23. Fattorini, HO: Second Order Linear Differential Equations in Banach Space. Notas de Matematica. North-Holland, Amsterdam (1985)
24. Piskarev, S, Shaw, Y: On certain operator families related to cosine operator function. Taiwan. J. Math. **1**(4), 3585-3592 (1997)
25. Guidetti, D, Karasözen, B, Piskarev, S: Approximation of abstract differential equations. J. Math. Sci. **122**(2), 3013-3054 (2004)
26. Ashyralyev, A, Sobolevskii, PE: A note on the difference schemes for hyperbolic equations. Abstr. Appl. Anal. **6**(2), 63-70 (2001)
27. Soltanov, H: A note on the Goursat problem for a multidimensional hyperbolic equation. Contemp. Anal. Appl. Math. **1**(2), 98-106 (2013)
28. Ashyralyev, A, Aggez, N: Finite difference method for hyperbolic equations with the nonlocal integral condition. Discrete Dyn. Nat. Soc. **2011**, 562385 (2011)
29. Sobolevskii, PE: Difference Methods for the Approximate Solution of Differential Equations. Izdat. Voronezh. Gosud. Univ., Voronezh (1975)
30. Ashyralyev, A, Yildirim, O: On multipoint nonlocal boundary value problems for hyperbolic differential and difference equations. Taiwan. J. Math. **14**(1), 165-194 (2010)
31. Ashyralyev, A, Aggez, N: On the solution of NBVP for multidimensional hyperbolic equations. Sci. World J. **2014**, 841602 (2014). doi:10.1155/2014/841602

doi:10.1186/s13661-014-0205-4

Cite this article as: Ashyralyev and Aggez: Nonlocal boundary value hyperbolic problems involving integral conditions. *Boundary Value Problems* 2014 **2014**:205.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
