

RESEARCH

Open Access



# Non-uniformly asymptotically linear fourth-order elliptic problems

Ruichang Pei<sup>1,2\*</sup> and Jihui Zhang<sup>2</sup>

\*Correspondence: prc211@163.com

<sup>1</sup>School of Mathematics and Statistics, Tianshui Normal University, Tianshui, 741001, P.R. China

<sup>2</sup>School of Mathematical Sciences, Nanjing Normal University, Nanjing, 210097, P.R. China

## Abstract

The existence of multiple solutions for a class of fourth-order elliptic equations with respect to the non-uniformly asymptotically linear conditions is established by using the minimax method and Morse theory.

**Keywords:** fourth-order elliptic boundary value problems; multiple solutions; mountain pass theorem; Morse theory

## 1 Introduction

Let  $\mathbf{H} = H^2(\Omega) \cap H_0^1(\Omega)$  be a Hilbert space equipped with the inner product

$$(u, v)_{\mathbf{H}} = \int_{\Omega} (\Delta u \Delta v + \nabla u \nabla v) dx,$$

and the deduced norm

$$\|u\|_{\mathbf{H}}^2 = \int_{\Omega} |\Delta u|^2 dx + \int_{\Omega} |\nabla u|^2 dx.$$

Let  $\lambda_k$  ( $k = 1, 2, \dots$ ) denote the eigenvalues and  $\varphi_k$  ( $k = 1, 2, \dots$ ) the corresponding eigenfunctions of the eigenvalue problem

$$\begin{cases} -\Delta u = \lambda u, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases}$$

where each eigenvalue  $\lambda_k$  is repeated as the multiplicity; recall that  $0 < \lambda_1 < \lambda_2 \leq \lambda_3 \leq \dots \leq \lambda_k \rightarrow \infty$  and that  $\varphi_1(x) > 0$  for  $x \in \Omega$ . We can easily observe that  $\Lambda_k = \lambda_k(\lambda_k - c)$ ,  $k = 1, 2, \dots$ , are eigenvalues of the eigenvalue problem

$$\begin{cases} \Delta^2 u(x) + c \Delta u = \Lambda u, & \text{in } \Omega, \\ u = \Delta u = 0, & \text{on } \partial\Omega, \end{cases}$$

and the corresponding eigenfunctions are still  $\varphi_k(x)$ .

The set of  $\{\varphi_k(x)\}$  is an orthogonal base on space  $\mathbf{H}$ ; thus one may denote an element  $u$  of  $\mathbf{H}$  as  $\sum_{k=1}^{\infty} u_k \varphi_k$ ,  $\sum_{k=1}^{\infty} u_k^2 < \infty$ .

Assume that  $c < \lambda_1$ ; let us define a norm of  $u \in \mathbf{H}$  as follows:

$$\|u\|^2 = \int_{\Omega} |\Delta u|^2 dx - c \int_{\Omega} |\nabla u|^2 dx.$$

It is easy to show that the norm  $\|\cdot\|$  is a equivalent norm on  $\mathbf{H}$  and the following Poincaré inequality holds:

$$\|u\| \geq \Lambda_1 \|u\|_{L^2}$$

for all  $u \in \mathbf{H}$ .

Consider the following Navier boundary value problem:

$$\begin{cases} \Delta^2 u(x) + c\Delta u = f(x, u), & \text{in } \Omega, \\ u = \Delta u = 0, & \text{on } \partial\Omega, \end{cases} \tag{1.1}$$

where  $\Delta^2$  is the biharmonic operator,  $\Omega$  is a bounded smooth domain in  $\mathbb{R}^N$  ( $N > 4$ ), and  $c < \lambda_1$ .

Let  $f$  be a continuous function on  $\Omega \times \mathbb{R}$ . Suppose that there are measurable real functions  $p(x)$  and  $q(x)$  on  $\Omega$  such that

$$\lim_{|t| \rightarrow 0} \frac{f(x, t)}{t} = p(x), \quad \forall x \in \Omega, \tag{1.2}$$

$$\lim_{|t| \rightarrow \infty} \frac{f(x, t)}{t} = q(x), \quad \forall x \in \Omega. \tag{1.3}$$

If the convergences in (1.2) and (1.3) are uniform in  $\Omega$ , we say that  $f$  is uniformly asymptotically linear at zero and infinity. This case has been studied by many authors under various assumptions on  $p(x)$  and  $q(x)$ .

In [1], An and Liu obtained the existence of one non-trivial solution of (1.1), if the following conditions are fulfilled:

- (AL1)  $f(x, t) \in C(\overline{\Omega} \times \mathbb{R})$ ;  $f(x, t) \equiv 0, \forall x \in \Omega, t \leq 0, f(x, t) \geq 0, \forall x \in \Omega, t > 0$ ;
- (AL2)  $\frac{f(x, t)}{t}$  is nondecreasing with respect to  $t \geq 0$  for a.e.  $x \in \Omega$ ;
- (AL3)  $\lim_{|t| \rightarrow 0} \frac{f(x, t)}{t} = \mu; \lim_{|t| \rightarrow \infty} \frac{f(x, t)}{t} = \nu$  uniformly for a.e.  $x \in \Omega$ , where  $\mu < \lambda_1(\lambda_1 - c) < \nu < +\infty, \nu \neq \Lambda_k = \lambda_k(\lambda_k - c)$  are constants.

In [2], under the above similar conditions, Qian and Li established the existence of three non-trivial solutions of (1.1) by use of mountain pass theorem and regularity of critical groups. Similarly, in [3], we also obtained three non-trivial solutions by using mountain pass theorem and Morse theory for problem (1.1) when the nonlinearity  $f$  is resonant at infinity or the nonlinearity  $f$  is not resonant at infinity. Pu *et al.* [4] proved the existence and multiplicity of solutions for the fourth Navier boundary value problems with concave term, which is similar to problem (1.1) when nonlinearity  $f$  is uniformly asymptotically linear at infinity. In [5], Wei established the existence of multiple solutions for problem (1.1) by means of bifurcation theory. Particularly, Liu and Huang [6] obtained one sign-changing solution for problem (1.1) with uniformly asymptotically linear nonlinearity term.

In the present paper, we study the problem in the case that  $f$  may be non-uniformly asymptotically linear. These new aspects with  $p$ -Laplacian were first presented by Duc and Huy in [7]. But they discussed asymmetric non-uniformly asymptotically linear situation and their methods are not directly to use the non-uniformly asymptotically linear Navier boundary value problems since  $u \in \mathbf{H}$  does not imply that  $u^\pm \in \mathbf{H}$ , where  $u^\pm = \max\{\pm u, 0\}$ . Our main results are as follows.

**Theorem 1.1** *Suppose:*

- (H1)  $f(x, 0) = 0, f(x, t)t \geq 0$  for all  $x \in \Omega, t \in \mathbb{R}$ .
- (H2) There exists  $r$  in the interval  $(\frac{N}{4}, \infty)$  such that  $q(x) \in L^r(\Omega)$  with  $\|q\|_{L^r} > 0$ .
- (H3) There exists a nonnegative measurable function  $W$  on  $\Omega$  such that:

(i)

$$|f(x, s)| \leq W(x)|s|, \quad \forall x \in \Omega, \forall s \in \mathbb{R}.$$

(ii) There exists a constant  $K_W$  such that

$$\int_{\Omega} W|u|^2 dx \leq K_W \|u\|^2, \quad \forall u \in \mathbf{H}.$$

(iii) For any sequence  $\{u_m\}$  converging weakly to  $u$  in  $\mathbf{H}$ , there exists a measurable function  $g$  on  $\Omega$  and a subsequence  $\{u_{m_k}\}$  of  $\{u_m\}$  having the following properties:  $|u_{m_k}| \leq g$  for a.e.  $x \in \Omega$ , for any  $k$  and

$$\int_{\Omega} Wg^2 dx < \infty.$$

(H4)  $\gamma(p) > 1$  and  $\gamma(q) < 1$ , where

$$\gamma(p) = \inf \left\{ \int_{\Omega} (|\Delta u|^2 - c|\nabla u|^2) dx, \int_{\Omega} p(x)u^2 dx = 1 \right\},$$

and the definition of  $\gamma(q)$  is similar.

Then problem (1.1) has at least two non-trivial solutions  $u_1$  and  $u_2$  such that  $u_1 > 0, u_2 < 0, I(u_1) > 0$ , and  $I(u_2) > 0$ , where

$$F(x, t) = \int_0^t f(x, s) ds, \quad \forall (x, s) \in \Omega \times \mathbb{R} \tag{1.4}$$

and

$$I(u) = \frac{1}{2} \int_{\Omega} (|\Delta u|^2 - c|\nabla u|^2) dx - \int_{\Omega} F(x, u) dx. \tag{1.5}$$

**Theorem 1.2** *Suppose:*

- (H1\*)  $f \in C^1(\overline{\Omega} \times \mathbb{R}, \mathbb{R}), f(x, 0) = 0, f(x, t)t \geq 0$  for all  $x \in \Omega, t \in \mathbb{R}$ .

(H3\*) There exists a nonnegative measurable function  $W$  on  $\Omega$  with  $W \in L^\infty(\Omega)$  such that

$$|f_s(x, s)| \leq W(x)$$

for all  $x \in \Omega$  and  $s \in \mathbb{R}$ .

(H4\*)  $\gamma(p) > 1$ .

If  $\Lambda_k < q(x) < \Lambda_{k+1}$  for  $k \geq 2$ , then problem (1.1) has at least three non-trivial solutions.

Here we introduce a non-quadratic condition.

(H5)  $\lim_{|t| \rightarrow \infty} [tf(x, t) - 2F(x, t)] = -\infty$  for every  $x \in \Omega$ .

**Theorem 1.3** Suppose (H1\*), (H3\*), (H4\*), and (H5) hold. If  $q(x) \equiv \Lambda_k$  for  $k \geq 2$ , then problem (1.1) has at least three non-trivial solutions.

### 2 Preliminary results

Let  $u$  be in  $\mathbf{H}$ ,  $F$ , and  $I$  be as in (1.4) and (1.5). Put

$$I_+(u) = \frac{1}{2} \int_{\Omega} (|\Delta u|^2 - c|\nabla u|^2) dx - \int_{\Omega} F_+(x, u) dx,$$

where

$$f_+(x, t) = \begin{cases} f(x, t), & t > 0, \\ 0, & t \leq 0. \end{cases}$$

Combining with the knowledge of nonlinear functional analysis, we have the following lemmas.

**Lemma 2.1** Under conditions (H1) and (H3), the functionals  $I$  and  $I_+$  belong to  $C^1(\mathbf{H}, \mathbb{R})$ . Moreover, for every  $u$  and  $v$  in  $\mathbf{H}$ ,

$$\begin{aligned} \langle DI(u), v \rangle &= \int_{\Omega} (\Delta u \Delta v - c \nabla u \nabla v) dx - \int_{\Omega} f(x, u) v dx, \\ \langle DI_+(u), v \rangle &= \int_{\Omega} (\Delta u \Delta v - c \nabla u \nabla v) dx - \int_{\Omega} f(x, u^+) v dx. \end{aligned}$$

*Proof* We only prove the lemma for  $I$  and the case of  $I^+$  is similar. From the knowledge of nonlinear functional analysis, we easily imply that the map  $u \mapsto \frac{1}{2} \|u\|^2$  is continuously Fréchet differentiable from  $\mathbf{H}$  to  $\mathbf{H}$ .

Let

$$G(u) = \int_{\Omega} F(x, u) dx$$

for all  $u \in \mathbf{H}$ . We prove that  $G \in C^1(\mathbf{H}, \mathbb{R})$  by the following steps.

(i) Given  $u, v \in \mathbf{H}$  and  $s \in \mathbb{R} \setminus \{0\}$  with  $|s| \leq 1$ . Using the mean-value theorem, by (H3)(i), one gets

$$\left| \frac{F(x, u + sv) - F(x, u)}{s} \right| \leq \int_0^1 |f(x, u + tsv)| |v| dt \leq 2W(x)(|u| + |v|)|v|. \tag{2.1}$$

Using Hölder’s inequality and by (H3)(ii), we have

$$\int_{\Omega} W(x)|u||v| \, dx \leq \left( \int_{\Omega} W|u|^2 \, dx \right)^{\frac{1}{2}} \left( \int_{\Omega} W|v|^2 \, dx \right)^{\frac{1}{2}} \leq K_W \|u\| \|v\|. \tag{2.2}$$

Since  $W|v|^2$  is integrable, combining (2.1), (2.2), by the Lebesgue dominated convergence theorem, we see that  $G$  is directional-differentiable on  $\mathbf{H}$  and

$$\langle DG(u), v \rangle = \int_{\Omega} f(x, u)v \, dx.$$

Moreover, by the estimate (2.2), it follows that

$$|\langle DG(u), v \rangle| \leq K_W \|u\| \|v\|, \quad \forall v \in \mathbf{H}.$$

Hence,  $DG(u)$  is a continuous linear functional on  $\mathbf{H}$  and  $G$  is Gâteaux-differentiable on  $\mathbf{H}$ .

(ii) We now prove that  $DG$  is continuous on  $\mathbf{H}$ . Let  $\{u_n\}$  converging to  $u$  in  $\mathbf{H}$ . Suppose by contradiction that  $DG(u_n)$  does not converge to  $D(u)$ . Then there exists  $\epsilon > 0$ , a subsequence of  $\{u_n\}$  (it will be also denoted by  $\{u_n\}$ ) and a sequence  $\{v_n\} \in \mathbf{H}$  with  $\|v_n\| = 1$  such that

$$\epsilon < |\langle DG(u_n) - DG(u), v_n \rangle| = \left| \int_{\Omega} (f(x, u_n) - f(x, u))v_n \, dx \right|, \quad \forall n \in N. \tag{2.3}$$

By condition (H3)(iii), there exist measurable functions  $g_1, g_2, v$ , and a strictly increasing sequence  $\{n_k\}$  of positive integer numbers such that  $Wg_i^2$  is integrable and

$$\begin{aligned} \lim_{k \rightarrow \infty} u_{n_k} &= u(x), & \lim_{k \rightarrow \infty} v_{n_k} &= v(x) \quad \text{a.e. } x \in \Omega, \\ |v_{n_k}(x)| &\leq g_1(x), & |u_{n_k}(x)| &\leq g_2(x) \quad \text{a.e. } x \in \Omega. \end{aligned}$$

It follows that

$$\lim_{k \rightarrow \infty} (f(x, u_{n_k}) - f(x, u))v_{n_k} = 0 \quad \text{a.e. } x \in \Omega$$

and by condition (H3)(i),

$$\begin{aligned} |f(x, u_{n_k}(x)) - f(x, u(x))||v_{n_k}(x)| &\leq (|f(x, u_{n_k}(x))| + |f(x, u(x))|)|v_{n_k}(x)| \\ &\leq W(x)|u_{n_k}| |v_{n_k}| + W(x)|u(x)| |v_{n_k}(x)| \\ &\leq W(x)g_2(x)g_1(x) + W(x)|u(x)|g_1(x). \end{aligned}$$

Let  $T = Wg_2g_1 + W|u|g_1$ . By (H3)(ii),  $W(x)|u|^2$  is integrable on  $\Omega$  and  $T$  is therefore integrable on  $\Omega$ . Indeed, it follows from Hölder’s inequality that

$$\begin{aligned} \int_{\Omega} |T(x)| \, dx &\leq \left( \int_{\Omega} W|g_2|^2 \, dx \right)^{\frac{1}{2}} \left( \int_{\Omega} W|g_1|^2 \, dx \right)^{\frac{1}{2}} \\ &\quad + \left( \int_{\Omega} W|u|^2 \, dx \right)^{\frac{1}{2}} \left( \int_{\Omega} W|g_1|^2 \, dx \right)^{\frac{1}{2}}. \end{aligned}$$

Using the Lebesgue dominated convergence theorem, we obtain

$$\lim_{k \rightarrow \infty} \left| \int_{\Omega} (f(x, u_{n_k}) - f(x, u)v_{n_k}) dx \right| = 0,$$

which contradicts (2.3). □

**Lemma 2.2** *Under conditions (H1)-(H4), the functional  $I_+$  satisfies the (PS) condition.*

*Proof* Let  $\{u_n\} \subset \mathbf{H}$  be a sequence such that  $|I'_+(u_n)| \leq c, \langle I'_+(u_n), \phi \rangle \rightarrow 0$  as  $n \rightarrow \infty$ . Note that

$$\langle I'_+(u_n), \phi \rangle = \int_{\Omega} (\Delta u_n \Delta \phi - c \nabla u_n \nabla \phi) dx - \int_{\Omega} f_+(x, u_n) \phi dx = o(\|\phi\|) \tag{2.4}$$

for all  $\phi \in \mathbf{H}$ .

1. We prove that  $\{\|u_n\|\}$  is bounded. Suppose by contradiction that there is a subsequence of  $\{u_n\}$  (also denoted by  $\{u_n\}$ ) such that  $\|u_n\| \rightarrow \infty$ . Put  $w_n = \frac{u_n}{\|u_n\|}$  for every  $n \in \mathbb{N}$ . We have  $\|w_n\| = 1$  for every  $n$ . Without loss of generality, we assume that  $w_n \rightarrow w$  in  $\mathbf{H}$ , then  $w_n \rightarrow w$  in  $L^2(\Omega)$ . Hence,  $w_n \rightarrow w$  a.e. in  $\Omega$ . Dividing both sides of (2.4) by  $\|u_n\|$ , we get

$$\int_{\Omega} (\Delta w_n \Delta \phi - c \nabla w_n \nabla \phi) dx - \int_{\Omega} \frac{f_+(x, u_n)}{\|u_n\|} \phi dx = o\left(\frac{\|\phi\|}{\|u_n\|}\right), \quad \forall \phi \in \mathbf{H}. \tag{2.5}$$

Note that if  $u_n(x) = 0$  then  $w_n(x) = 0$  and

$$\|u_n\|^{-2} (f(x, u_n^+) u_n) = 0 = \frac{f(x, u_n^+)}{|u_n^+|} w_n^2.$$

Taking  $\phi = w_n$  in (2.5), we have

$$\int_{\Omega} (|\Delta w_n|^2 - c \nabla w_n \nabla w_n) dx - \int_{\Omega} \frac{f_+(x, u_n)}{\|u_n\|} w_n dx = o\left(\frac{\|w_n\|}{\|u_n\|}\right). \tag{2.6}$$

Using (H3)(i), we obtain

$$\left| \frac{f(x, u_n^+)}{|u_n^+|} w_n^2 dx \right| \leq \int_{\Omega} W(x) w_n^2 dx.$$

By (H3)(iii), using the Lebesgue dominated convergence theorem, we have

$$\lim_{n \rightarrow \infty} \int_{\Omega} W(x) w_n^2 dx = 0,$$

and thus

$$\lim_{n \rightarrow \infty} \int_{\Omega} \frac{f(x, u_n^+)}{|u_n^+|} w_n^2 dx = 0. \tag{2.7}$$

Letting  $n \rightarrow \infty$  in (2.6), we get

$$0 = \lim_{n \rightarrow \infty} \|w_n\|^2 = 1,$$

which is impossible. Therefore, we conclude that  $w \neq 0$ .

Set  $D = \{x : w(x) \neq 0\}$ . We have  $u_n(x) \rightarrow \infty$  for all  $x \in D$ . Then condition (1.3) implies that

$$\lim_{n \rightarrow \infty} \frac{f(x, u_n^+)}{u_n^+} = q(x)$$

for all  $x \in D$ . Similar to (2.7), we get

$$\lim_{n \rightarrow \infty} \int_{\Omega} \frac{f(x, u_n^+)}{|u_n^+|} w_n^+ \phi \, dx = \int_{\Omega} q(x) w^+ \phi \, dx. \tag{2.8}$$

Since  $w_n \rightarrow w$ , combining (2.8) and letting  $n \rightarrow \infty$  in (2.5), we have

$$\int_{\Omega} (\Delta w \Delta \phi - c \nabla w \nabla \phi) \, dx = \int_{\Omega} q(x) w^+ \phi \, dx. \tag{2.9}$$

Then we have

$$\begin{cases} \Delta^2 w + c \Delta w = q(x) w^+, & x \in \Omega, \\ w|_{\partial \Omega} = \Delta w|_{\partial \Omega} = 0. \end{cases}$$

Meanwhile, let  $-\Delta w = u$ , by the comparison maximum principle  $w > 0$ . This contradicts our assumption (H4). So  $\{u_n\}$  is bounded in  $\mathbf{H}$ .

2. We prove that  $\{u_n\}$  has a strong convergent subsequence. Since  $\mathbf{H}$  is a Hilbert space, we only need to prove that  $\|u_n\| \rightarrow \|u\|$ . We may assume that  $u_n \rightarrow u$ . Using (H3) and the Lebesgue dominated theorem, we obtain

$$\lim_{n \rightarrow \infty} \int_{\Omega} f(x, u_n) u_n \, dx = \int_{\Omega} f(x, u) u \, dx. \tag{2.10}$$

Combining (2.4) and (2.10), we have

$$\lim_{n \rightarrow \infty} \|u_n\| = \|u\|. \tag{2.11} \quad \square$$

**Remark 2.1** Under conditions (H1\*), (H3\*), and (H4\*), this lemma still holds.

**Lemma 2.3** Under conditions (1.2), (H1), (H3), and (H4), there exist positive numbers  $\rho$  and  $\eta$  such that  $I_+(u) \geq \eta$  for all  $u \in \mathbf{H}$  with  $\|u\| = \rho$ .

*Proof* We adapt a new method from [7] to prove this conclusion. Suppose by contradiction that for every  $n \in \mathbb{N}$ , there exists  $u_n$  in  $\mathbf{H}$  such that  $\|u_n\| = n^{-1}$  and

$$I_+(u_n) = \frac{1}{2} \int_{\Omega} (|\Delta u_n|^2 - c |\nabla u_n|^2) \, dx - \int_{\Omega} F(x, u_n^+) \, dx < \frac{1}{n^3}. \tag{2.11}$$

Let  $w_n = nu_n$ ; then  $\|w_n\| = 1$  and we can suppose that  $\{w_n\}$  weakly converges to  $w$  in  $\mathbf{H}$ . Dividing both sides of (2.11) by  $\frac{1}{n^2}$ , one has

$$\frac{1}{2} \|w_n\|^2 - \int_{\Omega} n^2 F(x, u_n^+) \, dx < \frac{1}{n}. \tag{2.12}$$

Now, we claim that

$$\lim_{n \rightarrow \infty} \|w_n\| = \|w\|.$$

From (H3)(i), we have

$$\begin{aligned} & \int_{\Omega} |F(x, n^{-1}w_n) - F(x, n^{-1}w)| \, dx \\ & \leq \int_{\Omega} \int_0^1 |f(x, n^{-1}w + t(n^{-1}w_n - n^{-1}w))(n^{-1}w_n - n^{-1}w)| \, dt \, dx \\ & \leq \int_{\Omega} 2W(x)n^{-2}(|w||w_n - w| + |w_n - w|^2) \, dx. \end{aligned}$$

Therefore, by the Lebesgue dominated convergence theorem and (ii), (iii) of (H3),

$$\lim_{n \rightarrow \infty} \int_{\Omega} n^2 |F(x, n^{-1}w_n) - F(x, n^{-1}w)| \, dx = 0. \tag{2.13}$$

So, our claim holds. Combining this claim and  $w_n$  weakly converging to  $w$ , we get

$$\|w_n - w\| \rightarrow 0 \tag{2.14}$$

as  $n \rightarrow \infty$ .

On the other hand, arguing as in the proof of (2.13), we have

$$\lim_{n \rightarrow \infty} n^2 F(x, n^{-1}w_n^+) \, dx \leq \frac{1}{2} \int_{\Omega} p(x)w^2 \, dx. \tag{2.15}$$

Using (2.14), (2.15) and letting  $n \rightarrow \infty$  in (2.12), we obtain

$$\frac{1}{2} \left( \|w\|^2 - \int_{\Omega} p(x)w^2 \, dx \right) \leq 0. \tag{2.16}$$

According to assumption (H4), this leads to a contradiction. □

**Remark 2.2** Under conditions (1.2), (H1\*), (H3\*), and (H4\*), this lemma still holds.

**Lemma 2.4** Under conditions (H1)-(H4), the functional  $I_+$  satisfies

$$\lim_{t \rightarrow \infty} \frac{I_+(t\phi_1(q))}{t^2} < 0,$$

where  $\phi_1(q)$  is the first eigenfunction of the eigenvalue problem

$$\begin{cases} \Delta^2 u(x) + c\Delta u = \Lambda q(x)u, & \text{in } \Omega, \\ u = \Delta u = 0, & \text{on } \partial\Omega. \end{cases}$$

*Proof* Let  $u = \phi_1(q)$ . Using principal eigenvalue theorem, we have  $u > 0$  a.e. in  $\Omega$ . By the Lebesgue dominated convergence theorem and (H3), we obtain

$$\begin{aligned} \lim_{t \rightarrow \infty} \int_{\Omega} \frac{F(x, tu)}{t^2} \, dx &= \lim_{t \rightarrow \infty} \int_{\Omega} \frac{F(x, tu) - F(x, 0)}{t^2} \, dx = \lim_{t \rightarrow \infty} \int_{\Omega} \int_0^1 \frac{f(x, stu)tu}{t^2} \, ds \, dx \\ &= \int_{\Omega} \int_0^1 \lim_{t \rightarrow \infty} \frac{f(x, stu)}{stu} su^2 \, ds \, dx = \frac{1}{2} \int_{\Omega} q(x)u^2 \, dx. \end{aligned} \tag{2.17}$$



Combining (2.17) and (H4), we have

$$\lim_{t \rightarrow \infty} \frac{I_+(tu)}{t^2} = \frac{1}{2} \|u\|^2 - \frac{1}{2} \int_{\Omega} q(x)u^2 \, dx \leq -c\|u\|^2 < 0. \quad \square$$

**Remark 2.3** Under conditions (1.3), (H1\*), (H3\*), and (H4\*), this lemma still holds.

**Lemma 2.5** Let  $H = V \oplus W$ , where  $V = E_{\lambda_1} \oplus E_{\lambda_2} \oplus \dots \oplus E_{\lambda_k}$ . If satisfies (H1\*), (H3\*) and (H4\*) and  $\lambda_k \leq q(x) < \lambda_{k+1}$ , then:

(i) the functional  $I$  is coercive on  $W$ , that is,

$$I(u) \rightarrow +\infty \quad \text{as } \|u\| \rightarrow +\infty, u \in W$$

and bounded from below on  $W$ ,

(ii) the functional  $I$  is anti-coercive on  $V$ .

*Proof* (i) Suppose by contradiction that there exist  $M > 0$  and  $\{u_n\}$  in  $H$  such that  $\|u_n\| \rightarrow \infty$  and

$$I(u_n) = \frac{1}{2} \|u_n\|^2 - \int_{\Omega} F(x, u_n) \, dx \leq M. \quad (2.18)$$

Let  $w_n = \frac{u_n}{\|u_n\|}$ ; then  $\|w_n\| = 1$ . Now, we may assume that  $w_n \rightharpoonup w$  in  $W$  and  $w_n \rightarrow w$  a.e.  $x \in \Omega$ . It is obvious that  $w \neq 0$ .

Dividing both sides of (2.18) by  $\|u_n\|^2$ , we have

$$\frac{1}{2} - \int_{\Omega} \frac{F(x, u_n)}{\|u_n\|^2} \leq \frac{M}{\|u_n\|^2}. \quad (2.19)$$

By (1.3) and (H3\*), using the Lebesgue dominated theorem and letting  $n \rightarrow \infty$  in (2.19), we get

$$1 - \int_{\Omega} q(x)w^2 \, dx < 0. \quad (2.20)$$

Since  $w_n \rightharpoonup w$ , from (2.20)

$$\|w\|^2 \leq \int_{\Omega} q(x)w^2 \, dx,$$

which leads to a contradiction.

(ii) Case 1. When  $\lambda_k < q(x) < \lambda_{k+1}$ , similar to the proof of (i), it is easy to verify that the conclusion holds.

Case 2. When  $l = \lambda_k$ , write  $G(x, t) = F(x, t) - \frac{1}{2}\lambda_k t^2$ ,  $g(x, t) = f(x, t) - \lambda_k t$ . Then for every  $x \in \Omega$ , (H5), and (1.3) imply that

$$\lim_{|t| \rightarrow \infty} [g(x, t)t - 2G(x, t)] = -\infty \quad (2.21)$$

and

$$\lim_{|t| \rightarrow \infty} \frac{2G(x, t)}{t^2} = 0. \quad (2.22)$$

It follows from (2.21) that for every  $M > 0$ , there exists a constant  $T > 0$  ( $T$  depends on  $x$ ) such that

$$g(x, t)t - 2G(x, t) \leq -M, \quad \forall t \in \mathbb{R}, |t| \geq T. \tag{2.23}$$

For  $\tau > 0$ , we have

$$\frac{d}{d\tau} \frac{G(x, \tau)}{\tau^2} = \frac{g(x, \tau)\tau - 2G(x, \tau)}{\tau^3}. \tag{2.24}$$

Integrating (2.24) over  $[t, s] \subset [T, +\infty)$ , we deduce that

$$\frac{G(x, s)}{s^2} - \frac{G(x, t)}{t^2} \leq \frac{M}{2} \left( \frac{1}{s^2} - \frac{1}{t^2} \right). \tag{2.25}$$

Letting  $s \rightarrow +\infty$  and using (2.22), we see that  $G(x, t) \geq \frac{M}{2}$ , for  $t \in \mathbb{R}, t \geq T$ . A similar argument shows that  $G(x, t) \geq \frac{M}{2}$ , for  $t \in \mathbb{R}, t \leq -T$ . Hence, for every  $x \in \Omega$ , we have

$$\lim_{|t| \rightarrow \infty} G(x, t) \rightarrow +\infty. \tag{2.26}$$

By (2.26), we get

$$\begin{aligned} I(v) &= \frac{1}{2} \int_{\Omega} |\Delta v|^2 dx - \int_{\Omega} F(x, v) dx \\ &= \frac{1}{2} \int_{\Omega} |\Delta v|^2 dx - \frac{1}{2} \lambda_k \int_{\Omega} v^2 dx - \int_{\Omega} G(x, v) dx \\ &\leq -\delta \|v^-\|^2 - \int_{\Omega} G(x, v) dx \rightarrow -\infty \end{aligned}$$

for  $v \in V$  with  $\|v\| \rightarrow +\infty$ , where  $v^- \in E_{\lambda_1} \oplus E_{\lambda_2} \oplus \dots \oplus E_{\lambda_{k-1}}$ . □

**Lemma 2.6** *Under conditions (1.3), (H1\*), and (H3\*), I satisfies the (PS) condition for  $\lambda_k < q(x) < \lambda_{k+1}$ .*

*Proof* Let  $\{u_n\} \subset \mathbf{H}$  be a sequence such that  $|I'(u_n)| \leq c, \langle I'(u_n), \phi \rangle \rightarrow 0$  as  $n \rightarrow \infty$ . Note that

$$\langle I'(u_n), \phi \rangle = \int_{\Omega} (\Delta u_n \Delta \phi - c \nabla u_n \nabla \phi) dx - \int_{\Omega} f(x, u_n) \phi dx = o(\|\phi\|) \tag{2.27}$$

for all  $\phi \in \mathbf{H}$ .

We prove that  $\{\|u_n\|\}$  is bounded. Suppose by contradiction that there is a subsequence of  $\{u_n\}$  (also denoted by  $\{u_n\}$ ) such that  $\|u_n\| \rightarrow \infty$ . Put  $w_n = \frac{u_n}{\|u_n\|}$  for every  $n \in \mathbb{N}$ . We have  $\|w_n\| = 1$  for every  $n$ . Without loss of generality, we assume that  $w_n \rightharpoonup w$  in  $\mathbf{H}$ , then  $w_n \rightarrow w$  in  $L^2(\Omega)$ . Hence,  $w_n \rightarrow w$  a.e. in  $\Omega$ . Dividing both sides of (2.27) by  $\|u_n\|$ , we get

$$\int_{\Omega} (\Delta w_n \Delta \phi - c \nabla w_n \nabla \phi) dx - \int_{\Omega} \frac{f(x, u_n)}{\|u_n\|} \phi dx = o\left(\frac{\|\phi\|}{\|u_n\|}\right), \quad \forall \phi \in \mathbf{H}. \tag{2.28}$$

Note that if  $u_n(x) = 0$  then  $w_n(x) = 0$  and

$$\|u_n\|^{-2} [f(x, u_n)u_n] = 0 = \frac{f(x, u_n)}{|u_n|} w_n^2.$$

Taking  $\phi = w_n$  in (2.28), we have

$$\int_{\Omega} (|\Delta w_n|^2 - c \nabla w_n \nabla w_n) dx - \int_{\Omega} \frac{f(x, u_n)}{\|u_n\|} w_n dx = o\left(\frac{\|w_n\|}{\|u_n\|}\right). \tag{2.29}$$

Using (H3\*), we obtain

$$\left| \frac{f(x, u_n)}{u_n} w_n^2 dx \right| \leq \int_{\Omega} W(x) w_n^2 dx.$$

From the Lebesgue dominated convergence theorem, we have

$$\lim_{n \rightarrow \infty} \int_{\Omega} W(x) w_n^2 dx = 0,$$

and thus

$$\lim_{n \rightarrow \infty} \int_{\Omega} \frac{f(x, u_n)}{u_n} w_n^2 dx = 0. \tag{2.30}$$

Letting  $n \rightarrow \infty$  in (2.29), we get

$$0 = \lim_{n \rightarrow \infty} \|w_n\|^2 = 1,$$

which is impossible. Therefore, we conclude that  $w \neq 0$ .

Set  $D = \{x : w(x) \neq 0\}$ . We have  $u_n(x) \rightarrow \infty$  for all  $x \in D$ . Then condition (1.3) implies that

$$\lim_{n \rightarrow \infty} \frac{f(x, u_n)}{u_n} = q(x)$$

for all  $x \in D$ . Similar to (2.30), we get

$$\lim_{n \rightarrow \infty} \int_{\Omega} \frac{f(x, u_n)}{u_n} w_n \phi dx = \int_{\Omega} q(x) w \phi dx. \tag{2.31}$$

Since  $w_n \rightharpoonup w$ , combining (2.31) and letting  $n \rightarrow \infty$  in (2.28), we have

$$\int_{\Omega} (\Delta w \Delta \phi - c \nabla w \nabla \phi) dx = \int_{\Omega} q(x) w \phi dx. \tag{2.32}$$

Then we have

$$\begin{cases} \Delta^2 w + c \Delta w = q(x)w, & x \in \Omega, \\ w|_{\partial\Omega} = \Delta w|_{\partial\Omega} = 0. \end{cases}$$

This combined with our assumptions implies that  $w = 0$  and it leads to a contradiction. So  $\{u_n\}$  is bounded in  $\mathbf{H}$ . □

**Lemma 2.7** *Under conditions (1.3), (H1\*), (H3\*), and (H5), the functional  $I$  satisfies the (C) condition which is stated in [8] for  $q(x) = \Lambda_k$ .*

*Proof* Suppose  $u_n \in \mathbf{H}$  satisfies

$$I(u_n) \rightarrow c \in \mathbb{R}, \quad (1 + \|u_n\|) \|I'(u_n)\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \tag{2.33}$$

According to the proof of Lemma 2.2, it suffices to prove that  $u_n$  is bounded in  $\mathbf{H}$ . Similar to the proof of Lemma 2.6, we have

$$\int_{\Omega} (\Delta w \Delta \phi \, dx - c \nabla w \nabla \phi) \, dx - \int_{\Omega} lw \phi \, dx = 0, \quad \forall \phi \in \mathbf{H}. \tag{2.34}$$

Therefore  $w \neq 0$  is an eigenfunction of  $\lambda_k$ , then  $|u_n(x)| \rightarrow \infty$  for a.e.  $x \in \Omega$ . It follows from (H5) that

$$\lim_{n \rightarrow +\infty} [f(x, u_n(x))u_n(x) - 2F(x, u_n(x))] = -\infty$$

holds for every  $x \in \Omega$ , which implies that

$$\int_{\Omega} (f(x, u_n)u_n - 2F(x, u_n)) \, dx \rightarrow -\infty \quad \text{as } n \rightarrow \infty. \tag{2.35}$$

On the other hand, (2.33) implies that

$$2I(u_n) - \langle I'(u_n), u_n \rangle \rightarrow 2c \quad \text{as } n \rightarrow \infty.$$

Thus

$$\int_{\Omega} (f(x, u_n)u_n - 2F(x, u_n)) \, dx \rightarrow 2c \quad \text{as } n \rightarrow \infty,$$

which contradicts (2.35). Hence  $u_n$  is bounded. □

It is well known that critical groups and Morse theory are the main tools in solving elliptic partial differential equations. Let us recall some results which will be used later. We refer the reader to [9] for more information on Morse theory.

Let  $\mathbf{H}$  be a Hilbert space and  $I \in C^1(\mathbf{H}, \mathbb{R})$  be a functional satisfying the (PS) condition or the (C) condition, and  $H_q(X, Y)$  be the  $q$ th singular relative homology group with integer coefficients. Let  $u_0$  be an isolated critical point of  $I$  with  $I(u_0) = c, c \in \mathbb{R}$ , and  $U$  be a neighborhood of  $u_0$ . The group

$$C_q(I, u_0) := H_q(I^c \cap U, I^c \cap U \setminus \{u_0\}), \quad q \in \mathbb{Z},$$

is said to be the  $q$ th critical group of  $I$  at  $u_0$ , where  $I^c = \{u \in \mathbf{H} : I(u) \leq c\}$ .

Let  $K := \{u \in \mathbf{H} : I'(u) = 0\}$  be the set of critical points of  $I$  and  $a < \inf I(K)$ , the critical groups of  $I$  at infinity are formally defined by (see [10])

$$C_q(I, \infty) := H_q(\mathbf{H}, I^a), \quad q \in \mathbb{Z}.$$

The following result comes from [9, 10] and will be used to prove the result in this article.

**Proposition 2.1** [10] *Assume that  $\mathbf{H} = H_\infty^+ \oplus H_\infty^-$ ,  $I$  is bounded from below on  $H_\infty^+$  and  $I(u) \rightarrow -\infty$  as  $\|u\| \rightarrow \infty$  with  $u \in H_\infty^-$ . Then*

$$C_k(I, \infty) \not\cong 0, \quad \text{if } k = \dim H_\infty^- < \infty. \tag{2.36}$$

### 3 Proof of the main result

*Proof of Theorem 1.1* By Lemma 2.2, Lemma 2.3, Lemma 2.4, and the mountain pass theorem, the functional  $I_+$  has a critical point  $u_1$  satisfying  $I_+(u_1) \geq \beta$ . Since  $I_+(0) = 0$ ,  $u_1 \neq 0$  and by the maximum principle, we get  $u_1 > 0$ . Hence  $u_1$  is a positive solution of the problem (1.1). Similarly, we can obtain another negative critical point  $u_2$  of  $I$ .  $\square$

*Proof of Theorem 1.2* By Remarks 2.1 and 2.3 and the mountain pass theorem, the functional  $I_+$  has a critical point  $u_1$  satisfying  $I_+(u_1) \geq \beta$ . Since  $I_+(0) = 0$ ,  $u_1 \neq 0$ , and by the maximum principle, we get  $u_1 > 0$ . Hence  $u_1$  is a positive solution of the problem (1.1) and satisfies

$$C_1(I_+, u_1) \neq 0, \quad u_1 > 0. \tag{3.1}$$

From conditions (H1\*) and (H3\*), we easily verify that  $I$  is  $C^2$  on  $\mathbf{H}$ . Thus, by using the results in [2], we obtain

$$C_q(I, u_1) = C_q(I|_{C_0^1(\Omega)}, u_1) = C_q(I_+|_{C_0^1(\Omega)}, u_1) = C_q(I_+, u_1) = \delta_{q,1}Z. \tag{3.2}$$

Similarly, we can obtain another negative critical point  $u_2$  of  $I$  satisfying

$$C_q(I, u_2) = \delta_{q,1}Z. \tag{3.3}$$

Since  $\gamma(p) > 1$ , the zero function is a local minimizer of  $I$ , then

$$C_q(I, 0) = \delta_{q,0}Z. \tag{3.4}$$

On the other hand, by Lemma 2.5, Lemma 2.6, and Proposition 2.1, we have

$$C_k(I, \infty) \not\cong 0. \tag{3.5}$$

Hence  $I$  has a critical point  $u_3$  satisfying

$$C_k(I, u_3) \not\cong 0. \tag{3.6}$$

Since  $k \geq 2$ , it follows from (3.2)-(3.6) that  $u_1, u_2$ , and  $u_3$  are three different non-trivial solutions of the problem (1.1).  $\square$

*Proof of Theorem 1.3* By Lemma 2.5, Lemma 2.7, and Proposition 2.1, we can prove the conclusion (3.5). The other proof is similar to that of Theorem 1.2.  $\square$

**Competing interests**

The authors declare that they have no competing interests.

**Authors' contributions**

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

**Acknowledgements**

This study was supported by the National NSF (Grant No. 11571268) of China, Natural Science Foundation of Gansu Province China (Grant No. 1506RJZE114) and Planned Projects for Postdoctoral Research Funds of Jiangsu Province (Grant No. 1301038C). The authors are very grateful for reviewers' valuable comments and suggestions in improving this paper.

Received: 19 August 2015 Accepted: 4 November 2015 Published online: 14 November 2015

**References**

1. An, YK, Liu, RY: Existence of nontrivial solutions of an asymptotically linear fourth-order elliptic equation. *Nonlinear Anal. TMA* **68**, 3325-3331 (2008)
2. Qian, AX, Li, SJ: Multiple solutions for a fourth-order asymptotically linear elliptic problem. *Acta Math. Sin.* **22**, 1121-1126 (2006)
3. Pei, RC: Multiple solutions for biharmonic equations with asymptotically linear nonlinearities. *Bound. Value Probl.* **2010**, Article ID 241518 (2010)
4. Pu, Y, Wu, XP, Tang, CL: Fourth-order Navier boundary value problem with combined nonlinearities. *J. Math. Anal. Appl.* **398**, 798-813 (2013)
5. Wei, YH: Multiplicity results for some fourth-order elliptic equations. *J. Math. Anal. Appl.* **385**, 797-807 (2012)
6. Liu, XQ, Huang, YS: On sign-changing solution for a fourth-order asymptotically linear elliptic problem. *Nonlinear Anal. TMA* **72**, 2271-2276 (2010)
7. Duc, DM, Huy, NQ: Non-uniformly asymptotically linear  $p$ -Laplacian problems. *Nonlinear Anal. TMA* **92**, 183-197 (2013)
8. Zhou, HS: Existence of asymptotically linear Dirichlet problem. *Nonlinear Anal. TMA* **44**, 909-918 (2001)
9. Chang, KC: *Infinite Dimensional Morse Theory and Multiple Solutions Problems*. Birkhäuser, Boston (1993)
10. Bartsch, T, Li, SJ: *Critical point theory for asymptotically quadratic functionals and applications to problems with resonance*. *Nonlinear Anal. TMA* **28**, 419-441 (1997)

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)

---