

RESEARCH

Open Access



On G -invariant solutions of a singular biharmonic elliptic system involving multiple critical exponents in \mathbb{R}^N

Zhiying Deng^{1*} , Dong Xu¹ and Yisheng Huang²

*Correspondence:

dengzy@cqupt.edu.cn

¹Key Lab of Intelligent Analysis and Decision on Complex Systems, Chongqing University of Posts and Telecommunications, Chongqing, P.R. China

Full list of author information is available at the end of the article

Abstract

In this work, a biharmonic elliptic system is investigated in \mathbb{R}^N , which involves singular potentials and multiple critical exponents. By the Rellich inequality and the symmetric criticality principle, the existence and multiplicity of G -invariant solutions to the system are established. To our best knowledge, our results are new even in the scalar cases.

MSC: 35B33; 35J48; 35J50

Keywords: G -invariant solution; Rellich inequality; Symmetric criticality principle; Biharmonic elliptic system

1 Introduction

In this article, we study the singular fourth-order elliptic problem:

$$\begin{cases} \Delta^2 u = \mu \frac{u}{|x|^4} + Q(x) \sum_{i=1}^m \frac{\varsigma_i \alpha_i}{2^{**}} |u|^{\alpha_i-2} u |v|^{\beta_i} + \sigma h(x) |u|^{q-2} u, & \text{in } \mathbb{R}^N, \\ \Delta^2 v = \mu \frac{v}{|x|^4} + Q(x) \sum_{i=1}^m \frac{\varsigma_i \beta_i}{2^{**}} |u|^{\alpha_i} |v|^{\beta_i-2} v + \sigma h(x) |v|^{q-2} v, & \text{in } \mathbb{R}^N, \\ \int_{\mathbb{R}^N} (|\Delta u|^2 + |\Delta v|^2) dx < +\infty \quad \text{and} \quad u, v \not\equiv 0, & \text{in } \mathbb{R}^N, \end{cases} \quad (1.1)$$

where Δ^2 denotes the biharmonic operator, $N \geq 5$, $\sigma \geq 0$, $\mu \in [0, \bar{\mu})$ with $\bar{\mu} \triangleq \frac{1}{16} N^2 (N-4)^2$, $q \in (1, 2)$, $\varsigma_i \in (0, +\infty)$, and $\alpha_i, \beta_i > 1$ satisfy $\alpha_i + \beta_i = 2^{**}$ ($i = 1, \dots, m; 1 \leq m \in \mathbb{N}$), $2^{**} \triangleq \frac{2N}{N-4}$ is the critical Sobolev exponent; $Q(x)$ and $h(x)$ are G -invariant functions such that $Q(x) \in \mathcal{C}(\mathbb{R}^N) \cap L^\infty(\mathbb{R}^N)$ and $h(x) \in L^\theta(\mathbb{R}^N)$ with $\theta \triangleq 2^{**}/(2^{**} - q)$ (see Sect. 2 for details).

There have been by now a large number of papers concerning the existence, nonexistence as well as qualitative properties of nontrivial solutions to critical elliptic problems of second order. With no hope of being complete, we would like to mention some of them [1–4]. In most of these papers, the authors deal with the elliptic problems involving singular potentials and critical exponents. For instance, Deng and Jin in [4] handled the following singular equation:

$$-\Delta u = \mu \frac{u}{|x|^2} + Q(x) |x|^{-s} u^{2^{**}(s)-1} \quad \text{and} \quad u > 0 \quad \text{in } \mathbb{R}^N, \quad (1.2)$$

where $N > 2$, $\mu \in [0, \frac{1}{4}(N-2)^2]$, $s \in [0, 2)$, $2^*(s) = \frac{2(N-s)}{N-2}$, and $2^*(0) = 2^* \triangleq \frac{2N}{N-2}$, and Q is G -invariant with respect to a subgroup G of $O(\mathbb{N})$. By applying analytic techniques and critical point theory, several results on the existence and multiplicity of G -invariant solutions to (1.2) were obtained. Subsequently, Waliullah [5] extended the results in [4] to the weighted polyharmonic elliptic equations. In particular, Waliullah considered the following semilinear partial differential equation:

$$(-\Delta)^k u = Q(x)|u|^{2_{(k)}^*-2}u \quad \text{in } \mathbb{R}^N, \quad (1.3)$$

where $k > 1$, $N > 2k$, $2_{(k)}^* = \frac{2N}{N-2k}$, and Q is G -invariant. By employing the minimizing sequence and the concentration–compactness method, the author attained the existence of nontrivial G -invariant solution to (1.3). Borrowing ideas from [4, 5], Deng and Huang [6–8] recently established a few valuable results for the scalar elliptic problems in a bounded G -invariant domain. Moreover, let us also mention that when $\mu = 0$ and the right-hand side nonlinearity term $|x|^{-s}u^{2^*(s)-1}$ in (1.2) is substituted by u^{q-1} with $1 < q \leq 2^*$, there have been a variety of remarkable results on G -invariant solutions in [9–11]. Furthermore, for other results about this aspect, see [12] with singular Lane–Emden–Fowler equations, [13] with singular p -Laplacian equations, [14] with biharmonic operators and [15] with $p(x)$ -biharmonic operators [16], and monograph [17] with generalized Lane–Emden–Fowler equations or Gierer–Meinhardt systems involving singular nonlinearity.

For the systems of singular elliptic equations involving critical exponents, a wide range of works concerning the solutions structures have been presented in recent years. For example, Cai and Kang [18] studied the following elliptic system with multiple critical terms:

$$\begin{cases} \mathcal{L}_\mu u = \frac{\varsigma_1 \alpha_1}{2^*} |u|^{\alpha_1-2} u |v|^{\beta_1} + \frac{\varsigma_2 \alpha_2}{2^*} |u|^{\alpha_2-2} u |v|^{\beta_2} + a_1 |u|^{q_1-2} u + a_2 v, & \text{in } \Omega, \\ \mathcal{L}_\mu v = \frac{\varsigma_1 \beta_1}{2^*} |u|^{\alpha_1} |v|^{\beta_1-2} v + \frac{\varsigma_2 \beta_2}{2^*} |u|^{\alpha_2} |v|^{\beta_2-2} v + a_2 u + a_3 |v|^{q_2-2} v, & \text{in } \Omega, \\ u = v = 0, & \text{on } \partial\Omega, \end{cases} \quad (1.4)$$

where $N \geq 3$, $\Omega \subset \mathbb{R}^N$ is a smooth bounded domain such that $0 \in \Omega$, $\mathcal{L}_\mu = -\Delta - \mu|x|^{-2}$, $\mu < \frac{1}{4}(N-2)^2$, $a_j \in \mathbb{R}$ ($j = 1, 2, 3$), $\varsigma_i \in (0, +\infty)$, $q_i \in [2, 2^*)$, and $\alpha_i, \beta_i > 1$ fulfill $\alpha_i + \beta_i = 2^*$ ($i = 1, 2$). By a variational minimax method combined with a delicate analysis of Palais–Smale sequences, the authors proved the existence of positive solutions to (1.4). Very recently, Nyamoradi and Hsu [19] investigated the following quasilinear elliptic system involving multiple critical exponents:

$$\begin{cases} -\operatorname{div}(|x|^{-ap} |\nabla u|^{p-2} \nabla u) = \sum_{i=1}^m \frac{\varsigma_i \alpha_i |u|^{\alpha_i-2} u |v|^{\beta_i}}{p^*(a,b)|x|^{bp^*(a,b)}} + \sum_{i=1}^m \frac{\lambda_i f_i(x)}{|x|^\beta} |u|^{q-2} u, & \text{in } \Omega, \\ -\operatorname{div}(|x|^{-ap} |\nabla v|^{p-2} \nabla v) = \sum_{i=1}^m \frac{\varsigma_i \beta_i |u|^{\alpha_i} |v|^{\beta_i-2} v}{p^*(a,b)|x|^{bp^*(a,b)}} + \sum_{i=1}^m \frac{\mu_i f_i(x)}{|x|^\beta} |v|^{q-2} v, & \text{in } \Omega, \\ u = v = 0, & \text{on } \partial\Omega, \end{cases} \quad (1.5)$$

where $0 \in \Omega$ is a smooth bounded domain in \mathbb{R}^N , $1 < p < N$, $0 \leq a < \frac{N-p}{p}$, $a \leq b < a+1$, $0 < \varsigma_i, \lambda_i, \mu_i < +\infty$, $\alpha_i, \beta_i > 1$, $\alpha_i + \beta_i = p^*(a,b) = \frac{Np}{N-p(a+1-b)}$ for $i = 1, \dots, m$. By employing the analytic techniques of Nehari manifold, the authors established the existence and multiplicity of positive solutions to (1.5) under certain appropriate hypotheses on the parameters $q, \beta, \lambda_i, \mu_i$ and the weighted functions $f_i(x)$ ($i = 1, \dots, m$). Other results relating to

second-order elliptic systems can be found in [20–23] and the references therein. For the systems of fourth-order elliptic equations, we would like to refer the reader to the papers [24–26] for the elliptic problems related to nonlinearities with critical growth.

Nevertheless, elliptic systems involving the G -invariant solutions have seldom been studied; we only find a handful of results in [27–30]. To the best of our knowledge, there are few results on G -invariant solutions for the singular fourth-order elliptic problem (1.1) even in the scalar cases $\sigma = 0$, $0 < \mu < \bar{\mu}$, $m = 1$, and $u = v$. Therefore, it is necessary for us to investigate (1.1) thoroughly. Let $\bar{Q} > 0$ be a constant. This work is dedicated to seeking the G -invariant solutions for both the cases of $\sigma = 0$, $Q(x) \not\equiv \bar{Q}$ and $\sigma > 0$, $Q(x) \equiv \bar{Q}$ in (1.1). Our arguments are mainly based upon the symmetric criticality principle due to Palais [31] and variational methods.

The rest of this article is schemed as follows. The variational framework and the main results of this paper are presented in Sect. 2. The proofs of G -invariant solutions for the cases $\sigma = 0$ and $Q(x) \not\equiv \bar{Q}$ are detailed in Sect. 3, while the multiplicity results for the cases $\sigma > 0$ and $Q(x) \equiv \bar{Q}$ are proved in Sect. 4.

2 Preliminaries and main results

Let $\mathcal{D}^{2,2}(\mathbb{R}^N)$ denote the completion of $\mathcal{C}_0^\infty(\mathbb{R}^N)$ under the norm $(\int_{\mathbb{R}^N} |\Delta u|^2 dx)^{1/2}$, associated with the inner product given by $\langle u, \varphi \rangle = \int_{\mathbb{R}^N} \Delta u \Delta \varphi dx$. Recall the well-known Rellich inequality [32]

$$\int_{\mathbb{R}^N} |\Delta u|^2 dx \geq \bar{\mu} \int_{\mathbb{R}^N} \frac{u^2}{|x|^4} dx, \quad \forall u \in \mathcal{D}^{2,2}(\mathbb{R}^N), \quad (2.1)$$

where $N \geq 5$, $\bar{\mu} = \frac{1}{16}N^2(N-4)^2$. We now employ the following norm in $\mathcal{D}^{2,2}(\mathbb{R}^N)$:

$$\|u\|_\mu \triangleq \left[\int_{\mathbb{R}^N} (|\Delta u|^2 - \mu |x|^{-4} u^2) dx \right]^{\frac{1}{2}}, \quad 0 \leq \mu < \bar{\mu}.$$

Thanks to the Rellich inequality (2.1), we find that the above norm $\|\cdot\|_\mu$ is equivalent to the usual norm $(\int_{\mathbb{R}^N} |\Delta \cdot|^2 dx)^{1/2}$. Besides, we define the product space $(\mathcal{D}^{2,2}(\mathbb{R}^N))^2$ endowed with the norm

$$\|(u, v)\|_\mu = (\|u\|_\mu^2 + \|v\|_\mu^2)^{\frac{1}{2}}, \quad \forall (u, v) \in (\mathcal{D}^{2,2}(\mathbb{R}^N))^2. \quad (2.2)$$

As usual, we denote by G any closed subgroup of $O(N)$, the group of orthogonal linear transformations. Let $G_x = \{gx; g \in G\}$ be the orbit of $x \in \mathbb{R}^N$; $|G_x|$ denote the number of elements in G_x and $|G_0| = |G_\infty| = 1$. Denote $|G| = \inf_{x \in \mathbb{R}^N \setminus \{0\}} |G_x|$. Note that $|G|$ may be $+\infty$. We call Ω a G -invariant subset of \mathbb{R}^N , if $x \in \Omega$, then $gx \in \Omega$ for all $g \in G$. A function $f: \mathbb{R}^N \mapsto \mathbb{R}$ is called G -invariant if $f(gx) = f(x)$ for every $g \in G$ and $x \in \mathbb{R}^N$. In particular, an $O(N)$ -invariant function is called radial.

The natural functional space to frame the analysis of (1.1) by variational methods is the Hilbert space $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$, which is the subspace of $(\mathcal{D}^{2,2}(\mathbb{R}^N))^2$ consisting of all G -invariant functions. This work is devoted to the study of the following systems:

$$(\mathcal{D}_\sigma^Q) \begin{cases} \Delta^2 u = \mu \frac{u}{|x|^4} + Q(x) \sum_{i=1}^m \frac{\zeta_i \alpha_i}{2^{**}} |u|^{\alpha_i-2} u |v|^{\beta_i} + \sigma h(x) |u|^{q-2} u, & \text{in } \mathbb{R}^N, \\ \Delta^2 v = \mu \frac{v}{|x|^4} + Q(x) \sum_{i=1}^m \frac{\zeta_i \beta_i}{2^{**}} |u|^{\alpha_i} |v|^{\beta_i-2} v + \sigma h(x) |v|^{q-2} v, & \text{in } \mathbb{R}^N, \\ (u, v) \in (\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2 \quad \text{and} \quad u, v \neq 0, & \text{in } \mathbb{R}^N. \end{cases}$$

To clearly describe the results of this paper, several notations should be presented:

$$\mathcal{A}_\mu \triangleq \inf_{u \in \mathcal{D}^{2,2}(\mathbb{R}^N) \setminus \{0\}} \frac{\int_{\mathbb{R}^N} (|\Delta u|^2 - \mu \frac{u^2}{|x|^4}) dx}{(\int_{\mathbb{R}^N} |u|^{2^{**}} dx)^{\frac{2}{2^{**}}}}, \quad (2.3)$$

$$y_\epsilon(x) \triangleq C\epsilon^{-\Lambda_0} U_\mu \left(\frac{|x|}{\epsilon} \right), \quad (2.4)$$

where $\epsilon > 0$, $\Lambda_0 \triangleq \frac{N-4}{2}$, and the constant $C = C(N, \mu) > 0$, depending only on N and μ . From [26, 33], we mention that $y_\epsilon(x)$ satisfies the following equations:

$$\int_{\mathbb{R}^N} \left(|\Delta y_\epsilon|^2 - \mu \frac{y_\epsilon^2}{|x|^4} \right) dx = 1 \quad (2.5)$$

and

$$\int_{\mathbb{R}^N} y_\epsilon^{2^{**}-1} \varphi dx = \mathcal{A}_\mu^{-\frac{2^{**}}{2}} \int_{\mathbb{R}^N} \left(\Delta y_\epsilon \Delta \varphi - \mu \frac{y_\epsilon \varphi}{|x|^4} \right) dx$$

for all $\varphi \in \mathcal{D}^{2,2}(\mathbb{R}^N)$. Hence, we obtain (let $\varphi = y_\epsilon$)

$$\int_{\mathbb{R}^N} y_\epsilon^{2^{**}} dx = \mathcal{A}_\mu^{-\frac{2^{**}}{2}}. \quad (2.6)$$

According to [26, Lemma 2.1] and [33, Theorem 2], we remark that the function $U_\mu(x)$ in (2.4) is positive, radial symmetric, radially decreasing, and solves

$$\begin{cases} \Delta^2 u = \mu \frac{u}{|x|^4} + u^{2^{**}-1}, & \text{in } \mathbb{R}^N \setminus \{0\}, \\ u \in \mathcal{D}^{2,2}(\mathbb{R}^N) \quad \text{and} \quad u > 0, & \text{in } \mathbb{R}^N \setminus \{0\}. \end{cases}$$

By setting $r = |x|$, there holds that

$$U_\mu(r) = O_1(r^{-l_1(\mu)}), \quad \text{as } r \rightarrow 0, \quad (2.7)$$

$$U_\mu(r) = O_1(r^{-l_2(\mu)}), \quad U'_\mu(r) = O_1(r^{-l_2(\mu)-1}), \quad \text{as } r \rightarrow +\infty, \quad (2.8)$$

where $O_1(r^t)$ ($r \rightarrow r_0$) means that there exist constants $C_1, C_2 > 0$ such that $C_1 r^t \leq O_1(r^t) \leq C_2 r^t$ as $r \rightarrow r_0$, $l_1(\mu) \triangleq \Lambda_0 \vartheta(\mu)$, $l_2(\mu) \triangleq \Lambda_0(2 - \vartheta(\mu))$, $\Lambda_0 = \frac{N-4}{2}$, and $\vartheta(\mu) : [0, \bar{\mu}] \mapsto [0, 1]$ is defined as

$$\vartheta(\mu) \triangleq 1 - \frac{\sqrt{N^2 - 4N + 8 - 4\sqrt{(N-2)^2 + \mu}}}{N-4}.$$

This implies $\vartheta(0) = 0$, $\vartheta(\bar{\mu}) = 1$ and

$$0 \leq l_1(\mu) < \Lambda_0 < l_2(\mu) \leq 2\Lambda_0, \quad \forall \mu \in [0, \bar{\mu}]. \quad (2.9)$$

Moreover, there exist positive constants $C_3 = C_3(N, \mu)$ and $C_4 = C_4(N, \mu)$ such that

$$0 < C_3 \leq U_\mu(x) \left(|x|^{\frac{l_1(\mu)}{\Lambda_0}} + |x|^{\frac{l_2(\mu)}{\Lambda_0}} \right)^{\Lambda_0} \leq C_4, \quad \forall x \in \mathbb{R}^N \setminus \{0\}. \quad (2.10)$$

The following hypotheses are needed.

- (q.1) $Q(x)$ is G -invariant.
- (q.2) $Q(x) \in \mathcal{C}(\mathbb{R}^N) \cap L^\infty(\mathbb{R}^N)$, and $Q_+(x) \not\equiv 0$, where $Q_+(x) = \max\{0, Q(x)\}$.
- (h.1) $h(x)$ is G -invariant.
- (h.2) $h(x)$ is a nonnegative function in \mathbb{R}^N such that

$$0 < \|h\|_\theta \triangleq \left(\int_{\mathbb{R}^N} h^\theta(x) dx \right)^{\frac{1}{\theta}} < +\infty \quad \text{with } \theta = \frac{2^{**}}{2^{**} - q}.$$

The main results of this work can be stated in the following.

Theorem 2.1 *Assume that (q.1) and (q.2) hold. If*

$$\int_{\mathbb{R}^N} Q(x) y_\epsilon^{2^{**}} dx \geq \max\{|G|^{\frac{2-2^{**}}{2}} \mathcal{A}_0^{-\frac{2^{**}}{2}} \|Q_+\|_\infty, \mathcal{A}_\mu^{-\frac{2^{**}}{2}} Q_+(0), \mathcal{A}_\mu^{-\frac{2^{**}}{2}} Q_+(\infty)\} > 0 \quad (2.11)$$

for certain $\epsilon > 0$, where $Q_+(\infty) = \limsup_{|x| \rightarrow \infty} Q_+(x)$, then problem (\mathcal{P}_0^Q) possesses at least one nontrivial solution in $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$.

Corollary 2.1 *Assume that (q.1) and (q.2) hold. Then we have the following statements.*

- (1) *Problem (\mathcal{P}_0^Q) admits at least one nontrivial solution if*

$$Q(0) > 0, \quad Q(0) \geq \max\{|G|^{\frac{2-2^{**}}{2}} (\mathcal{A}_0/\mathcal{A}_\mu)^{-\frac{2^{**}}{2}} \|Q_+\|_\infty, Q_+(\infty)\},$$

*and either (i) $Q(x) \geq Q(0) + \xi_0 |x|^{2^{**}(l_2(\mu) - \Lambda_0)}$ for some $\xi_0 > 0$ and $|x|$ small, or (ii) $|Q(x) - Q(0)| \leq \xi_1 |x|^\varsigma$ for some constants $\xi_1 > 0$, $\varsigma > 2^{**}(l_2(\mu) - \Lambda_0) > 0$ and $|x|$ small and*

$$\int_{\mathbb{R}^N} (Q(x) - Q(0)) |x|^{-2^{**}l_2(\mu)} dx > 0. \quad (2.12)$$

- (2) *Problem (\mathcal{P}_0^Q) has at least one nontrivial solution if $\lim_{|x| \rightarrow \infty} Q(x) = Q(\infty)$ exists and is positive,*

$$Q(\infty) \geq \max\{|G|^{\frac{2-2^{**}}{2}} (\mathcal{A}_0/\mathcal{A}_\mu)^{-\frac{2^{**}}{2}} \|Q_+\|_\infty, Q_+(0)\},$$

*and either (i) $Q(x) \geq Q(\infty) + \xi_2 |x|^{-2^{**}(\Lambda_0 - l_1(\mu))}$ for certain $\xi_2 > 0$ and large $|x|$, or (ii) $|Q(x) - Q(\infty)| \leq \xi_3 |x|^{-\kappa}$ for some constants $\xi_3 > 0$, $\kappa > 2^{**}(\Lambda_0 - l_1(\mu)) > 0$ and large $|x|$ and*

$$\int_{\mathbb{R}^N} (Q(x) - Q(\infty)) |x|^{-2^{**}l_1(\mu)} dx > 0. \quad (2.13)$$

- (3) *If $Q(x) \geq Q(\infty) = Q(0) > 0$ on \mathbb{R}^N and*

$$Q(\infty) = Q(0) \geq |G|^{\frac{2-2^{**}}{2}} (\mathcal{A}_0/\mathcal{A}_\mu)^{-\frac{2^{**}}{2}} \|Q_+\|_\infty,$$

then problem (\mathcal{P}_0^Q) possesses at least one nontrivial solution.

Remark 2.1 Conditions (q.1) and (q.2) are essentially introduced in [9]. According to (q.2), we only presume that $Q(x)$ is bounded and continuous on \mathbb{R}^N . Hence, the above results do not require the continuity of $Q(x)$ at infinity.

Theorem 2.2 Assume that $|G| = +\infty$ and $Q_+(0) = Q_+(\infty) = 0$. Then there exist infinitely many G -invariant solutions to problem (\mathcal{P}_0^Q) .

Corollary 2.2 If Q is a radial function such that $Q_+(0) = Q_+(\infty) = 0$, then there exist infinitely many radial solutions to problem (\mathcal{P}_0^Q) .

Theorem 2.3 Let $\bar{Q} > 0$ be a constant. Assume that $Q(x) \equiv \bar{Q}$ and (h.1), (h.2) hold. Then there exists $\sigma^* > 0$ such that, for any $\sigma \in (0, \sigma^*)$, problem $(\mathcal{P}_\sigma^{\bar{Q}})$ possesses at least two non-trivial solutions in $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$.

Remark 2.2 The main results of this paper extend and complement those of [4, 5, 26, 29, 30]. Even in the scalar cases $\sigma = 0$, $0 < \mu < \bar{\mu}$, $m = 1$, and $u = v$, the above results in the whole space are new.

Throughout this paper, we denote various positive constants as C_i ($i = 1, 2, \dots$) or C . The dual space of $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$ ($(\mathcal{D}^{2,2}(\mathbb{R}^N))^2$, resp.) is denoted by $(\mathcal{D}_G^{-2,2}(\mathbb{R}^N))^2$ ($(\mathcal{D}^{-2,2}(\mathbb{R}^N))^2$, resp.). The ball of center x and radius r is denoted by $B_r(x)$. $o_n(1)$ is a generic infinitesimal value as $n \rightarrow \infty$. For any $\epsilon > 0$, $t \in \mathbb{R}$, $O(\epsilon^t)$ denotes the quantity satisfying $|O(\epsilon^t)|/\epsilon^t \leq C$, and $O_1(\epsilon^t)$ ($\epsilon \rightarrow \epsilon_0$) means that there exist constants $C_1, C_2 > 0$ such that $C_1\epsilon^t \leq O_1(\epsilon^t) \leq C_2\epsilon^t$ as $\epsilon \rightarrow \epsilon_0$. In a Banach space X , we denote by ‘ \rightarrow ’ and ‘ \rightharpoonup ’ strong and weak convergence, respectively. A functional $\mathcal{F} \in \mathcal{C}^1(X, \mathbb{R})$ is called to satisfy the $(PS)_c$ condition if each sequence $\{w_n\}$ in X satisfying $\mathcal{F}(w_n) \rightarrow c$ in \mathbb{R} , $\mathcal{F}'(w_n) \rightarrow 0$ in X^* contains a strongly convergent subsequence.

3 Existence and multiplicity results for problem (\mathcal{P}_0^Q)

The energy functional corresponding to problem (\mathcal{P}_0^Q) is defined on $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$ by

$$\mathcal{F}(u, v) = \frac{1}{2} \|(u, v)\|_\mu^2 - \frac{1}{2^{**}} \int_{\mathbb{R}^N} Q(x) \sum_{i=1}^m \varsigma_i |u|^{\alpha_i} |v|^{\beta_i} dx. \quad (3.1)$$

It follows from (q.2) and the Rellich inequality (2.1) that \mathcal{F} is a well-defined \mathcal{C}^1 functional on $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$. Then the critical points of \mathcal{F} correspond to weak solutions of problem (\mathcal{P}_0^Q) . According to the principle of symmetric criticality (see Lemma 3.1), any critical point of \mathcal{F} in $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$ is also a solution of (\mathcal{P}_0^Q) in $(\mathcal{D}^{2,2}(\mathbb{R}^N))^2$. This means that $(u, v) \in (\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$ satisfies (\mathcal{P}_0^Q) if and only if, for any $(\varphi_1, \varphi_2) \in (\mathcal{D}^{2,2}(\mathbb{R}^N))^2$,

$$\begin{aligned} & \langle \mathcal{F}'(u, v), (\varphi_1, \varphi_2) \rangle \\ &= \int_{\mathbb{R}^N} \left(\Delta u \Delta \varphi_1 + \Delta v \Delta \varphi_2 - \mu \frac{u\varphi_1 + v\varphi_2}{|x|^4} \right) dx \\ & \quad - \frac{1}{2^{**}} \int_{\mathbb{R}^N} Q(x) \left(\varphi_1 \sum_{i=1}^m \varsigma_i \alpha_i |u|^{\alpha_i-2} u |v|^{\beta_i} + \varphi_2 \sum_{i=1}^m \varsigma_i \beta_i |u|^{\alpha_i} |v|^{\beta_i-2} v \right) dx = 0. \end{aligned} \quad (3.2)$$

Lemma 3.1 *If $Q(x)$ is a G -invariant function, then $\mathcal{F}'(u, v) = 0$ in $(\mathcal{D}_G^{-2,2}(\mathbb{R}^N))^2$ implies $\mathcal{F}'(u, v) = 0$ in $(\mathcal{D}^{-2,2}(\mathbb{R}^N))^2$.*

Proof The proof is similar to that of [9, Lemma 1] and is omitted here. \square

For $\mu \in [0, \overline{\mu})$, $\varsigma_i \in (0, +\infty)$, $\alpha_i, \beta_i > 1$, and $\alpha_i + \beta_i = 2^{**}$ ($i = 1, \dots, m$), we define

$$\mathcal{A}_{\mu,m} \triangleq \inf_{(u,v) \in (\mathcal{D}^{2,2}(\mathbb{R}^N) \setminus \{0\})^2} \frac{\int_{\mathbb{R}^N} (|\Delta u|^2 + |\Delta v|^2 - \mu \frac{u^2 + v^2}{|x|^4}) dx}{(\int_{\mathbb{R}^N} \sum_{i=1}^m \varsigma_i |u|^{\alpha_i} |v|^{\beta_i} dx)^{\frac{2}{2^{**}}}}, \quad (3.3)$$

$$\mathcal{B}(\tau) \triangleq \frac{1 + \tau^2}{(\sum_{i=1}^m \varsigma_i \tau^{\beta_i})^{\frac{2}{2^{**}}}}, \quad \tau \geq 0, \quad (3.4)$$

$$\mathcal{B}(\tau_{\min}) \triangleq \min_{\tau \geq 0} \mathcal{B}(\tau) > 0, \quad (3.5)$$

where $\tau_{\min} > 0$ is a minimal point of $\mathcal{B}(\tau)$ and hence a root of the equation

$$\sum_{i=1}^m \varsigma_i \tau^{\beta_i-1} (\alpha_i \tau^2 - \beta_i) = 0, \quad \tau \geq 0. \quad (3.6)$$

Lemma 3.2 *Let $y_\epsilon(x)$ be the minimizer of \mathcal{A}_μ defined in (2.4), $\mu \in [0, \overline{\mu})$, $\varsigma_i \in (0, +\infty)$, $\alpha_i, \beta_i > 1$, and $\alpha_i + \beta_i = 2^{**}$ ($i = 1, \dots, m$). Then we have the following statements.*

- (i) $\mathcal{A}_{\mu,m} = \mathcal{B}(\tau_{\min}) \mathcal{A}_\mu$;
- (ii) $\mathcal{A}_{\mu,m}$ has the minimizer $(y_\epsilon(x), \tau_{\min} y_\epsilon(x))$ for all $\epsilon > 0$.

Proof The proof is a repeat of that in [19, Theorem 2.2] (see also [21, Theorem 5]) and hence is omitted here. \square

To find conditions under which the Palais–Smale condition holds, we need the following concentration compactness principle due to Lions [34].

Lemma 3.3 *Let $\{(u_n, v_n)\}$ be a weakly convergent sequence to (u, v) in $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$ such that $|\Delta u_n|^2 \rightharpoonup \eta^{(1)}$, $|\Delta v_n|^2 \rightharpoonup \eta^{(2)}$, $|u_n|^{\alpha_i} |v_n|^{\beta_i} \rightharpoonup v^{(i)}$ ($i = 1, \dots, m$), $|x|^{-4} |u_n|^2 \rightharpoonup \gamma^{(1)}$, and $|x|^{-4} |v_n|^2 \rightharpoonup \gamma^{(2)}$ in the sense of measures. Then there exists some at most countable set \mathcal{J} , $\{\eta_j^{(1)} \geq 0\}_{j \in \mathcal{J} \cup \{0\}}$, $\{\eta_j^{(2)} \geq 0\}_{j \in \mathcal{J} \cup \{0\}}$, $\{v_j^{(i)} \geq 0\}_{j \in \mathcal{J} \cup \{0\}}$, $\gamma_0^{(1)} \geq 0$, $\gamma_0^{(2)} \geq 0$, $\{x_j\}_{j \in \mathcal{J}} \subset \mathbb{R}^N \setminus \{0\}$ such that*

- (a) $\eta^{(1)} \geq |\Delta u|^2 + \sum_{j \in \mathcal{J}} \eta_j^{(1)} \delta_{x_j} + \eta_0^{(1)} \delta_0$, $\eta^{(2)} \geq |\Delta v|^2 + \sum_{j \in \mathcal{J}} \eta_j^{(2)} \delta_{x_j} + \eta_0^{(2)} \delta_0$,
- (b) $v^{(i)} = |u|^{\alpha_i} |v|^{\beta_i} + \sum_{j \in \mathcal{J}} v_j^{(i)} \delta_{x_j} + v_0^{(i)} \delta_0$, $i = 1, \dots, m$,
- (c) $\gamma^{(1)} = |x|^{-4} |u|^2 + \gamma_0^{(1)} \delta_0$, $\gamma^{(2)} = |x|^{-4} |v|^2 + \gamma_0^{(2)} \delta_0$,
- (d) $\mathcal{A}_{0,m} (\sum_{i=1}^m \varsigma_i v_j^{(i)})^{\frac{2}{2^{**}}} \leq \eta_j^{(1)} + \eta_j^{(2)}$,
- (e) $\mathcal{A}_{\mu,m} (\sum_{i=1}^m \varsigma_i v_0^{(i)})^{\frac{2}{2^{**}}} \leq \eta_0^{(1)} + \eta_0^{(2)} - \mu (\gamma_0^{(1)} + \gamma_0^{(2)})$,

where δ_{x_j} , $j \in \mathcal{J} \cup \{0\}$, is a Dirac mass of 1 concentrated at $x_j \in \mathbb{R}^N$.

To establish the existence results for problem (\mathcal{P}_0^Q) , we need the following local $(PS)_c$ condition, which is indispensable for the proof of Theorem 2.1.

Lemma 3.4 Assume that (q.1) and (q.2) hold. Then the $(PS)_c$ condition in $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$ holds for \mathcal{F} if

$$c < c_0^* \triangleq \frac{2}{N} \min \left\{ |G| \mathcal{A}_{0,m}^{\frac{N}{4}} \|Q_+\|_\infty^{1-\frac{N}{4}}, \mathcal{A}_{\mu,m}^{\frac{N}{4}} Q_+(0)^{1-\frac{N}{4}}, \mathcal{A}_{\mu,m}^{\frac{N}{4}} Q_+(\infty)^{1-\frac{N}{4}} \right\}. \quad (3.7)$$

Proof We follow closely the arguments in [9, Proposition 2]. It is trivial to check that the $(PS)_c$ sequence $\{(u_n, v_n)\}$ of \mathcal{F} is bounded in $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$. Then we may assume that $(u_n, v_n) \rightharpoonup (u, v)$ in $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$. In view of Lemma 3.3, there exist measures $\eta^{(1)}$, $\eta^{(2)}$, $\nu^{(i)}$ ($i = 1, \dots, m$), $\gamma^{(1)}$, and $\gamma^{(2)}$ such that relations (a)–(e) of this lemma hold. We begin by considering the concentration at the point $x_j \in \mathbb{R}^N \setminus \{0\}$, $j \in \mathcal{J}$. For $\epsilon > 0$ small, we define the cut-off function $\psi_{x_j}^\epsilon(x) \in \mathcal{C}_0^\infty(\mathbb{R}^N)$ such that $0 \leq \psi_{x_j}^\epsilon(x) \leq 1$, $\psi_{x_j}^\epsilon(x) = 1$ in $B_\epsilon(x_j)$, $\psi_{x_j}^\epsilon(x) = 0$ on $\mathbb{R}^N \setminus B_{2\epsilon}(x_j)$, $|\nabla \psi_{x_j}^\epsilon| \leq 2/\epsilon$, and $|\Delta \psi_{x_j}^\epsilon| \leq 2/\epsilon^2$ on \mathbb{R}^N . Then, by Lemma 3.1, $\lim_{n \rightarrow \infty} \langle \mathcal{F}'(u_n, v_n), (u_n \psi_{x_j}^\epsilon, v_n \psi_{x_j}^\epsilon) \rangle = 0$; hence, combining (3.2), the Hölder inequality, and the Sobolev inequality, we derive

$$\begin{aligned} & \int_{\mathbb{R}^N} \psi_{x_j}^\epsilon \left\{ d\eta^{(1)} + d\eta^{(2)} - \mu(d\gamma^{(1)} + d\gamma^{(2)}) - Q(x) \sum_{i=1}^m \frac{S_i}{2^{**}} (\alpha_i + \beta_i) dv^{(i)} \right\} \\ & \leq \overline{\lim}_{n \rightarrow \infty} \int_{\mathbb{R}^N} \left\{ 2|\Delta u_n \langle \nabla u_n, \nabla \psi_{x_j}^\epsilon \rangle + \Delta v_n \langle \nabla v_n, \nabla \psi_{x_j}^\epsilon \rangle| + |(u_n \Delta u_n + v_n \Delta v_n) \Delta \psi_{x_j}^\epsilon| \right\} dx \\ & \leq \sup_{n \geq 1} \left(\int_{\mathbb{R}^N} |\Delta u_n|^2 dx \right)^{\frac{1}{2}} \left[2 \overline{\lim}_{n \rightarrow \infty} \left(\int_{\mathbb{R}^N} |\nabla u_n|^2 |\nabla \psi_{x_j}^\epsilon|^2 dx \right)^{\frac{1}{2}} \right. \\ & \quad \left. + \overline{\lim}_{n \rightarrow \infty} \left(\int_{\mathbb{R}^N} |u_n|^2 |\Delta \psi_{x_j}^\epsilon|^2 dx \right)^{\frac{1}{2}} \right] \\ & \quad + \sup_{n \geq 1} \left(\int_{\mathbb{R}^N} |\Delta v_n|^2 dx \right)^{\frac{1}{2}} \left[2 \overline{\lim}_{n \rightarrow \infty} \left(\int_{\mathbb{R}^N} |\nabla v_n|^2 |\nabla \psi_{x_j}^\epsilon|^2 dx \right)^{\frac{1}{2}} \right. \\ & \quad \left. + \overline{\lim}_{n \rightarrow \infty} \left(\int_{\mathbb{R}^N} |v_n|^2 |\Delta \psi_{x_j}^\epsilon|^2 dx \right)^{\frac{1}{2}} \right] \\ & \leq C \left\{ \left(\int_{\mathbb{R}^N} |\nabla u|^2 |\nabla \psi_{x_j}^\epsilon|^2 dx \right)^{\frac{1}{2}} + \left(\int_{\mathbb{R}^N} |u|^2 |\Delta \psi_{x_j}^\epsilon|^2 dx \right)^{\frac{1}{2}} + \left(\int_{\mathbb{R}^N} |v|^2 |\Delta \psi_{x_j}^\epsilon|^2 dx \right)^{\frac{1}{2}} \right. \\ & \quad \left. + \left(\int_{\mathbb{R}^N} |\nabla v|^2 |\nabla \psi_{x_j}^\epsilon|^2 dx \right)^{\frac{1}{2}} \right\} \leq C \left\{ \left(\int_{B_{2\epsilon}(x_j)} |\nabla u|^{\frac{2N}{N-2}} dx \right)^{\frac{N-2}{2N}} \left(\int_{\mathbb{R}^N} |\nabla \psi_{x_j}^\epsilon|^N dx \right)^{\frac{1}{N}} \right. \\ & \quad + \left(\int_{B_{2\epsilon}(x_j)} |u|^{2^{**}} dx \right)^{\frac{1}{2^{**}}} \left(\int_{\mathbb{R}^N} |\Delta \psi_{x_j}^\epsilon|^{\frac{N}{2}} dx \right)^{\frac{2}{N}} \\ & \quad + \left(\int_{B_{2\epsilon}(x_j)} |v|^{2^{**}} dx \right)^{\frac{1}{2^{**}}} \left(\int_{\mathbb{R}^N} |\Delta \psi_{x_j}^\epsilon|^{\frac{N}{2}} dx \right)^{\frac{2}{N}} \\ & \quad + \left(\int_{B_{2\epsilon}(x_j)} |\nabla v|^{\frac{2N}{N-2}} dx \right)^{\frac{N-2}{2N}} \left(\int_{\mathbb{R}^N} |\nabla \psi_{x_j}^\epsilon|^N dx \right)^{\frac{1}{N}} \Big\} \leq C \left\{ \left(\int_{B_{2\epsilon}(x_j)} |\nabla u|^{\frac{2N}{N-2}} dx \right)^{\frac{N-2}{2N}} \right. \\ & \quad \left. + \left(\int_{B_{2\epsilon}(x_j)} |\Delta u|^2 dx \right)^{\frac{1}{2}} + \left(\int_{B_{2\epsilon}(x_j)} |\Delta v|^2 dx \right)^{\frac{1}{2}} + \left(\int_{B_{2\epsilon}(x_j)} |\nabla v|^{\frac{2N}{N-2}} dx \right)^{\frac{N-2}{2N}} \right\}. \quad (3.8) \end{aligned}$$

As $\epsilon \rightarrow 0$, it follows from (3.8) and Lemma 3.3 that

$$Q(x_j) \sum_{i=1}^m \varsigma_i v_j^{(i)} \geq \eta_j^{(1)} + \eta_j^{(2)}. \quad (3.9)$$

This means that the concentration of the measures $v^{(i)}$ ($i = 1, \dots, m$) cannot occur at points where $Q(x_j) \leq 0$. By virtue of (3.9) and (d) of Lemma 3.3, we conclude that either (i) $v_j^{(i)} = 0$ ($i = 1, \dots, m$) or (ii) $\sum_{i=1}^m \varsigma_i v_j^{(i)} \geq (\mathcal{A}_{0,m}/\|Q_+\|_\infty)^{N/4}$. Let us now study the possibility of concentration at $x = 0$ and at ∞ . By the argument similar to that of $x_j \in \mathbb{R}^N \setminus \{0\}$, we find $\eta_0^{(1)} + \eta_0^{(2)} - \mu(\gamma_0^{(1)} + \gamma_0^{(2)}) - Q(0) \sum_{i=1}^m \varsigma_i v_0^{(i)} \leq 0$. Together with (e) of Lemma 3.3, it follows that either (iii) $v_0^{(i)} = 0$ ($i = 1, \dots, m$) or (iv) $\sum_{i=1}^m \varsigma_i v_0^{(i)} \geq (\mathcal{A}_{\mu,m}/Q_+(0))^{N/4}$. To discuss the concentration at infinity of the sequence $\{(u_n, v_n)\}$, we define the following quantities:

$$\begin{aligned} (1) \quad & \eta_\infty^{(1)} = \lim_{R \rightarrow \infty} \overline{\lim_{n \rightarrow \infty}} \int_{|x| > R} |\Delta u_n|^2 dx, \quad \eta_\infty^{(2)} = \lim_{R \rightarrow \infty} \overline{\lim_{n \rightarrow \infty}} \int_{|x| > R} |\Delta v_n|^2 dx, \\ (2) \quad & v_\infty^{(i)} = \lim_{R \rightarrow \infty} \overline{\lim_{n \rightarrow \infty}} \int_{|x| > R} |u_n|^{\alpha_i} |v_n|^{\beta_i} dx, \quad i = 1, \dots, m, \\ (3) \quad & \gamma_\infty^{(1)} = \lim_{R \rightarrow \infty} \overline{\lim_{n \rightarrow \infty}} \int_{|x| > R} |x|^{-4} |u_n|^2 dx, \quad \gamma_\infty^{(2)} = \lim_{R \rightarrow \infty} \overline{\lim_{n \rightarrow \infty}} \int_{|x| > R} |x|^{-4} |v_n|^2 dx. \end{aligned}$$

It is obvious that $\eta_\infty^{(1)}, \eta_\infty^{(2)}, v_\infty^{(i)}$ ($i = 1, \dots, m$), $\gamma_\infty^{(1)}$, and $\gamma_\infty^{(2)}$ defined by (1)–(3) exist and are finite. For $R > 1$, let $\psi_R(x) \in \mathcal{C}^\infty(\mathbb{R}^N)$ be a function such that $0 \leq \psi_R(x) \leq 1$, $\psi_R(x) = 1$ for $|x| > R + 1$, $\psi_R(x) = 0$ for $|x| < R$, $|\nabla \psi_R| \leq 2/R$, and $|\Delta \psi_R| \leq 2/R^2$. Because the sequence $\{(u_n \psi_R, v_n \psi_R)\}$ is bounded in $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$, we deduce from (3.2) and the fact that $\alpha_i + \beta_i = 2^{**}$ ($i = 1, \dots, m$) that

$$\begin{aligned} 0 &= \lim_{n \rightarrow \infty} \langle \mathcal{F}'(u_n, v_n), (u_n \psi_R, v_n \psi_R) \rangle \\ &= \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \left\{ \left(|\Delta u_n|^2 + |\Delta v_n|^2 - \mu \frac{|u_n|^2 + |v_n|^2}{|x|^4} - Q(x) \sum_{i=1}^m \varsigma_i |u_n|^{\alpha_i} |v_n|^{\beta_i} \right) \psi_R \right. \\ &\quad \left. + (2\Delta u_n \langle \nabla u_n, \nabla \psi_R \rangle + u_n \Delta u_n \Delta \psi_R + 2\Delta v_n \langle \nabla v_n, \nabla \psi_R \rangle + v_n \Delta v_n \Delta \psi_R) \right\} dx. \quad (3.10) \end{aligned}$$

Furthermore, by utilizing the Hölder inequality and the Sobolev inequality, we obtain

$$\begin{aligned} & \lim_{R \rightarrow \infty} \overline{\lim_{n \rightarrow \infty}} \int_{\mathbb{R}^N} (2|\Delta u_n \langle \nabla u_n, \nabla \psi_R \rangle| + |u_n \Delta u_n \Delta \psi_R|) dx \\ & \leq \lim_{R \rightarrow \infty} \overline{\lim_{n \rightarrow \infty}} \left(\int_{\mathbb{R}^N} |\Delta u_n|^2 dx \right)^{\frac{1}{2}} \left[2 \left(\int_{\mathbb{R}^N} |\nabla u_n|^2 |\nabla \psi_R|^2 dx \right)^{\frac{1}{2}} \right. \\ & \quad \left. + \left(\int_{\mathbb{R}^N} |u_n|^2 |\Delta \psi_R|^2 dx \right)^{\frac{1}{2}} \right] \\ & \leq C \lim_{R \rightarrow \infty} \left\{ \left(\int_{R < |x| < R+1} |\nabla u|^2 |\nabla \psi_R|^2 dx \right)^{\frac{1}{2}} + \left(\int_{R < |x| < R+1} |u|^2 |\Delta \psi_R|^2 dx \right)^{\frac{1}{2}} \right\} \\ & \leq C \lim_{R \rightarrow \infty} \left\{ \left(\int_{R < |x| < R+1} |\nabla u|^{\frac{2N}{N-2}} dx \right)^{\frac{N-2}{2N}} + \left(\int_{R < |x| < R+1} |\Delta u|^2 dx \right)^{\frac{1}{2}} \right\} = 0. \end{aligned}$$

Similarly, we have $\lim_{R \rightarrow \infty} \overline{\lim_{n \rightarrow \infty}} \int_{\mathbb{R}^N} (2|\Delta v_n \langle \nabla v_n, \nabla \psi_R \rangle| + |v_n \Delta v_n \Delta \psi_R|) dx = 0$. Consequently, it follows from (3.10) and definitions (1)–(3) of the quantities $\eta_\infty^{(1)}, \eta_\infty^{(2)}, v_\infty^{(i)}$ ($i =$

$1, \dots, m)$, $\gamma_\infty^{(1)}$, and $\gamma_\infty^{(2)}$ that

$$Q_+(\infty) \sum_{i=1}^m \varsigma_i v_\infty^{(i)} \geq \eta_\infty^{(1)} + \eta_\infty^{(2)} - \mu(\gamma_\infty^{(1)} + \gamma_\infty^{(2)}). \quad (3.11)$$

Moreover, in view of (3.3), we find $\mathcal{A}_{\mu,m}(\sum_{i=1}^m \varsigma_i v_\infty^{(i)})^{\frac{2}{2^{**}}} \leq \eta_\infty^{(1)} + \eta_\infty^{(2)} - \mu(\gamma_\infty^{(1)} + \gamma_\infty^{(2)})$. This, combined with (3.11), implies that either (v) $v_\infty^{(i)} = 0$ ($i = 1, \dots, m$) or (vi) $\sum_{i=1}^m \varsigma_i v_\infty^{(i)} \geq (\mathcal{A}_{\mu,m}/Q_+(\infty))^{N/4}$. In the following, we claim that (ii), (iv), and (vi) cannot occur. For every continuous nonnegative function ψ such that $0 \leq \psi(x) \leq 1$ on \mathbb{R}^N , we find

$$\begin{aligned} c &= \lim_{n \rightarrow \infty} \left(\mathcal{F}(u_n, v_n) - \frac{1}{2^{**}} \langle \mathcal{F}'(u_n, v_n), (u_n, v_n) \rangle \right) \\ &= \frac{2}{N} \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \left(|\Delta u_n|^2 + |\Delta v_n|^2 - \mu \frac{|u_n|^2 + |v_n|^2}{|x|^4} \right) dx \\ &\geq \frac{2}{N} \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \left(|\Delta u_n|^2 + |\Delta v_n|^2 - \mu \frac{|u_n|^2 + |v_n|^2}{|x|^4} \right) \psi(x) dx. \end{aligned}$$

Note that the measures $v^{(i)}$ ($i = 1, \dots, m$) are bounded and G -invariant. This means that if (ii) holds, then the set \mathcal{J} must be finite. Moreover, if $x_j \neq 0$ is a singular point of $v^{(i)}$ ($i = 1, \dots, m$), so is gx_j for each $g \in G$, and the mass of $v^{(i)}$ ($i = 1, \dots, m$) concentrated at gx_j is the same for every $g \in G$. Assuming that (ii) occurs for some $j \in \mathcal{J}$ with $x_j \neq 0$, we choose ψ with compact support so that $\psi(gx_j) = 1$ for every $g \in G$, and we derive

$$\begin{aligned} c &\geq \frac{2}{N} |G| (\eta_j^{(1)} + \eta_j^{(2)}) \geq \frac{2}{N} |G| \mathcal{A}_{0,m} \left(\sum_{i=1}^m \varsigma_i v_j^{(i)} \right)^{\frac{2}{2^{**}}} \\ &\geq \frac{2}{N} |G| \mathcal{A}_{0,m} (\mathcal{A}_{0,m} / \|Q_+\|_\infty)^{\frac{2}{2^{**}-2}} = \frac{2}{N} |G| \mathcal{A}_{0,m}^{\frac{N}{4}} \|Q_+\|_\infty^{1-\frac{N}{4}}, \end{aligned}$$

which is impossible. Similarly, assuming that (iv) holds for $x = 0$, we take ψ with compact support so that $\psi(0) = 1$, and we have

$$\begin{aligned} c &\geq \frac{2}{N} (\eta_0^{(1)} + \eta_0^{(2)} - \mu \gamma_0^{(1)} - \mu \gamma_0^{(2)}) \geq \frac{2}{N} \mathcal{A}_{\mu,m} \left(\sum_{i=1}^m \varsigma_i v_0^{(i)} \right)^{\frac{2}{2^{**}}} \\ &\geq \frac{2}{N} \mathcal{A}_{\mu,m} (\mathcal{A}_{\mu,m} / Q_+(0))^{\frac{2}{2^{**}-2}} = \frac{2}{N} \mathcal{A}_{\mu,m}^{\frac{N}{4}} Q_+(0)^{1-\frac{N}{4}}, \end{aligned}$$

a contradiction to (3.7). Finally, if (vi) occurs, we choose $\psi = \psi_R$ to obtain

$$\begin{aligned} c &\geq \frac{2}{N} (\eta_\infty^{(1)} + \eta_\infty^{(2)} - \mu \gamma_\infty^{(1)} - \mu \gamma_\infty^{(2)}) \geq \frac{2}{N} \mathcal{A}_{\mu,m} \left(\sum_{i=1}^m \varsigma_i v_\infty^{(i)} \right)^{\frac{2}{2^{**}}} \\ &\geq \frac{2}{N} \mathcal{A}_{\mu,m} (\mathcal{A}_{\mu,m} / Q_+(\infty))^{\frac{2}{2^{**}-2}} = \frac{2}{N} \mathcal{A}_{\mu,m}^{\frac{N}{4}} Q_+(\infty)^{1-\frac{N}{4}}, \end{aligned}$$

which contradicts (3.7). Hence, $v_j^{(i)} = 0$ ($i = 1, \dots, m$) for all $j \in \mathcal{J} \cup \{0, \infty\}$, and this yields

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \sum_{i=1}^m \varsigma_i |u_n|^{\alpha_i} |v_n|^{\beta_i} dx = \int_{\mathbb{R}^N} \sum_{i=1}^m \varsigma_i |u|^{\alpha_i} |v|^{\beta_i} dx.$$

Finally, taking into account $\lim_{n \rightarrow \infty} \langle \mathcal{F}'(u_n, v_n) - \mathcal{F}'(u, v), (u_n - u, v_n - v) \rangle = 0$, we naturally deduce $(u_n, v_n) \rightarrow (u, v)$ as $n \rightarrow \infty$ in $(\mathcal{D}^{2,2}(\mathbb{R}^N))^2$. \square

Thanks to Lemma 3.4, we immediately obtain the following result.

Corollary 3.1 *If $|G| = +\infty$ and $Q_+(0) = Q_+(\infty) = 0$, then the functional \mathcal{F} satisfies the $(PS)_c$ condition for every $c \in \mathbb{R}$.*

Proof of Theorem 2.1 Let y_ϵ be the extremal function satisfying (2.4)–(2.10). We now choose $\epsilon > 0$ such that (2.11) is fulfilled. It is clear from (q.2), (3.1), and (3.2) that there exist constants $\alpha_0 > 0$ and $\rho > 0$ such that $\mathcal{F}(u, v) \geq \alpha_0$ for all $\|(u, v)\|_\mu = \rho$. Moreover, if we set $u = y_\epsilon$, $v = \tau_{\min} y_\epsilon$, and

$$\begin{aligned} \Phi(t) = \mathcal{F}(ty_\epsilon, t\tau_{\min}y_\epsilon) &= \frac{t^2}{2} (1 + \tau_{\min}^2) \int_{\mathbb{R}^N} (|\Delta y_\epsilon|^2 - \mu \frac{y_\epsilon^2}{|x|^4}) dx \\ &\quad - \frac{t^{2^{**}}}{2^{**}} \sum_{i=1}^m \varsigma_i \tau_{\min}^{\beta_i} \int_{\mathbb{R}^N} Q(x) y_\epsilon^{2^{**}} dx \end{aligned}$$

with $t \geq 0$, then $\max_{t \geq 0} \Phi(t)$ is attained for some finite $\bar{t} > 0$ with $\Phi'(\bar{t}) = 0$. This yields

$$\max_{t \geq 0} \Phi(t) = \mathcal{F}(\bar{t}y_\epsilon, \bar{t}\tau_{\min}y_\epsilon) = \frac{2}{N} \left\{ \frac{(1 + \tau_{\min}^2) \int_{\mathbb{R}^N} (|\Delta y_\epsilon|^2 - \mu \frac{y_\epsilon^2}{|x|^4}) dx}{(\sum_{i=1}^m \varsigma_i \tau_{\min}^{\beta_i} \int_{\mathbb{R}^N} Q(x) y_\epsilon^{2^{**}} dx)^{\frac{2}{2^{**}}}} \right\}^{\frac{2^{**}}{2^{**}-2}}. \quad (3.12)$$

Besides, because $\mathcal{F}(ty_\epsilon, t\tau_{\min}y_\epsilon) \rightarrow -\infty$ as $t \rightarrow +\infty$, there exists $t_0 > 0$ such that $\|(t_0y_\epsilon, t_0\tau_{\min}y_\epsilon)\|_\mu > \rho$ and $\mathcal{F}(t_0y_\epsilon, t_0\tau_{\min}y_\epsilon) < 0$. Now, we define

$$c_0 = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} \mathcal{F}(\gamma(t)), \quad (3.13)$$

where $\Gamma = \{\gamma \in \mathcal{C}([0,1], (\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2); \gamma(0) = (0,0), \mathcal{F}(\gamma(1)) < 0, \|\gamma(1)\|_\mu > \rho\}$. It follows directly from (2.5), (2.11), (3.4), (3.5), (3.7), (3.12), (3.13), and Lemma 3.2 that

$$\begin{aligned} c_0 &\leq \mathcal{F}(\bar{t}y_\epsilon, \bar{t}\tau_{\min}y_\epsilon) = \frac{2}{N} \left\{ \frac{(1 + \tau_{\min}^2) \int_{\mathbb{R}^N} (|\Delta y_\epsilon|^2 - \mu \frac{y_\epsilon^2}{|x|^4}) dx}{(\sum_{i=1}^m \varsigma_i \tau_{\min}^{\beta_i} \int_{\mathbb{R}^N} Q(x) y_\epsilon^{2^{**}} dx)^{\frac{2}{2^{**}}}} \right\}^{\frac{2^{**}}{2^{**}-2}} \\ &\leq \frac{2}{N} \left\{ \frac{\mathcal{B}(\tau_{\min}) \int_{\mathbb{R}^N} (|\Delta y_\epsilon|^2 - \mu \frac{y_\epsilon^2}{|x|^4}) dx}{(\max\{|G|^{\frac{2-2^{**}}{2}} \mathcal{A}_0^{-\frac{2^{**}}{2}} \|Q_+\|_\infty, \mathcal{A}_\mu^{-\frac{2^{**}}{2}} Q_+(0), \mathcal{A}_\mu^{-\frac{2^{**}}{2}} Q_+(\infty)\})^{\frac{2}{2^{**}}}} \right\}^{\frac{2^{**}}{2^{**}-2}} \\ &= \frac{2}{N} \min\{|G|^{\frac{N}{4}} \mathcal{A}_{0,m}^{\frac{N}{4}} \|Q_+\|_\infty^{1-\frac{N}{4}}, \mathcal{A}_{\mu,m}^{\frac{N}{4}} Q_+(0)^{1-\frac{N}{4}}, \mathcal{A}_{\mu,m}^{\frac{N}{4}} Q_+(\infty)^{1-\frac{N}{4}}\} = c_0^*. \end{aligned}$$

If $c_0 < c_0^*$, then the $(PS)_c$ condition holds by Lemma 3.4. Thus we arrive at the conclusion by the mountain pass theorem in [35]. If $c_0 = c_0^*$, then $\gamma(t) = (tt_0y_\epsilon, tt_0\tau_{\min}y_\epsilon)$, with $0 \leq t \leq 1$, is a path in Γ such that $\max_{t \in [0,1]} \mathcal{F}(\gamma(t)) = c_0$. Hence, either $\Phi'(\bar{t}) = 0$ and we are done, or γ can be deformed to a path $\tilde{\gamma} \in \Gamma$ with $\max_{t \in [0,1]} \mathcal{F}(\tilde{\gamma}(t)) < c_0$ and we have a contradiction. Thus we conclude from Lemma 3.1 that there exists a nontrivial G -invariant solution $(u_0, v_0) \in (\mathcal{D}_G^{2,2}(\mathbb{R}^N) \setminus \{0\})^2$ to problem (\mathcal{P}_0^Q) and the results follow. \square

Proof of Corollary 2.1 In view of (2.6) and Theorem 2.1, it is sufficient to prove that

$$\int_{\mathbb{R}^N} (Q(x) - \tilde{Q}) U_{\mu}^{2^{**}} \left(\frac{|x|}{\epsilon} \right) dx \geq 0 \quad (3.14)$$

for some $\epsilon > 0$, where $\tilde{Q} = \max\{|G|^{\frac{2-2^{**}}{2}} (\mathcal{A}_0/\mathcal{A}_{\mu})^{-\frac{2^{**}}{2}} \|Q_+\|_{\infty}, Q_+(0), Q_+(\infty)\}$.

Part (1), case (i). By virtue of (3.14), we need to show that

$$\epsilon^{-2^{**}l_2(\mu)} \int_{\mathbb{R}^N} (Q(x) - Q(0)) U_{\mu}^{2^{**}} \left(\frac{|x|}{\epsilon} \right) dx \geq 0 \quad (3.15)$$

for certain $\epsilon > 0$. By the hypothesis, we choose $\varrho_0 > 0$ so that $Q(x) \geq Q(0) + \xi_0|x|^{2^{**}(l_2(\mu)-\Lambda_0)}$ for $|x| \leq \varrho_0$. It follows from $2^{**}\Lambda_0 = N$ and (2.8) that

$$\begin{aligned} & \epsilon^{-2^{**}l_2(\mu)} \int_{|x| \leq \varrho_0} (Q(x) - Q(0)) U_{\mu}^{2^{**}} \left(\frac{|x|}{\epsilon} \right) dx \\ & \geq \xi_0 \int_{|x| \leq \varrho_0} \epsilon^{-2^{**}l_2(\mu)} |x|^{2^{**}(l_2(\mu)-\Lambda_0)} U_{\mu}^{2^{**}} \left(\frac{|x|}{\epsilon} \right) dx \\ & = \xi_0 \int_{|x| \leq \varrho_0} \left[\left(\frac{|x|}{\epsilon} \right)^{l_2(\mu)} U_{\mu} \left(\frac{|x|}{\epsilon} \right) \right]^{2^{**}} |x|^{-N} dx \rightarrow +\infty \end{aligned} \quad (3.16)$$

as $\epsilon \rightarrow 0$. On the other hand, for any $\epsilon > 0$, we deduce from (2.8), (2.9), and the fact that $2^{**}l_2(\mu) > N$ that

$$\begin{aligned} & \left| \epsilon^{-2^{**}l_2(\mu)} \int_{|x| > \varrho_0} (Q(x) - Q(0)) U_{\mu}^{2^{**}} \left(\frac{|x|}{\epsilon} \right) dx \right| \\ & \leq \int_{|x| > \varrho_0} \frac{|Q(x) - Q(0)|}{|x|^{2^{**}l_2(\mu)}} \left[\left(\frac{|x|}{\epsilon} \right)^{l_2(\mu)} U_{\mu} \left(\frac{|x|}{\epsilon} \right) \right]^{2^{**}} dx \\ & \leq C \int_{|x| > \varrho_0} \frac{1}{|x|^{2^{**}l_2(\mu)}} dx \leq \bar{C}_1 \end{aligned} \quad (3.17)$$

for some constant $\bar{C}_1 > 0$ independent of ϵ . Combining (3.16) and (3.17), we obtain (3.15) for ϵ sufficiently small.

Part (1), case (ii). By the hypothesis, we choose $\varrho_1 > 0$ so that $|Q(x) - Q(0)| \leq \xi_1|x|^{\varsigma}$ for $|x| \leq \varrho_1$. Taking into account $\varsigma > 2^{**}(l_2(\mu) - \Lambda_0) > 0$, $N - 1 + \varsigma - 2^{**}l_2(\mu) > -1$ and $N - 1 - 2^{**}l_2(\mu) < -1$, we derive

$$\begin{aligned} & \epsilon^{-2^{**}l_2(\mu)} \int_{\mathbb{R}^N} |Q(x) - Q(0)| U_{\mu}^{2^{**}} \left(\frac{|x|}{\epsilon} \right) dx \\ & = \int_{\mathbb{R}^N} \frac{|Q(x) - Q(0)|}{|x|^{2^{**}l_2(\mu)}} \left[\left(\frac{|x|}{\epsilon} \right)^{l_2(\mu)} U_{\mu} \left(\frac{|x|}{\epsilon} \right) \right]^{2^{**}} dx \\ & \leq C \int_{\mathbb{R}^N} \frac{|Q(x) - Q(0)|}{|x|^{2^{**}l_2(\mu)}} dx \\ & \leq C \left(\xi_1 \int_{|x| \leq \varrho_1} |x|^{\varsigma-2^{**}l_2(\mu)} dx + \int_{|x| > \varrho_1} |Q(x) - Q(0)| |x|^{-2^{**}l_2(\mu)} dx \right) \\ & \leq C \left(\int_0^{\varrho_1} r^{N-1+\varsigma-2^{**}l_2(\mu)} dr + \int_{\varrho_1}^{+\infty} r^{N-1-2^{**}l_2(\mu)} dr \right) < +\infty. \end{aligned}$$

Thus, by (2.8), (2.12), and the Lebesgue dominated convergence theorem, we have

$$\begin{aligned} & \lim_{\epsilon \rightarrow 0} \int_{\mathbb{R}^N} \epsilon^{-2^{**}l_2(\mu)} (Q(x) - Q(0)) U_\mu^{2^{**}} \left(\frac{|x|}{\epsilon} \right) dx \\ &= \lim_{\epsilon \rightarrow 0} \int_{\mathbb{R}^N} (Q(x) - Q(0)) |x|^{-2^{**}l_2(\mu)} \left[\left(\frac{|x|}{\epsilon} \right)^{l_2(\mu)} U_\mu \left(\frac{|x|}{\epsilon} \right) \right]^{2^{**}} dx \\ &= C \int_{\mathbb{R}^N} (Q(x) - Q(0)) |x|^{-2^{**}l_2(\mu)} dx > 0. \end{aligned}$$

Hence (3.15) holds for ϵ small enough.

Part (2), case (i). According to (3.14), we need to prove that

$$\epsilon^{-2^{**}l_1(\mu)} \int_{\mathbb{R}^N} (Q(x) - Q(\infty)) U_\mu^{2^{**}} \left(\frac{|x|}{\epsilon} \right) dx \geq 0 \quad (3.18)$$

for certain $\epsilon > 0$. By the assumption, we take $\varrho_2 > 0$ such that $Q(x) \geq Q(\infty) + \xi_2 |x|^{-2^{**}(\Lambda_0 - l_1(\mu))}$ for all $|x| \geq \varrho_2$. It follows from (2.7) that

$$\begin{aligned} & \epsilon^{-2^{**}l_1(\mu)} \int_{|x| \geq \varrho_2} (Q(x) - Q(\infty)) U_\mu^{2^{**}} \left(\frac{|x|}{\epsilon} \right) dx \\ &= \int_{|x| \geq \varrho_2} (Q(x) - Q(\infty)) |x|^{-2^{**}l_1(\mu)} \left[\left(\frac{|x|}{\epsilon} \right)^{l_1(\mu)} U_\mu \left(\frac{|x|}{\epsilon} \right) \right]^{2^{**}} dx \\ &\geq \xi_2 \int_{|x| \geq \varrho_2} |x|^{-N} \left[\left(\frac{|x|}{\epsilon} \right)^{l_1(\mu)} U_\mu \left(\frac{|x|}{\epsilon} \right) \right]^{2^{**}} dx \rightarrow +\infty \end{aligned}$$

as $\epsilon \rightarrow +\infty$. On the other hand, for any $\epsilon > 0$, we conclude from (2.7), (q.2), and the fact that $N - 1 - 2^{**}l_1(\mu) > -1$ that

$$\begin{aligned} & \left| \int_{|x| \leq \varrho_2} \epsilon^{-2^{**}l_1(\mu)} (Q(x) - Q(\infty)) U_\mu^{2^{**}} \left(\frac{|x|}{\epsilon} \right) dx \right| \\ &\leq \int_{|x| \leq \varrho_2} \frac{|Q(x) - Q(\infty)|}{|x|^{2^{**}l_1(\mu)}} \left[\left(\frac{|x|}{\epsilon} \right)^{l_1(\mu)} U_\mu \left(\frac{|x|}{\epsilon} \right) \right]^{2^{**}} dx \\ &\leq C \int_{|x| \leq \varrho_2} \frac{|Q(x) - Q(\infty)|}{|x|^{2^{**}l_1(\mu)}} dx \leq C \int_0^{\varrho_2} r^{N-1-2^{**}l_1(\mu)} dr \leq \bar{C}_2 \end{aligned}$$

for some constant $\bar{C}_2 > 0$ independent of $\epsilon > 0$. By putting these two estimates together, we obtain (3.18) for $\epsilon > 0$ large enough.

Part (2), case (ii). By the assumption, we take $\varrho_3 > 0$ such that $|Q(x) - Q(\infty)| \leq \xi_3 |x|^{-\kappa}$ for all $|x| \geq \varrho_3$. Taking into account $\kappa > 2^{**}(\Lambda_0 - l_1(\mu)) > 0$, $N - 1 - \kappa - 2^{**}l_1(\mu) < -1$ and $N - 1 - 2^{**}l_1(\mu) > -1$, we find

$$\begin{aligned} & \epsilon^{-2^{**}l_1(\mu)} \int_{\mathbb{R}^N} |Q(x) - Q(\infty)| U_\mu^{2^{**}} \left(\frac{|x|}{\epsilon} \right) dx \\ &= \int_{\mathbb{R}^N} \frac{|Q(x) - Q(\infty)|}{|x|^{2^{**}l_1(\mu)}} \left[\left(\frac{|x|}{\epsilon} \right)^{l_1(\mu)} U_\mu \left(\frac{|x|}{\epsilon} \right) \right]^{2^{**}} dx \\ &\leq C \int_{\mathbb{R}^N} \frac{|Q(x) - Q(\infty)|}{|x|^{2^{**}l_1(\mu)}} dx \end{aligned}$$

$$\begin{aligned}
&\leq C \left(\int_{|x| \geq \varrho_3} |x|^{-\kappa-2^{**}l_1(\mu)} dx + \int_{|x| \leq \varrho_3} |Q(x) - Q(\infty)| |x|^{-2^{**}l_1(\mu)} dx \right) \\
&\leq C \left(\int_{\varrho_3}^{+\infty} r^{N-1-\kappa-2^{**}l_1(\mu)} dr + \int_0^{\varrho_3} r^{N-1-2^{**}l_1(\mu)} dr \right) < +\infty.
\end{aligned}$$

Therefore, by (2.7), (2.13), and the Lebesgue dominated convergence theorem, we obtain

$$\begin{aligned}
&\lim_{\epsilon \rightarrow +\infty} \int_{\mathbb{R}^N} \epsilon^{-2^{**}l_1(\mu)} (Q(x) - Q(\infty)) U_\mu^{2^{**}} \left(\frac{|x|}{\epsilon} \right) dx \\
&= \lim_{\epsilon \rightarrow +\infty} \int_{\mathbb{R}^N} \frac{Q(x) - Q(\infty)}{|x|^{2^{**}l_1(\mu)}} \left[\left(\frac{|x|}{\epsilon} \right)^{l_1(\mu)} U_\mu \left(\frac{|x|}{\epsilon} \right) \right]^{2^{**}} dx \\
&= C \int_{\mathbb{R}^N} (Q(x) - Q(\infty)) |x|^{-2^{**}l_1(\mu)} dx > 0.
\end{aligned}$$

Thus (3.18) holds for $\epsilon > 0$ enough large. Similar to the above, we find that part (3) follows. \square

To prove Theorem 2.2, we need the following symmetric mountain pass theorem (see [36] or [37, Theorem 9.12]).

Lemma 3.5 *Let X be an infinite dimensional Banach space, and let $\mathcal{F} \in \mathcal{C}^1(X, \mathbb{R})$ be an even functional satisfying the $(PS)_c$ condition for each c and $\mathcal{F}(0) = 0$. Furthermore, one supposes that:*

- (i) *there exist constants $\tilde{\alpha} > 0$ and $\rho > 0$ such that $\mathcal{F}(w) \geq \tilde{\alpha}$ for all $\|w\| = \rho$;*
- (ii) *there exists an increasing sequence of subspaces $\{X_k\}$ of X , with $\dim X_k = k$, such that for every k one can find a constant $R_k > 0$ such that $\mathcal{F}(w) \leq 0$ for all $w \in X_k$ with $\|w\| \geq R_k$.*

Then \mathcal{F} possesses a sequence of critical values $\{c_k\}$ tending to ∞ as $k \rightarrow \infty$.

Proof of Theorem 2.2 We follow closely the arguments in [9, Theorem 3] (see also [38, Theorem 3]). By virtue of Lemma 3.5 with $X = (\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$ and $w = (u, v) \in X$, we easily see from (q.2), (2.2), (3.1), and (3.3) that

$$\mathcal{F}(u, v) \geq \frac{1}{2} \|(u, v)\|_\mu^2 - \frac{1}{2^{**}} \|Q\|_\infty \mathcal{A}_{\mu, m}^{-\frac{2^{**}}{2}} \|(u, v)\|_\mu^{2^{**}}.$$

Thanks to $2^{**} > 2$, there exist constants $\tilde{\alpha} > 0$ and $\rho > 0$ such that $\mathcal{F}(u, v) \geq \tilde{\alpha}$ for any (u, v) with $\|(u, v)\|_\mu = \rho$. To find an appropriate sequence of finite dimensional subspaces of $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$, we set $\Omega = \{x \in \mathbb{R}^N; Q(x) > 0\}$. The set Ω is G -invariant, and we can define $(\mathcal{D}_G^{2,2}(\Omega))^2$, which is the subspace of G -invariant functions of $(\mathcal{D}^{2,2}(\Omega))^2$. Extending functions in $(\mathcal{D}_G^{2,2}(\Omega))^2$ by 0 outside Ω , we can presume that $(\mathcal{D}_G^{2,2}(\Omega))^2 \subset (\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$. Let $\{X_k\}$ be an increasing sequence of subspaces of $(\mathcal{D}_G^{2,2}(\Omega))^2$ with $\dim X_k = k$ for every k . As in [38, Theorem 3], we define $\varphi_{1,k}, \dots, \varphi_{k,k} \in \mathcal{C}_0^\infty(\mathbb{R}^N)$ such that $0 \leq \varphi_{i,k} \leq 1$, $\text{supp}(\varphi_{i,k}) \cap \text{supp}(\varphi_{j,k}) = \emptyset$, $i \neq j$, and

$$|\text{supp}(\varphi_{i,k}) \cap \Omega| > 0, \quad |\text{supp}(\varphi_{j,k}) \cap \Omega| > 0, \quad \forall i, j \in \{1, \dots, k\}.$$

Taking $e_{i,k} = (a\varphi_{i,k}, b\varphi_{i,k}) \in X_k$, $i = 1, \dots, k$, and $X_k = \text{span}\{e_{1,k}, \dots, e_{k,k}\}$, where a and b are two positive constants, we conclude from the construction of X_k that $\dim X_k = k$ for every k . Therefore, there exists a constant $\epsilon(k) > 0$ such that

$$\begin{aligned} & \int_{\Omega} Q(x) (\varsigma_1 |\tilde{u}|^{\alpha_1} |\tilde{v}|^{\beta_1} + \dots + \varsigma_m |\tilde{u}|^{\alpha_m} |\tilde{v}|^{\beta_m}) dx \\ &= \int_{\Omega} Q(x) \left(\varsigma_1 \left| \sum_{i=1}^k a t_{i,k} \varphi_{i,k} \right|^{\alpha_1} \left| \sum_{i=1}^k b t_{i,k} \varphi_{i,k} \right|^{\beta_1} + \dots \right. \\ & \quad \left. + \varsigma_m \left| \sum_{i=1}^k a t_{i,k} \varphi_{i,k} \right|^{\alpha_m} \left| \sum_{i=1}^k b t_{i,k} \varphi_{i,k} \right|^{\beta_m} \right) dx \geq \epsilon(k) \end{aligned}$$

for all $(\tilde{u}, \tilde{v}) = \sum_{i=1}^k t_{i,k} e_{i,k} \in X_k$, with $\|(\tilde{u}, \tilde{v})\|_{\mu} = 1$. Hence, if $(u, v) \in X_k \setminus \{(0, 0)\}$, then we write $(u, v) = t(\tilde{u}, \tilde{v})$, with $t = \|(u, v)\|_{\mu}$ and $\|(\tilde{u}, \tilde{v})\|_{\mu} = 1$. Therefore, we derive

$$\mathcal{F}(u, v) = \frac{1}{2} t^2 - \frac{1}{2^{**}} \int_{\Omega} Q(x) \sum_{i=1}^m \varsigma_i |\tilde{u}|^{\alpha_i} |\tilde{v}|^{\beta_i} dx \leq \frac{1}{2} t^2 - \frac{\epsilon(k)}{2^{**}} t^{2^{**}} \leq 0$$

for $t > 0$ sufficiently large. By Corollary 3.1 and Lemma 3.5, we conclude that there exists a sequence of critical values $c_k \rightarrow \infty$ as $k \rightarrow \infty$ and the results follow. \square

Proof of Corollary 2.2 Because $Q(x)$ is radial, we know that the corresponding group $G = O(\mathbb{N})$ and $|G| = +\infty$. By Corollary 3.1, \mathcal{F} satisfies the $(PS)_c$ condition for every $c \in \mathbb{R}$. Therefore, we deduce from Theorem 2.2 that the results follow. \square

4 Multiplicity results for problem $(\mathcal{P}_{\sigma}^{\bar{Q}})$

The purpose of this section is to investigate problem $(\mathcal{P}_{\sigma}^{\bar{Q}})$ and prove Theorem 2.3; here we always presume that $\sigma > 0$ and $Q(x) \equiv \bar{Q} > 0$ is a constant. The corresponding energy functional of problem $(\mathcal{P}_{\sigma}^{\bar{Q}})$ is defined on $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$ by

$$\mathcal{E}_{\sigma}(u, v) = \frac{1}{2} \|(u, v)\|_{\mu}^2 - \frac{\bar{Q}}{2^{**}} \int_{\mathbb{R}^N} \sum_{i=1}^m \varsigma_i |u|^{\alpha_i} |v|^{\beta_i} dx - \frac{\sigma}{q} \int_{\mathbb{R}^N} h(x) (|u|^q + |v|^q) dx, \quad (4.1)$$

where $1 < q < 2$. In view of (h.2), (2.3), and the Hölder inequality, we find

$$\begin{aligned} & \int_{\mathbb{R}^N} h(x) (|u|^q + |v|^q) dx \\ & \leq \left(\int_{\mathbb{R}^N} h^{\theta}(x) dx \right)^{\frac{1}{\theta}} \left\{ \left(\int_{\mathbb{R}^N} |u|^{2^{**}} dx \right)^{\frac{q}{2^{**}}} + \left(\int_{\mathbb{R}^N} |v|^{2^{**}} dx \right)^{\frac{q}{2^{**}}} \right\} \\ & \leq \mathcal{A}_{\mu}^{-\frac{q}{2}} \|h\|_{\theta} (\|u\|_{\mu}^q + \|v\|_{\mu}^q) \leq C \|h\|_{\theta} \|(u, v)\|_{\mu}^q. \end{aligned} \quad (4.2)$$

It follows from (4.1) and (4.2) that $\mathcal{E}_{\sigma} \in \mathcal{C}^1((\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2, \mathbb{R})$ and there exists a one-to-one correspondence between the weak solutions of $(\mathcal{P}_{\sigma}^{\bar{Q}})$ and the critical points of \mathcal{E}_{σ} . We now observe that an analogously symmetric criticality principle of Lemma 3.1 clearly holds. Consequently, the weak solutions of problem $(\mathcal{P}_{\sigma}^{\bar{Q}})$ are exactly the critical points of the functional \mathcal{E}_{σ} .

Lemma 4.1 Assume that (h.1) and (h.2) hold. Then there exists a positive constant M depending only on N , q , \mathcal{A}_μ , and $\|h\|_\theta$, such that any bounded sequence $\{(u_n, v_n)\} \subset (\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$ satisfying

$$\begin{aligned}\mathcal{E}_\sigma(u_n, v_n) &\rightarrow c < \frac{2}{N} \overline{Q}^{1-\frac{N}{4}} \mathcal{A}_{\mu,m}^{\frac{N}{4}} - M\sigma^{\frac{2}{2-q}}, \\ \mathcal{E}'_\sigma(u_n, v_n) &\rightarrow 0 \quad (n \rightarrow \infty)\end{aligned}\tag{4.3}$$

contains a convergent subsequence.

Proof By the hypothesis, $\{(u_n, v_n)\}$ is bounded in $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$. Hence we obtain a subsequence, still denoted by $\{(u_n, v_n)\}$, satisfying $(u_n, v_n) \rightharpoonup (u, v)$ in $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$, $(u_n, v_n) \rightarrow (u, v)$ a.e. in \mathbb{R}^N and $(u_n, v_n) \rightarrow (u, v)$ in $(L^r_{\text{loc}}(\mathbb{R}^N))^2$ for all $r \in [1, 2^{**})$. By virtue of (h.2), the Hölder inequality and the Lebesgue dominated theorem, we derive

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} h(x)(|u_n|^q + |v_n|^q) dx = \int_{\mathbb{R}^N} h(x)(|u|^q + |v|^q) dx.\tag{4.4}$$

Applying the standard argument, we easily check from (4.4) that (u, v) is a critical point of \mathcal{E}_σ . Further, in view of (h.2), (4.1), (4.2), and the Hölder inequality, by direct calculation, we obtain

$$\begin{aligned}\mathcal{E}_\sigma(u, v) &= \mathcal{E}_\sigma(u, v) - \frac{1}{2^{**}} \langle \mathcal{E}'_\sigma(u, v), (u, v) \rangle \\ &= \frac{2}{N} \|(u, v)\|_\mu^2 - \frac{\sigma}{2^{**}q} (2^{**} - q) \int_{\mathbb{R}^N} h(x)(|u|^q + |v|^q) dx \\ &\geq \frac{2}{N} (\|u\|_\mu^2 + \|v\|_\mu^2) - \frac{\sigma}{2^{**}q} (2^{**} - q) \mathcal{A}_\mu^{-\frac{q}{2}} \|h\|_\theta (\|u\|_\mu^q + \|v\|_\mu^q) \\ &\geq -(2 - q) \left(\frac{qN}{4} \right)^{\frac{q}{2-q}} \left(\frac{2^{**} - q}{2^{**}q} \mathcal{A}_\mu^{-\frac{q}{2}} \|h\|_\theta \right)^{\frac{2}{2-q}} \sigma^{\frac{2}{2-q}} \triangleq -M\sigma^{\frac{2}{2-q}},\end{aligned}\tag{4.5}$$

where $M = (2 - q) \left(\frac{qN}{4} \right)^{\frac{q}{2-q}} \left(\frac{2^{**} - q}{2^{**}q} \mathcal{A}_\mu^{-\frac{q}{2}} \|h\|_\theta \right)^{\frac{2}{2-q}}$ is a positive constant. We now set $\tilde{u}_n = u_n - u$ and $\tilde{v}_n = v_n - v$. Then, by the Brezis–Lieb lemma [39] and arguing as in [40, Lemma 2.1], we have

$$\|(\tilde{u}_n, \tilde{v}_n)\|_\mu^2 = \|(u_n, v_n)\|_\mu^2 - \|(u, v)\|_\mu^2 + o_n(1),\tag{4.6}$$

$$\int_{\mathbb{R}^N} |\tilde{u}_n|^{\alpha_i} |\tilde{v}_n|^{\beta_i} dx = \int_{\mathbb{R}^N} |u_n|^{\alpha_i} |v_n|^{\beta_i} dx - \int_{\mathbb{R}^N} |u|^{\alpha_i} |v|^{\beta_i} dx + o_n(1), \quad i = 1, \dots, m.\tag{4.7}$$

Taking into account $\mathcal{E}_\sigma(u_n, v_n) = c + o_n(1)$ and $\mathcal{E}'_\sigma(u_n, v_n) = o_n(1)$, we conclude from (4.1), (4.4), (4.6), and (4.7) that

$$\begin{aligned}c + o_n(1) &= \mathcal{E}_\sigma(u_n, v_n) = \mathcal{E}_\sigma(u, v) + \frac{1}{2} \|(\tilde{u}_n, \tilde{v}_n)\|_\mu^2 \\ &\quad - \frac{\overline{Q}}{2^{**}} \int_{\mathbb{R}^N} \sum_{i=1}^m \varsigma_i |\tilde{u}_n|^{\alpha_i} |\tilde{v}_n|^{\beta_i} dx + o_n(1)\end{aligned}\tag{4.8}$$

and

$$\|(\tilde{u}_n, \tilde{v}_n)\|_\mu^2 - \overline{Q} \int_{\mathbb{R}^N} \sum_{i=1}^m \varsigma_i |\tilde{u}_n|^{\alpha_i} |\tilde{v}_n|^{\beta_i} dx = o_n(1). \quad (4.9)$$

As a result, for a subsequence $\{(\tilde{u}_n, \tilde{v}_n)\}$, we find

$$\|(\tilde{u}_n, \tilde{v}_n)\|_\mu^2 \rightarrow \bar{\xi} \geq 0, \quad \overline{Q} \int_{\mathbb{R}^N} \sum_{i=1}^m \varsigma_i |\tilde{u}_n|^{\alpha_i} |\tilde{v}_n|^{\beta_i} dx \rightarrow \bar{\xi}$$

as $n \rightarrow \infty$. It follows from (3.3) that $\mathcal{A}_{\mu,m}(\bar{\xi}/\overline{Q})^{\frac{2}{2^{**}}} \leq \bar{\xi}$. This yields either $\bar{\xi} = 0$ or $\bar{\xi} \geq \overline{Q}^{1-\frac{N}{4}} \mathcal{A}_{\mu,m}^{\frac{N}{4}}$. If $\bar{\xi} \geq \overline{Q}^{1-\frac{N}{4}} \mathcal{A}_{\mu,m}^{\frac{N}{4}}$, then we deduce from (4.5), (4.8), and (4.9) that

$$c = \mathcal{E}_\sigma(u, v) + \left(\frac{1}{2} - \frac{1}{2^{**}}\right) \bar{\xi} \geq \frac{2}{N} \overline{Q}^{1-\frac{N}{4}} \mathcal{A}_{\mu,m}^{\frac{N}{4}} - M\sigma^{\frac{2}{2-q}},$$

which contradicts (4.3). Therefore, we obtain $\|(\tilde{u}_n, \tilde{v}_n)\|_\mu^2 \rightarrow 0$ as $n \rightarrow +\infty$, and hence, $(u_n, v_n) \rightarrow (u, v)$ in $(\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$. The conclusion of this lemma follows. \square

Lemma 4.2 Assume that (h.1) and (h.2) hold. Then there exists $\sigma_1^* > 0$ such that for any $\sigma \in (0, \sigma_1^*)$ the following geometric conditions for $\mathcal{E}_\sigma(u, v)$ hold:

- (i) $\mathcal{E}_\sigma(0, 0) = 0$; there exist constants $\overline{\alpha} > 0$ and $\rho > 0$ such that $\mathcal{E}_\sigma(u, v) \geq \overline{\alpha}$ for all $\|(u, v)\|_\mu = \rho$;
- (ii) there exists $(e_u, e_v) \in (\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2$ such that $\|(e_u, e_v)\|_\mu > \rho$ and $\mathcal{E}_\sigma(e_u, e_v) < 0$.

Proof In view of (h.2), (3.3), (4.1), (4.2), and the Hölder inequality, by direct computation, we derive

$$\begin{aligned} \mathcal{E}_\sigma(u, v) &\geq \frac{1}{2} \|(u, v)\|_\mu^2 - \frac{\overline{Q}}{2^{**}} \mathcal{A}_{\mu,m}^{\frac{2^{**}}{2}} \|(u, v)\|_\mu^{2^{**}} - \frac{\sigma}{q} C \|h\|_\theta \|(u, v)\|_\mu^q \\ &\geq \left(\frac{1}{2} - \varsigma_0\right) \|(u, v)\|_\mu^2 - \frac{\overline{Q}}{2^{**}} \mathcal{A}_{\mu,m}^{\frac{2^{**}}{2}} \|(u, v)\|_\mu^{2^{**}} - C(\varsigma_0) \sigma^{\frac{2}{2-q}} \end{aligned} \quad (4.10)$$

for any $\varsigma_0 \in (0, \frac{1}{2})$, where $C(\varsigma_0) = (\frac{2}{q} - 1) \varsigma_0 [C \|h\|_\theta / (2\varsigma_0)]^{2/(2-q)}$ is a positive constant. It follows from the last inequality in (4.10) that there exist constants $\overline{\alpha} > 0$, $\rho > 0$, and $\sigma_1^* > 0$ such that $\mathcal{E}_\sigma(u, v) \geq \overline{\alpha} > 0$ for all $\|(u, v)\|_\mu = \rho$, $\varsigma_0 \in (0, \frac{1}{2})$ and $\sigma \in (0, \sigma_1^*)$. This yields (i).

On the other hand, taking into account $\int_{\mathbb{R}^N} h(x)(|u|^q + |v|^q) dx \geq 0$, we deduce from (4.1) that there exists $(\check{u}, \check{v}) \in (\mathcal{D}_G^{2,2}(\mathbb{R}^N) \setminus \{0\})^2$ such that $\mathcal{E}_\sigma(t\check{u}, t\check{v}) \rightarrow -\infty$ as $t \rightarrow +\infty$. Therefore, we can choose $(e_u, e_v) = (T\check{u}, T\check{v})$ ($T > 0$ large enough) such that $\|(e_u, e_v)\|_\mu > \rho$ and $\mathcal{E}_\sigma(e_u, e_v) < 0$. Thus (ii) follows. \square

Lemma 4.3 Assume that (h.1) and (h.2) hold. Then there exists $\sigma_2^* > 0$ such that

$$\sup_{t \geq 0} \mathcal{E}_\sigma(t\gamma_\epsilon, t\tau_{\min}\gamma_\epsilon) < \frac{2}{N} \overline{Q}^{1-\frac{N}{4}} \mathcal{A}_{\mu,m}^{\frac{N}{4}} - M\sigma^{\frac{2}{2-q}} \quad (4.11)$$

for any $\sigma \in (0, \sigma_2^*)$ and small $\epsilon > 0$, where $M > 0$ is given in Lemma 4.1 and $\tau_{\min} > 0$ satisfies (3.4)–(3.6).

Proof Similar to the proof in Alves [41, Theorem 3], we define the functions

$$\begin{aligned}\Psi(t) = \mathcal{E}_\sigma(t y_\epsilon, t \tau_{\min} y_\epsilon) &= \frac{t^2}{2} (1 + \tau_{\min}^2) \int_{\mathbb{R}^N} \left(|\Delta y_\epsilon|^2 - \mu \frac{y_\epsilon^2}{|x|^4} \right) dx \\ &\quad - \frac{t^{2^{**}}}{2^{**}} \left(\sum_{i=1}^m \varsigma_i \tau_{\min}^{\beta_i} \right) \overline{Q} \int_{\mathbb{R}^N} y_\epsilon^{2^{**}} dx - \frac{\sigma}{q} t^q (1 + \tau_{\min}^q) \int_{\mathbb{R}^N} h(x) y_\epsilon^q dx\end{aligned}\quad (4.12)$$

and

$$\widetilde{\Psi}(t) = \frac{t^2}{2} (1 + \tau_{\min}^2) \int_{\mathbb{R}^N} \left(|\Delta y_\epsilon|^2 - \mu \frac{y_\epsilon^2}{|x|^4} \right) dx - \frac{t^{2^{**}}}{2^{**}} \left(\sum_{i=1}^m \varsigma_i \tau_{\min}^{\beta_i} \right) \overline{Q} \int_{\mathbb{R}^N} y_\epsilon^{2^{**}} dx \quad (4.13)$$

with $t \geq 0$. Note that $\widetilde{\Psi}(0) = 0$, $\widetilde{\Psi}(t) > 0$ for $t \rightarrow 0^+$, and $\lim_{t \rightarrow +\infty} \widetilde{\Psi}(t) = -\infty$. Hence, $\sup_{t \geq 0} \widetilde{\Psi}(t)$ can be achieved at some finite $t_\epsilon^0 > 0$ at which $\widetilde{\Psi}'(t)$ becomes zero. In view of (2.5), (2.6), (3.4)–(3.6), (4.13), and Lemma 3.2, by simple arithmetic, we derive

$$\begin{aligned}\sup_{t \geq 0} \widetilde{\Psi}(t) &= \widetilde{\Psi}(t_\epsilon^0) = \left(\frac{1}{2} - \frac{1}{2^{**}} \right) \left\{ \frac{(1 + \tau_{\min}^2) \int_{\mathbb{R}^N} (|\Delta y_\epsilon|^2 - \mu \frac{y_\epsilon^2}{|x|^4}) dx}{[(\sum_{i=1}^m \varsigma_i \tau_{\min}^{\beta_i}) \overline{Q} \int_{\mathbb{R}^N} y_\epsilon^{2^{**}} dx]^{\frac{2}{2^{**}}}} \right\}^{\frac{2^{**}}{2^{**}-2}} \\ &= \frac{2}{N} \overline{Q}^{1-\frac{N}{4}} (\mathcal{B}(\tau_{\min}) \mathcal{A}_\mu)^{\frac{N}{4}} = \frac{2}{N} \overline{Q}^{1-\frac{N}{4}} \mathcal{A}_{\mu,m}^{\frac{N}{4}}.\end{aligned}\quad (4.14)$$

Let $\overline{\sigma} > 0$ be such that

$$\frac{2}{N} \overline{Q}^{1-\frac{N}{4}} \mathcal{A}_{\mu,m}^{\frac{N}{4}} - M \sigma^{\frac{2}{2-q}} > 0, \quad \forall \sigma \in (0, \overline{\sigma}).$$

On the one hand, by virtue of (h.1), (h.2), (2.5), and (4.12), we conclude that

$$\Psi(t) = \mathcal{E}_\sigma(t y_\epsilon, t \tau_{\min} y_\epsilon) \leq \frac{t^2}{2} (1 + \tau_{\min}^2), \quad \forall t \geq 0, \sigma > 0,$$

and there exists $T_0 \in (0, 1)$ independent of ϵ such that

$$\sup_{0 \leq t \leq T_0} \Psi(t) \leq \frac{T_0^2}{2} (1 + \tau_{\min}^2) < \frac{2}{N} \overline{Q}^{1-\frac{N}{4}} \mathcal{A}_{\mu,m}^{\frac{N}{4}} - M \sigma^{\frac{2}{2-q}}, \quad \forall \sigma \in (0, \overline{\sigma}). \quad (4.15)$$

On the other hand, it follows from (4.12), (4.13), and (4.14) that

$$\begin{aligned}\sup_{t \geq T_0} \Psi(t) &\leq \sup_{t \geq 0} \widetilde{\Psi}(t) - \frac{\sigma}{q} T_0^q (1 + \tau_{\min}^q) \int_{\mathbb{R}^N} h(x) y_\epsilon^q dx \\ &= \frac{2}{N} \overline{Q}^{1-\frac{N}{4}} \mathcal{A}_{\mu,m}^{\frac{N}{4}} - \frac{\sigma}{q} T_0^q (1 + \tau_{\min}^q) \int_{\mathbb{R}^N} h(x) y_\epsilon^q dx.\end{aligned}\quad (4.16)$$

Now, taking $\sigma > 0$ such that $-\frac{\sigma}{q} T_0^q (1 + \tau_{\min}^q) \int_{\mathbb{R}^N} h(x) y_\epsilon^q dx < -M \sigma^{\frac{2}{2-q}}$, namely

$$0 < \sigma < \left[\frac{T_0^q}{qM} (1 + \tau_{\min}^q) \int_{\mathbb{R}^N} h(x) y_\epsilon^q dx \right]^{\frac{2-q}{q}} \triangleq \widetilde{\sigma},$$

we find from (4.16) that

$$\sup_{t \geq T_0} \Psi(t) < \frac{2}{N} \bar{Q}^{1-\frac{N}{4}} \mathcal{A}_{\mu,m}^{\frac{N}{4}} - M\sigma^{\frac{2}{2-q}}, \quad \forall \sigma \in (0, \tilde{\sigma}). \quad (4.17)$$

Choosing $\sigma_2^* = \min\{\bar{\sigma}, \tilde{\sigma}\}$, we deduce from (4.15) and (4.17) that

$$\sup_{t \geq 0} \Psi(t) < \frac{2}{N} \bar{Q}^{1-\frac{N}{4}} \mathcal{A}_{\mu,m}^{\frac{N}{4}} - M\sigma^{\frac{2}{2-q}}, \quad \forall \sigma \in (0, \sigma_2^*),$$

which implies (4.11). Hence the results of this lemma follow. \square

Proof of Theorem 2.3 Taking $\rho > 0$ and $\sigma^* = \min\{\sigma_1^*, \sigma_2^*\}$, for $0 < \sigma < \sigma^*$, given in the proofs of Lemmas 4.2 and 4.3, we define

$$c_1 \triangleq \inf_{\bar{\mathcal{B}}_\rho(0)} \mathcal{E}_\sigma(u, v),$$

where $\bar{\mathcal{B}}_\rho(0) = \{(u, v) \in (\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2; \|(u, v)\|_\mu \leq \rho\}$. It is easy to see that the metric space $\bar{\mathcal{B}}_\rho(0)$ is complete. According to the Ekeland variational principle [42], we deduce that there exists a sequence $\{(u_n, v_n)\} \subset \bar{\mathcal{B}}_\rho(0)$ such that $\mathcal{E}_\sigma(u_n, v_n) \rightarrow c_1$ and $\mathcal{E}'_\sigma(u_n, v_n) \rightarrow 0$ as $n \rightarrow \infty$.

Let $\varphi_0, \psi_0 \in \mathcal{C}_0^\infty(\mathbb{R}^N)$ be the G -invariant functions such that $\varphi_0, \psi_0 > 0$. It follows from (h.1) and (h.2) that $\int_{\mathbb{R}^N} h(x)(\varphi_0^q + \psi_0^q) dx > 0$. In view of $1 < q < 2 < 2^{**}$, we find that there exists $\tilde{t}_0 = \tilde{t}_0(\varphi_0, \psi_0) > 0$ sufficiently small such that

$$\begin{aligned} \mathcal{E}_\sigma(\tilde{t}_0 \varphi_0, \tilde{t}_0 \psi_0) &= \frac{\tilde{t}_0^2}{2} \|(\varphi_0, \psi_0)\|_\mu^2 \\ &\quad - \frac{\bar{Q}}{2^{**}} \tilde{t}_0^{2^{**}} \int_{\mathbb{R}^N} \sum_{i=1}^m s_i \varphi_0^{\alpha_i} \psi_0^{\beta_i} dx - \frac{\sigma}{q} \tilde{t}_0^q \int_{\mathbb{R}^N} h(x)(\varphi_0^q + \psi_0^q) dx < 0. \end{aligned}$$

This yields

$$c_1 < 0 < \frac{2}{N} \bar{Q}^{1-\frac{N}{4}} \mathcal{A}_{\mu,m}^{\frac{N}{4}} - M\sigma^{\frac{2}{2-q}}, \quad \forall \sigma \in (0, \sigma^*).$$

By virtue of Lemma 4.1, \mathcal{E}_σ admits a nontrivial critical point (u_1, v_1) with $\mathcal{E}_\sigma(u_1, v_1) = c_1 < 0$. Applying the principle of symmetric criticality, we obtain that (u_1, v_1) is a nontrivial G -invariant solution of problem (\mathcal{P}_σ^Q) .

Furthermore, we now define

$$c_2 \triangleq \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} \mathcal{E}_\sigma(\gamma(t)),$$

where $\Gamma = \{\gamma \in \mathcal{C}([0,1], (\mathcal{D}_G^{2,2}(\mathbb{R}^N))^2); \gamma(0) = (0,0), \gamma(1) = (e_u, e_v)\}$. It follows from Lemmas 4.2 and 4.3 that

$$0 < \bar{\alpha} \leq c_2 < \frac{2}{N} \bar{Q}^{1-\frac{N}{4}} \mathcal{A}_{\mu,m}^{\frac{N}{4}} - M\sigma^{\frac{2}{2-q}}, \quad \forall \sigma \in (0, \sigma^*).$$

This, combined with the mountain pass theorem, implies that c_2 is another nonzero critical value of \mathcal{E}_σ . Similar to the above arguments, problem (\mathcal{P}_σ^Q) possesses another non-trivial G -invariant solution (u_2, v_2) with $\mathcal{E}_\sigma(u_2, v_2) = c_2 > 0$. \square

Acknowledgements

ZD and DX are supported by the National Natural Science Foundation of China (Nos. 11471235; 11601052) and Chongqing Research Program of Basic Research and Frontier Technology (No. cstc2017jcyjBX0037). YH is supported by the National Natural Science Foundation of China (No. 11471235).

Availability of data and materials

Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

The authors declare that this study was independently finished. All authors read and approved the final manuscript.

Author details

¹Key Lab of Intelligent Analysis and Decision on Complex Systems, Chongqing University of Posts and Telecommunications, Chongqing, P.R. China. ²Department of Mathematics, Soochow University, Suzhou, P.R. China.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 19 December 2017 Accepted: 2 April 2018 Published online: 11 April 2018

References

- Brezis, H., Nirenberg, L.: Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents. *Commun. Pure Appl. Math.* **36**, 437–477 (1983)
- Cerami, G., Zhong, X.-X., Zou, W.-M.: On some nonlinear elliptic PDEs with Sobolev–Hardy critical exponents and a Li–Lin open problem. *Calc. Var. Partial Differ. Equ.* **54**, 1793–1829 (2015)
- Bhakta, M., Santra, S.: On singular equations with critical and supercritical exponents. *J. Differ. Equ.* **263**, 2886–2953 (2017)
- Deng, Y.-B., Jin, L.-Y.: On symmetric solutions of a singular elliptic equation with critical Sobolev–Hardy exponent. *J. Math. Anal. Appl.* **329**, 603–616 (2007)
- Waliullah, S.: Minimizers and symmetric minimizers for problems with critical Sobolev exponent. *Topol. Methods Nonlinear Anal.* **34**, 291–326 (2009)
- Deng, Z.-Y., Huang, Y.-S.: Existence and multiplicity of symmetric solutions for semilinear elliptic equations with singular potentials and critical Hardy–Sobolev exponents. *J. Math. Anal. Appl.* **393**, 273–284 (2012)
- Deng, Z.-Y., Huang, Y.-S.: Existence and multiplicity of symmetric solutions for the weighted critical quasilinear problems. *Appl. Math. Comput.* **219**, 4836–4846 (2013)
- Deng, Z.-Y., Huang, Y.-S.: On positive G -symmetric solutions of a weighted quasilinear elliptic equation with critical Hardy–Sobolev exponent. *Acta Math. Sci.* **34B**, 1619–1633 (2014)
- Bianchi, G., Chabrowski, J., Szulkin, A.: On symmetric solutions of an elliptic equations with a nonlinearity involving critical Sobolev exponent. *Nonlinear Anal.* **25**, 41–59 (1995)
- Bartsch, T., Willem, M.: Infinitely Many Non-radial Solutions of an Euclidean Scalar Field Equation. *Mathematisches Institut, Universität, Heidelberg* (1992)
- Chabrowski, J.: On the existence of G -symmetric entire solutions for semilinear elliptic equations. *Rend. Circ. Mat. Palermo* **41**, 413–440 (1992)
- Ghergu, M., Rădulescu, V.: Multi-parameter bifurcation and asymptotics for the singular Lane–Emden–Fowler equation with a convection term. *Proc. R. Soc. Edinb. A* **135A**, 61–83 (2005)
- Maultsby, B.: Uniqueness of solutions to singular p -Laplacian equations with subcritical nonlinearity. *Adv. Nonlinear Anal.* **6**, 37–59 (2017)
- Demarque, R., Miyagaki, O.: Radial solutions of inhomogeneous fourth order elliptic equations and weighted Sobolev embeddings. *Adv. Nonlinear Anal.* **4**, 135–151 (2015)
- Kong, L.: Multiple solutions for fourth order elliptic problems with $p(x)$ -biharmonic operators. *Opusc. Math.* **36**, 253–264 (2016)
- Ghergu, M., Rădulescu, V.: Singular elliptic problems with lack of compactness. *Ann. Mat. Pura Appl.* **185**, 63–79 (2006)
- Ghergu, M., Rădulescu, V.: *Singular Elliptic Problems: Bifurcation and Asymptotic Analysis*. Oxford Lecture Series in Mathematics and Its Applications, vol. 37. Oxford University Press, Oxford (2008)
- Cai, M.-J., Kang, D.-S.: Elliptic systems involving multiple strongly coupled critical terms. *Appl. Math. Lett.* **25**, 417–422 (2012)
- Nyamoradi, N., Hsu, T.-S.: Existence of multiple positive solutions for semilinear elliptic systems involving m critical Hardy–Sobolev exponents and m sign-changing weight function. *Acta Math. Sci.* **34B**, 483–500 (2014)
- Cortázar, C., Elgueta, M., García-Melián, J.: Analysis of an elliptic system with infinitely many solutions. *Adv. Nonlinear Anal.* **6**, 1–12 (2017)
- Alves, C.O., de Moraes Filho, D.C., Souto, M.A.S.: On systems of elliptic equations involving subcritical or critical Sobolev exponents. *Nonlinear Anal.* **42**, 771–787 (2000)

22. Kang, D.-S.: Positive minimizers of the best constants and solutions to coupled critical quasilinear systems. *J. Differ. Equ.* **260**, 133–148 (2016)
23. Benrhouma, M.: On a singular elliptic system with quadratic growth in the gradient. *J. Math. Anal. Appl.* **448**, 1120–1146 (2017)
24. Lü, D.-F.: Multiple solutions for a class of biharmonic elliptic systems with Sobolev critical exponent. *Nonlinear Anal.* **74**, 6371–6382 (2011)
25. Alvarez-Caudevilla, P., Colorado, E., Galaktionov, V.A.: Existence of solutions for a system of coupled nonlinear stationary bi-harmonic Schrödinger equations. *Nonlinear Anal., Real World Appl.* **23**, 78–93 (2015)
26. Kang, D.-S., Xiong, P.: Ground state solutions to biharmonic equations involving critical nonlinearities and multiple singular potentials. *Appl. Math. Lett.* **66**, 9–15 (2017)
27. Deng, Z.-Y., Huang, Y.-S.: Existence of symmetric solutions for singular semilinear elliptic systems with critical Hardy–Sobolev exponents. *Nonlinear Anal., Real World Appl.* **14**, 613–625 (2013)
28. Kang, D.-S., Yang, F.: Elliptic systems involving multiple critical nonlinearities and symmetric multi-polar potentials. *Sci. China Math.* **57**, 1011–1024 (2014)
29. Deng, Z.-Y., Huang, Y.-S.: Symmetric solutions for a class of singular biharmonic elliptic systems involving critical exponents. *Appl. Math. Comput.* **264**, 323–334 (2015)
30. Kang, D.-S., Huang, Y.-S.: Multiple symmetric results for a class of biharmonic elliptic systems with critical homogeneous nonlinearity in \mathbb{R}^N . *Acta Math. Sci.* **37B**, 1665–1684 (2017)
31. Palais, R.: The principle of symmetric criticality. *Commun. Math. Phys.* **69**, 19–30 (1979)
32. Rellich, F.: *Perturbation Theory of Eigenvalue Problems*. Courant Institute of Mathematical Sciences, New York University, New York (1954)
33. D'Ambrosio, L., Jannelli, E.: Nonlinear critical problems for the biharmonic operator with Hardy potential. *Calc. Var. Partial Differ. Equ.* **54**, 365–396 (2015)
34. Lions, P.L.: The concentration-compactness principle in the calculus of variations, the limit case. *Rev. Mat. Iberoam.* **1** (part I), 145–201 (1985) 1 (part II) (1985) 45–121
35. Ambrosetti, A., Rabinowitz, P.H.: Dual variational methods in critical point theory and applications. *J. Funct. Anal.* **14**, 349–381 (1973)
36. Pucci, P., Rădulescu, V.: The impact of the mountain pass theory in nonlinear analysis: a mathematical survey. *Boll. Unione Mat. Ital.* (9) **3**, 543–582 (2010)
37. Rabinowitz, H.: *Methods in Critical Point Theory with Applications to Differential Equations*. CBMS. Amer. Math. Soc., Providence (1986)
38. Nyamoradi, N.: Solutions of the quasilinear elliptic systems with combined critical Sobolev–Hardy terms. *Ukr. Math. J.* **67**, 891–915 (2015)
39. Brezis, H., Lieb, E.: A relation between pointwise convergence of functions and convergence of functionals. *Proc. Am. Math. Soc.* **88**, 486–490 (1983)
40. Han, P.-G.: The effect of the domain topology on the number of positive solutions of some elliptic systems involving critical Sobolev exponents. *Houst. J. Math.* **32**, 1241–1257 (2006)
41. Alves, C.O.: Multiple positive solutions for equations involving critical exponent in \mathbb{R}^N . *Electron. J. Differ. Equ.* **1997**, 13 (1997)
42. Ekeland, I.: Nonconvex minimization problems. *Bull. Am. Math. Soc.* **3**, 443–474 (1979)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)